


THE
CHEMICAL CHANGES AND PRODUCTS
RESULTING FROM
FERMENTATIONS

UNIVERSITY OF TORONTO



3 1761 01349291 3


R. H. ADERS PLIMMER

v2 12



Digitized by the Internet Archive
in 2007 with funding from
Microsoft Corporation

THE
CHEMICAL CHANGES AND PRODUCTS
RESULTING FROM
FERMENTATIONS



MPhy
P

THE
CHEMICAL CHANGES AND PRODUCTS
RESULTING FROM
FERMENTATIONS

BY
R. H. ADERS PLIMMER, D.Sc. (LOND.)
GROCERS' RESEARCH STUDENT, JENNER INSTITUTE OF PREVENTIVE MEDICINE



LONGMANS, GREEN, AND CO.
39 PATERNOSTER ROW, LONDON
NEW YORK AND BOMBAY

1903

All rights reserved

99429
511109



CONTENTS

CHAPTER	PAGE
INTRODUCTION	1
I. CHANGES IN THE POLYSACCHARIDES	5
II. CHANGES IN THE POLYSACCHARIDES— <i>continued</i>	26
III. CHANGES IN THE TRISACCHARIDES AND DISACCHARIDES	31
IV. CHANGES IN THE MONOSACCHARIDES	40
V. CHANGES IN THE GLUCOSIDES	51
VI. CHANGES IN THE MONOSACCHARIDES— <i>continued</i>	61
VII. CHANGES RESULTING IN THE FORMATION OF OPTICALLY ACTIVE PRODUCTS	67
VIII. CHANGES IN ESTERS	71
IX. CHANGES IN UREA AND URIC ACID	75
X. CHANGES OCCURRING AS THE RESULT OF OXIDATION	78
XI. CHANGES OCCURRING AS THE RESULT OF OXIDATION— <i>continued</i>	82
XII. CHANGES OCCURRING AS THE RESULT OF REDUCTION	91
XIII. NITRIFICATION AND DENITRIFICATION	95
XIV. VARIOUS CHANGES OCCURRING AS THE RESULT OF FERMENTATIONS	99
XV. CHANGES IN BLOOD	105
XVI. CHANGES IN MILK, MUSCLE, AND IN THE LIQUID OF THE PROSTATE GLAND	110
XVII. CHANGES OCCURRING IN ALBUMINS AS THE RESULT OF THE ACTION OF PEPSIN	114

CHAPTER	PAGE
XVIII. CHANGES OCCURRING IN ALBUMINS AS THE RESULT OF THE ACTION OF TRYPSIN	122
XIX. PROTEOLYSIS BY FERMENTS OTHER THAN PEPSIN AND TRYPSIN	129
XX. THE CHANGES AND PRODUCTS OCCURRING AS THE RESULT OF PUTREFACTION	134
BIBLIOGRAPHY	143
INDEX OF AUTHORS	177
INDEX OF SUBJECTS	181

THE CHEMICAL CHANGES AND PRODUCTS RESULTING FROM FERMENTATIONS

INTRODUCTION.

THE majority of the chemical changes which are the result of fermentation occur in two large classes of compounds—the carbohydrates and the albumins. These are the materials used by young plants and animals as food-stuffs; as such, however, they cannot be assimilated, but must first undergo the changes which will be described in the following pages, in order that they may be made assimilable and really serviceable as food-stuffs.

The changes, so far as they are known, generally consist in a splitting up of a complex molecule into simpler ones, but, at present, only in a few cases can the stages in the transformation be followed. The reverse changes, or synthetic processes, no doubt, also take place both in plants and animals, and they consist in the building up of a complex molecule from simpler ones; at the present time, however, we have no knowledge as to how they are produced, but during the last few years experimental

synthetic fermentation processes with the ferments maltase (Hill, Emmerling), lipase (Kastle and Loevenhart, Hanriot), lactase, and emulsin (Fischer and Armstrong), have been carried out in the laboratory, and it is very possible that processes of this kind take place in nature on similar lines.

All these changes are brought about either directly or indirectly by means of living organisms. In the latter case the change is caused by a soluble ferment, or enzyme, which is secreted by the organism, and which can be employed in the laboratory as a reagent; similar changes to those which occur in nature can thus be produced, and the different products which result at the various stages of the process can be obtained. In the former case, so far as is at present known, these changes are produced by the living organism as the result of its metabolism, though, in all probability, an enzyme is secreted, not excreted, which produces these metabolic changes in the interior of their constituting cells. The researches of recent times upon this subject have made this supposition very probable; for example, a soluble ferment, or enzyme, has been extracted by Buchner from yeast, which can cause the alcoholic fermentation of saccharine liquids—a change which until then had been regarded solely as the result of the life of the yeast-cells.

The chemical changes which occur as results of fermentation are generally brought about by hydrolysis; but oxidation, as well as reduction, also takes place. All enzymes, as pointed out by Schönbein, are capable of decomposing hydrogen peroxide with the evolution of oxygen, and of producing a blue colouration with guaiacum tincture,

{ beer yeasts only being exceptions with regard to the latter test. It has lately been shown that the blue colouration produced in guaiacum tincture is due to the action of a separate enzyme, belonging to the class of oxydases, and Loew has still more recently made experiments showing that the power of decomposing hydrogen peroxide is due to the action of another separate enzyme of general occurrence, termed by him catalase.

The changes which the carbohydrates undergo are by far the most important, on account of the fact that the products obtained from them have been the substances upon which most investigation and research have been expended, and many of them are of commercial importance, *e.g.* the production of alcohol from starch. It will, therefore, be more advantageous to describe firstly the changes which these undergo.

The carbohydrates are generally divided into four classes: (1) the Polysaccharides; (2) the Trisaccharides, or Saccharotrioses; (3) the Disaccharides, or Saccharobioses; (4) the Monosaccharides, or Monoses.

To the monosaccharides, the glucosides, most of which are derivatives of glucose, are very closely allied, and the chemical changes which these undergo as the result of fermentation are best considered before those which take place in the monosaccharides, especially glucose, and which result in the formation of products other than alcohol, *e.g.* the production of lactic acid. Hydrolytic changes also occur in the fats and oils, and in urea, and these are best reviewed before the changes caused by oxidation, the most important of which is the production of acetic acid from alcohol; we shall then come to

reduction-changes and the processes of nitrification and denitrification, and finally to the changes occurring in the albumins, which are of a much more complicated nature, but no whit less interesting or important than those occurring in the carbohydrates.

CHAPTER I.

CHANGES IN THE POLYSACCHARIDES.

Of the polysaccharides, starch is by far the most important, not only on account of the great *rôle* which it plays in nature as a reserve material in plants, but also on account of the numerous investigations which have been carried out by various observers, in order to elucidate the chemical changes which it undergoes in plants as the result of the action of the ferment, or enzyme, known as diastase, upon it. These investigations have been made with the ultimate view of determining its constitution, as well as the constitution of the several decomposition products obtained from it during the process.

As is well known, a thin starch paste, containing about one or two per cent. of starch, to which a little diastase has been added, gradually becomes clear and the starch dissolves. This solution now is found to possess the power of reducing Fehling's solution, a property which it did not possess before the addition of the diastase, and, if it be tested at intervals, during the process of this transformation, with a solution of iodine in potassium iodide, a series of colour-changes will be observed: at first a blue colour is formed, showing still the presence of starch, but quickly this colour changes to purple, and then gradually to reddish-brown, until finally no colour-change

at all is seen. The purple colour is due to a mixture of starch and dextrin, the reddish-brown to a dextrin, and the cessation of colour-change indicates the presence of another dextrin and maltose. These changes, then, are the result of the hydrolysis of the starch through a series of dextrins to a reducing sugar, now known to be maltose.

Payen and Persoz, who, in 1833, first prepared diastase, were the first to examine the products resulting from this change in starch, and they found that a reducing sugar was not the only product obtained, but that a dextrin, not coloured by iodine and soluble in dilute alcohol, was also formed.

In 1860 Musculus observed that the dextrin and the sugar were formed simultaneously, and that the dextrin was not further altered. Independently of each other, about ten years later, Griessmayer, O'Sullivan, and Brücke ascertained that at least two dextrins were formed in this change, one of which was coloured brown by iodine, whilst the other was not visibly affected. Brücke named these respectively erythrodextrin and achroodextrin. O'Sullivan considered them to be identical, and said that they were convertible into maltose by malt extract.

Further investigations were made by Musculus and Gruber in 1878, who came to the conclusion that at least three achroodextrins were produced, the starch molecule breaking down by a series of hydrations and decompositions, maltose being obtained at each stage, together with a dextrin of decreasing, or lower, molecular weight.

In 1879 Brown and Heron published the results of the work which they had done upon the decomposition of starch by the action of diastase. Their conclusions, which

do not differ on the whole from the theories put forward by Musculus and Gruber in the previous year, are to the following effect: Starch has a formula of at least $10C_{12}H_{20}O_{10}$; in the hydrolysis by diastase, groups of $C_{12}H_{20}O_{10}$ are gradually split off, a dextrin of lower molecular weight being left at each stage, and eight of these dextrins are possible; the group of $C_{12}H_{20}O_{10}$, which is split off, is hydrolysed to maltose, $C_{12}H_{22}O_{11}$, the elements of a molecule of water being taken up.


In the same year O'Sullivan stated that four dextrins were formed, one of them being erythrodextrin and the other three achroodextrins.

About the same time Herzfeld brought forward still another view of the changes produced in starch, as the result of his own experiments. He observed that the products obtained were different when the hydrolysis was carried out below or above 65° C.; at the lower temperature maltose and achroodextrin were formed, and the latter was subsequently converted into a body, which he called maltodextrin, and maltose; at the higher temperature, in addition to these products, dextrin and erythrodextrin were formed. He concluded, therefore, that there was consecutively a conversion of the starch into soluble starch, erythrodextrin, and achroodextrin, and that the latter was then transformed into maltodextrin and maltose. For maltodextrin he suggested the composition of two dextrin groups united to a sugar group of the

formula $C_6H_{12}O_6$; thus: $\left\{ \begin{array}{c} C_6H_{10}O_5 \\ C_6H_{10}O_5 \\ C_6H_{12}O_6 \end{array} \right\}$.

In 1885 the results of a very thorough investigation

by Brown and Morris of the products obtained from starch by the action of diastase were published. These investigators found that 81 per cent. of maltose and 19 per cent. of dextrin were formed as end products, and that the dextrin was further hydrolysed only with great difficulty. They obtained also an intermediate product which they called maltodextrin, and this was found to be produced in varying amounts; this product was different to that of the same name obtained by Herzfeld, which they considered to be impure. The following theory was framed to account for these facts: Since only four-fifths of the starch is readily converted into maltose, the formula of starch is at least $5(\text{C}_{12}\text{H}_{20}\text{O}_{10})_3$, and consists of five amylin or dextrin-like groups, four of which are arranged symmetrically round the fifth, which is changed with greater difficulty, and finally remains as dextrin. At each stage a group of $(\text{C}_{12}\text{H}_{20}\text{O}_{10})_3$ is split off before complete hydration takes place, and a dextrin-like residue is left. The group which is split off is hydrolysed and forms maltodextrin



It is then further changed into maltose by the assimilation of two molecules of water.

That maltodextrin is a definite chemical compound, and not a mixture of maltose and dextrin, is shown by the following properties: (1) it cannot be separated into maltose and dextrin by dialysis; (2) it is completely converted into maltose by the action of diastase, whereas a mixture would leave a residue of dextrin; (3) it is not fermentable by yeast directly.


In 1874 Naegeli obtained, by the action of dilute acids upon starch, a substance which he called amylo-dextrin. This substance formed the subject of a paper by Brown and Morris in 1889; they found it to be a definite chemical compound, and not a mixture of maltose and dextrin, although its composition could be expressed in the terms of maltose and dextrin, and that it was analogous to maltodextrin in composition. They represented it by the formula $\left\{ \begin{array}{l} \text{C}_{12}\text{H}_{22}\text{O}_{11} \\ (\text{C}_{12}\text{H}_{20}\text{O}_{10})_6 \end{array} \right\}$, *i.e.* constituted as consisting of six amylin or dextrin groups, and one amylon or maltose group. It was converted, by the action of diastase, immediately and completely into maltose, and could not therefore contain the stable amylin group. It was also quite different from soluble starch, with which it was supposed by Naegeli and others to be identical.

In the same year the same investigators published their further views upon the constitution of starch, and the hydrolytic action of diastase upon it. The starch molecule was still supposed to consist of four complex amylin groups arranged symmetrically round a fifth similar group, which constituted a molecular nucleus; each of the five amylin groups had a formula of $(\text{C}_{12}\text{H}_{20}\text{O}_{10})_{12}$, and a molecular weight of 6480 (from results obtained by Raoult's method), which was considerably greater than they had originally supposed. The entire starch molecule, or, more correctly speaking, that of soluble starch, was represented by the formula $5(\text{C}_{12}\text{H}_{20}\text{O}_{10})_{20}$, with a molecular weight of 32,400. The first action of hydrolysis by diastase was to break up this

molecular complex, thereby liberating all the five amylin groups; four of these, by successive hydrolysis through maltodextrins, were completely converted into maltose, whilst the central amylin nucleus, by a closing up of the molecule, withstood hydrolysis, and constituted the stable dextrin which is so slowly acted upon. The four readily hydrolysable amylin groups were regarded as of equal value, which is in accordance with the view put forward by O'Sullivan in 1879, which stated that the dextrin molecules do not become smaller at each step, and that "the difference in their behaviour must be accounted for by a difference of relation in the arrangement of the molecules to one another, probably in solution alone." This hypothesis, they maintain, provided for intermediate maltodextrins or amyloextrins, whose number was only limited by the size of the original amylin group.

In 1890 Emil Fischer synthesised a new disaccharide from glucose, which he termed isomaltose; and Lintner, in 1891, who had been investigating the diastatic hydrolysis of starch, obtained a sugar of which he made the osazone, which corresponded in its properties with the osazone obtained by Fischer from his sugar: he therefore concluded that isomaltose was a product of the hydrolysis of starch, and not maltose, as had been previously supposed. Two years later, together with Düll, he criticised Brown and Morris's theory, and suggested that the various amyloins were mixtures of dextrin and isomaltose. At the same time, they brought forward their theory upon the action of diastase on starch, which is as follows: In the hydrolysis of starch five compounds are formed—isomaltose, maltose, and three dextrins, viz.

amylodextrin, erythrodextrin, and achroodextrin, of which they made molecular weight determinations. The first product is amylodextrin $(C_{12}H_{20}O_{10})_{54}$, which is the chief constituent of amidulin, or soluble starch; then erythrodextrin $(C_{12}H_{20}O_{10})_{17} \cdot C_{12}H_{22}O_{11}$ is formed; and subsequently achroodextrin $(C_{12}H_{20}O_{10})_5 \cdot C_{12}H_{22}O_{11}$, isomaltose, and maltose. They give the following equations to represent the formation of these substances:—



All these changes are supposed to occur simultaneously.

Schiffer also considered that isomaltose occurs amongst the transformation-products of starch. About the same time (1893), Scheibler and Mittelmeier evolved another theory, as they did not agree with that of Lintner and Düll. Scheibler and Mittelmeier thought that there was no real difference between the constitution of starch, dextrin, and the bioses, etc., except in the difference of the number of glucose residues in the molecule. The glucose molecules are capable of combining in two ways, either one or both of the carbonyl groups taking part; in the former case an aldehyde is formed, but not in the latter; thus there are monocarbonyl and dicarbonyl linkings. In starch the glucose residues are combined by

dicarbonyl linkings since starch cannot reduce Fehling's solution; hydrolysis by acids or by diastase takes place at the carbonyl linkings, two molecules, each containing an aldehyde group, being formed, and the particular glucose residue is connected with the remainder of the molecule by a monocarbonyl linking. These intermediate substances form dextrins, and as hydrolysis takes place probably simultaneously at several of the dicarbonyl linkings, a number of dextrins are simultaneously formed; by the further action of diastase, dextrins of smaller molecular weight, and finally maltose, are obtained.

Brown and Morris, in 1895, noticed that maltose, when liberated from starch by diastase, was in the state of half-rotation, and that on standing, or on heating, it acquired the normal rotatory power; O'Sullivan and Tompson have observed this same peculiarity with dextrose when obtained by the inversion of cane sugar.

In the same year, Ling and Baker, and Brown and Morris investigated Lintner's isomaltose, and concluded that the sugar obtained by Lintner was not identical with that obtained by E. Fischer, as it could be separated by fractionation with alcohol into maltose and dextrins of the maltodextrin type. Further, that maltose is the only sugar obtained in the hydrolysis of starch which can give a crystalline osazone.

In 1895 Ulrich repeated Lintner's work, and found that maltose and not isomaltose was formed; Jalowetz obtained the same result as Ulrich, and considered that maltose was the only sugar obtained. Ost, in 1896, again confirmed the presence of maltose, and Pottevin, in 1899, also came to the same conclusion. These various

observations definitely prove that maltose is the sugar resulting from the hydrolysis of starch by the action of diastase.

Mittelmeier, in 1895, promulgated a still different view upon the diastatic resolution of starch. According to him there are two stages: in the first stage, a small portion of the starch is rapidly converted into amylo-, achroo-, and erythro- dextrins, and sugar; these are not identical with those formed later; and he calls them, according to the stage, primary and secondary dextrins. From the primary dextrins maltose, and from the secondary a new sugar, which he terms metamaltose, are obtained. This view he modified two years later (1897), thus: the first action of diastase on starch results in the formation of equal quantities of two different amylodextrins, one of which is more easily attacked by diastase than the other; thus, while the one is in the erythro-dextrin stage, the other has passed from dextrin to sugar; the existence of a sugar molecule and a dextrin molecule in one molecular complex was thus said to have been made intelligible.


In 1898 Pottevin described a number of experiments which tended to show (1) that the conversion of starch into maltose took place in two stages, dextrin being the intermediate product; (2) that the various dextrins differed only in their physical states; (3) that the different parts of the starch granule differed in the ease with which they were converted into dextrin, and the dextrin then into maltose. Thus, the conversion of starch proceeds with unequal rapidity in the different portions of the mass. In the next year he stated that

the dextrins could be divided into three groups: (1) achroodextrins not coloured; (2) amylo-dextrins coloured blue; (3) erythro-dextrins, coloured red, by iodine. Amylo-dextrin, he considered, was not a definite substance, as it gave α - and β - amylo-dextrins; the formation of dextrin was independent and distinct from the formation of maltose from dextrin. Brown and Morris's maltodextrin he considered to be a mixture, as he had separated it into maltose and dextrin.


In 1899 Brown and Millar published two very important papers upon maltodextrin and dextrin. To maltodextrin the empirical formula $\left\{ \begin{array}{l} \text{C}_{12}\text{H}_{22}\text{O}_{11} \\ (\text{C}_{12}\text{H}_{20}\text{O}_{10})_2 \end{array} \right\}$ was assigned by Brown and Morris in 1885.

When this was prepared in a pure state—a very laborious process—and oxidised with mercuric oxide and barium hydroxide until it no longer had any reducing power, a definite and complex carboxylic acid, called maltodextrinic acid A, was obtained. By the hydrolytic action of diastase 40 per cent. of maltose and 60 per cent. of another carboxylic acid—maltodextrinic acid B—were formed; dilute acids gave 85·8 per cent. of d-glucose and a simpler acid containing five carbon atoms, which is a quite different result. The B acid gave the same products, on hydrolysis, as the A acid. The five carbon atom acid appeared to be a normal acid derived from a pentose, and had properties in common with those of xylonic acid. From these results they conceived an idea of the constitution of maltodextrin and the derived acids which is as follows: maltodextrin is made up of a number of C_{12} - or maltose residues, since it is completely


hydrolysable to maltose, but still retains a terminal CHO group owing to its possessing reducing properties; oxidation goes further than the conversion of this CHO group into a COOH group, since it splits up the terminal C_{12} -group into a C_5 - acid residue, which remains attached to the hydrolysable portion of the polysaccharide. If now Fischer's constitutional formula for maltose—



be accepted, the polysaccharide maltodextrin may be regarded as made up of three such molecules, with the elimination of two molecules of water, and may be represented as follows :—



or abbreviated thus—




the sign $\cdots$ being used to denote the open terminal carbonyl group, as suggested by Scheibler and Mittelmeier.

The C_{12} - groups and the C_6 - groups are linked to one another through oxygen atoms, and at these points the molecule breaks down in hydrolysis.

The difference between acid and diastatic hydrolysis, in that glucose is only formed by the former, may be accounted for by spatial relations between the adjacent C_{12} - groups, which may determine the differential action of diastase, or by the relative size or mass of the C_6 - or C_{12} - groups. This resembles the selective action of maltase or glucase, and emulsin, which was shown by E. Fischer to exist in their behaviour to glucosides.


The constitution of maltodextrinic acid A can only be explained by assuming that the terminal C_{12} - group of the maltodextrin is attacked in such a way that by simultaneous or successive oxidation $\text{C}_7\text{H}_{12}\text{O}_5$ is removed, and $\text{C}_5\text{H}_{10}\text{O}_6$ is left still in combination with the polysaccharide residue. The following is thus the only formula which it can possess :—




the reducing C_{12} - group of maltodextrin having become $-O-CH_2-(CHOH)_3-COOH-$ the residue of a normal C_5 - carboxylic acid.

The hydrolytic decompositions of maltodextrinic acid A can be expressed by the equations :—

1. *Diastatic*—




2. *Acids*—




The dextrin, which is formed together with maltose in the diastatic hydrolysis of starch, was obtained in the purest possible state by fractionation with alcohol and occasional fermentation, and was found to possess a feeble reducing power. On oxidation with mercuric oxide and baryta, it gave a complex carboxylic acid, called dextrinic acid. By hydrolysis with acids it was resolved into d-glucose and a C_5 - acid identical with that obtained from maltodextrin, but diastatic hydrolysis gave maltose and glucose, which are also obtained in almost equal quantities from dextrin itself by the action of diastase. Here there is no selective action of diastase to split off maltose rather than glucose, and the difference in linkings or spatial

relations between the C_{12} -groups and their constituent C_6 -groups cannot account for these results. The stable dextrin must therefore be made up of C_6 - complexes, or as a combination of 39 groups of $C_6H_{10}O_5$ and one $C_6H_{12}O_6$ group, or as $40(C_6H_{10}O_5) + H_2O$.

Its constitution may be regarded as a condensation of 40 glucose molecules with the elimination of 39 molecules of water, expressed by the formula :—





The constitution of dextrinic acid will then be given by converting the right-hand group into—




the residue of a C_5 - acid given by complete hydrolysis.

These formulæ can be abbreviated thus :—




The size and constitution of the starch molecule are suggested from the total results which Brown has obtained with his various co-workers. The stable dextrin obtained with maltose cannot have a simpler formula than $40 \text{ C}_6\text{H}_{10}\text{O}_5 + \text{H}_2\text{O}$, and the equation representing its formation from starch together with maltose in the proportion of 1 to 4 becomes :—




The starch molecule cannot be less than five times that of stable dextrin, on account of this result, and dextrin from molecular weight determinations has a formula of $39 (\text{C}_6\text{H}_{10}\text{O}_5) \cdot \text{C}_6\text{H}_{12}\text{O}_6$, giving a molecular weight of 6498.

Its molecular weight cannot be less than 32,400, and its empirical formula can be represented by :—



Starch also does not possess a carbonyl group, as it is a non-reducing polysaccharide, and its constitution can be expressed most simply as made up of 80 maltose groups and 40 dextrin groups, linked in ring form through oxygen atoms, thus :—



On hydrolysis with diastase, the dextrin complex A is split off, forming stable dextrin, $39 C_6H_{10}O_5 \cdot C_6H_{12}O_6$, whilst the maltose portion B of the ring is attacked at the oxygen linkings of the C_{12} - groups, forming, by successive stages of hydrolysis, maltodextrins or reducing dextrins, and finally maltose.


Also in 1899 W. Syniewski obtained from starch by the action of water, alkalies, and sodium peroxide, a number of substances which he termed amyloextrins, of which

the simplest was amylogen. Starch and the amylo-dextrins are therefore considered to be made up of a number of these amylogen molecules associated together in the form of anhydrides, by loss of water. The amylogen obtained by the action of sodium peroxide has the formula $C_{54}H_{96}O_{48}$, from molecular weight determinations, and consists of three maltose residues combined with a dextrin residue of eighteen carbon atoms; and the dextrin consists of three glucose residues, two of which are combined together in the form of isomaltose.


Diastase hydrolysis was performed with the amylo-dextrin obtained by the action of boiling water under pressure, and the process is supposed to be a gradual splitting off of maltose groups leaving a maltodextrin, until a dextrin remains which does not contain maltose, and is called "Grenzdextrin," a glucodextrin. This is then converted into simpler glucodextrins and glucose, then into isomaltose and glucose, and finally into glucose. The final result, which is a mixture of dextrin and maltose, is explained by supposing that the maltose is not so easily removed from some of the intermediate dextrins as it is from others.

By the action of malt extract, previously heated to 78° , upon amylo-dextrin, Syniewski has recently obtained a substance, which he has termed Grenzdextrin II.; it differs from Grenzdextrin I., which he formerly obtained by the action of fresh unheated malt extract, and is identical with Brown and Morris's maltodextrin. To account for these results he supposes that the amylogen residue contains three kinds of carbonyl linkings; he denotes them by α , β , γ , and the glucose residue by (C_6) ;


and the following schematic formula is given to the amylogen residue:—




When fresh malt extract acts upon the amylogen residues of amylopectin, the α linkings are attacked, maltose $(\text{C}_6) - \gamma - (\text{C}_6)$ and Grenz-dextrin I., made up of the residues—





being obtained; when heated malt extract acts upon the amylogen residues, the β linkings are attacked, and Grenz-dextrin II., made up of the residues—



is obtained. He has previously shown that Grenz-dextrin I. has the formula $\text{C}_{72}\text{H}_{124}\text{O}_{62}$, and consequently it contains four of these residues; amylogen, which has the formula $\text{C}_{54}\text{H}_{96}\text{O}_{48}$, can only give one of these residues, so that starch must contain at least four amylogen residues; if (C_6) and (C_{12}) represent glucose and maltose residues respectively, the following formulæ are then obtained:—



or



where the single lines between the (C_{18}) -residues represent β -carbonyl linkings, and the double lines two carbinol linkings.

H. Brown's last publication, with Glendinning, is upon the velocity of diastase hydrolysis, and he finds that it does not follow the law of mass action. A. J. Brown has obtained a similar result in the case of invertin.

Before passing to the next polysaccharide, the other sugars which have been said by different investigators to be products of diastase hydrolysis might be enumerated. Some observers have supposed glucose to be a product, but the conditions of the experiment were probably the cause of this observation, other enzymes being present in the diastase solution. Nasse has obtained a sugar which he calls ptyalose, differing from both glucose and maltose, by the action of ptyalin upon starch.

The action of diastase upon starch is considered by Arthur Meyer as catalytic, the amylose taking up the elements of water and breaking up into two or more molecules of amyloextrin, which subsequently forms

dextrin and isomaltose; maltose is formed by intramolecular rearrangement from isomaltose as well as from dextrin.

From the foregoing it will be seen that very many theories have been advanced by many workers to explain the hydrolysis of starch by diastase. Two facts, however, stand out clearly from the mass of complicated results: firstly, the production of maltose; and secondly, the production of a dextrin much more difficult to convert into maltose, but eventually convertible. Intermediate between the starch and maltose many dextrins, no doubt, are formed, and they have been named with the various terms enumerated above. Of these, Brown's maltodextrin only has been prepared in anything like a state of purity; but this may possibly be the simplest of them, as its constitution has been determined: the other dextrins are no doubt mixtures of several of the intermediate ones, and have not yet been prepared in such a state of purity that their properties could be established. Opposite is a table showing schematically the various results which have been obtained.


Name.	Date.	Products, according to the various Authors.
Payen and Persoz	1833	Dextrin and reducing sugar.
Museulus	1860	Ditto
Griessmayer	1871-72	Two dextrans : one coloured brown, one not coloured, by iodine. Sugar.
Brücke	Ditto	Ditto : erythro-dextrin, achroo-dextrin.
O'Sullivan	Ditto	Ditto ; but identical $\rightarrow$ maltose.
Museulus and Gruber	1878	Erythro-dextrin, and 3 achroo-dextrans. Maltose at each stage.
Brown and Heron	1879	8 possible dextrans. Maltose at each stage.
O'Sullivan	Ditto	1 erythro-dextrin and 3 achroo-dextrans. Maltose.
Herzfeld	Ditto	Starch $\rightarrow$ soluble starch $\rightarrow$ erythro-dextrin $\rightarrow$ achroo-dextrin $\rightarrow$ maltodextrin + maltose.
Brown and Morris	1885	Dextrin $\frac{2}{3}$ and maltodextrin $\rightarrow$ maltose $\frac{1}{3}$.
Ditto	1889	Same results as above, but starch molecule larger than originally supposed.
Lintner and Düll	1893	Starch $\rightarrow$ amylo-dextrin $\rightarrow$ 3 erythro-dextrin $\rightarrow$ 9 achroo-dextrin $\rightarrow$ 54 iso-maltose $\rightarrow$ 54 maltose.
Scheibler and Mittelmeier	Ditto	2 dextrans $\rightarrow$ dextrans of smaller molecular weight $\rightarrow$ maltose.
Mittelmeier	1895	(1) Primary dextrans (amylo-, erythro-, achroo-) $\rightarrow$ maltose.
		(2) Secondary dextrans $\rightarrow$ metamaltose.
Ditto	1897	2 amylo-dextrans $\rightleftharpoons$ one quickly to sugar.
Pottevin	1898	Starch $\rightarrow$ dextrans $\rightarrow$ maltose.
Ditto	1899	Dextrans = α and β amylo-, erythro-, achroo-dextrans.
Brown and Millar	1899	Dextrin and maltodextrin $\rightarrow$ maltose.
Syniewski	1899	$\downarrow$ Maltose, with difficulty. Starch $\rightarrow$ amylo-dextrans (simplest amylogen) $\rightarrow$ maltodextrans + maltose. $\downarrow$ Glucose + isomaltose $\leftarrow$ simpler glucodextrans $\leftarrow$ grenzdextrin + maltose. $\downarrow$ glucose.

CHAPTER II.

CHANGES IN THE POLYSACCHARIDES.—Continued.

THE next polysaccharide, inulin, which is also found as a reserve material in certain plants, undergoes hydrolysis by the action of the ferment known as inulase, discovered by Reynolds Green in 1887. The formula $C_{12}H_{20}O_{10} \cdot 2H_2O$ is usually given to this polysaccharide, which, according to Bourquelot, is converted by fermentation into lævulose or fructose.

Kiliani, from results of combustion, gave the formula $C_{36}H_{62}O_{31}$ or $6C_6H_{10}O_5 + H_2O$ to this carbohydrate; but Brown and Morris, in 1889, from determinations made by Raoult's method, considered the formula to be $2C_{36}H_{62}O_{31}$. They also stated that inulin resembled amyloextrin in its properties very closely, and they gave the following formula to express its constitution:—



which is closely analogous to that of amyloextrin. The ultimate products of hydrolysis of inulin are different, and they therefore thought that the amylin and amylo groups in its molecule were different to those in amyloextrin, because lævulose is obtained, whereas in the latter case maltose is obtained. Also, that it is a product

of the hydrolysis of an unknown carbohydrate, and stands in the same relation to this body as amyloextrin does to starch.

According to Düll, the molecular weight of inulin is 2950, which agrees with the formula $(C_6H_{10}O_5)_{18}$ or $(C_6H_{10}O_5)_{18} + H_2O$; this observer has found that the lævulins obtained from inulin by acid hydrolysis are really reversion products of lævulose, and therefore they are not hydrolytic decomposition products.

In animals, carbohydrates are stored up in the form of glycogen; this reserve material is found in greatest amount in the liver, where it is converted into maltose and glucose when required for use by the animal.

The chemical changes which this body undergoes by the action of diastase have been investigated by Miss Tebb, who has compared this hydrolysis with that produced by acids. The former method gives achroodextrin and another dextrin named dystropodextrin, whilst dilute acids give soluble glycogen, erythrodextrin, and achroodextrin. The sugar formed is supposed to be maltose, from which, by the action of glucase, glucose is produced in the liver. Miss Tebb has identified her dystropodextrin with the product of the same name obtained from glycogen by Seegen.

As regards the molecular weight of glycogen, the most recent determinations have been made by Jackson, who used carefully purified glycogen, and found that the amount of lowering of the freezing-point corresponded to a molecular weight of 9500–10,000.

Caroubin, a polysaccharide of the formula $C_6H_{10}O_5$, has been obtained by Effront from the seeds of the carob-bean

(*Ceratonia siliqua*); it is hydrolysed by the enzyme, caroubinase, obtained from the germinating seeds, to a non-crystallisable reducing sugar, which is a hexose. This hexose is fermentable with yeast, and gives an osazone melting at 183°C ; its reducing power is the same as that of glucose, but its specific rotatory power of $[\alpha]_D = +24^{\circ}$ is different. Effront called this sugar caroubinose, but Marlière considers it to be a mixture of glucose, fructose, and galactose; Von Ekenstein has since identified it as d-mannose.

Bourquelot and Hérissé have found that the so-called horn-albumin, which occurs in the fruit of the carob-tree in the species of *Medicago sativa* and *Trigonella fœnum græcum*, is a manno-galactan; this is hydrolysed by the enzyme seminase, which can be extracted from the germinating seeds, and converted into mannose and galactose.

The albumin of the seeds of *Phoenix canariensis* contains mannans; these are decomposed by a soluble ferment present in the germinating seeds into mannose (Bourquelot and Hérissé). Mannose, according to Hérissé, is also the product of the mannan of orchid tubers when they are acted upon by a soluble ferment present in them.

The cell-walls of plants consist of cellulose and certain bodies closely allied to, and found together with it, termed pectoses. The changes which these undergo when acted upon by the enzyme cytase, are probably very complicated, like those of starch; the ultimate products being a reducing sugar from cellulose and metapectic acid from the pectose; metapectic acid is probably identical with

arabic acid, which is found in gum arabic, and which can be hydrolysed to galactose and arabinose.

Cross and Bevan, who have investigated the hydrolytic decomposition products of the various celluloses, state that glucose and mannose are formed from oxycelluloses; in addition to these two sugars, galactose, arabinose, and xylose are formed from pectocelluloses. Tollens and Ganz state that they have obtained a sugar—seminose—from cellulose by acid hydrolysis; and Reiss has also found the same carbohydrate.

According to Omeliansky, cellulose is fermented by *Bacillus fermentationis cellulosa*, with the result that 70 per cent. fatty acids, chiefly acetic and butyric acids, together with 30 per cent. of gases (carbon dioxide, hydrogen, and in the last phases methane), are formed.

Besides cellulose, lignin, the woody substance, is contained in the cell-walls of plants. Czapek has observed that this is dissolved by an enzyme, which he has termed hadromase. The woody substance is regarded as an ester—a compound of cellulose and hadromal—and it is decomposed by the soluble ferment into its constituents. Hadromal has the properties of a phenol and of an aldehyde, and is probably a para-compound. The cellulose formed in this decomposition is then acted upon by cytase, which is also secreted by the mould fungi from which the hadromase was obtained.

The transformation, which many fruit juices undergo, into a jelly, is, according to Frémy, due to an enzyme called pectase. The substance so changed is the pectose which is found with the cellulose. Frémy states that pectine is formed from pectose by the action of acids, and that

the conversion by the enzyme takes place in two stages, resulting in the successive formation of pectosic and pectic acids, which are gelatinous bodies.

The most recent investigations have been carried out by Bertrand and Mallèvre, who have found that calcium is contained in the jelly, which consists of calcium pectate, and not of free pectic acid, and that gelatinisation depends upon the presence of calcium; this can, however, be replaced by barium or strontium. Free mineral acids, and even malic, citric, or oxalic acids, hinder the formation of this coagulum, but the effect of acids is decreased by excess of pectase or calcium salts. Coagulation, therefore, depends on the proportion of pectase, calcium salts, and free acid.


The power of gelatinising possessed by pectine under the influence of pectase, is, according to Bourquelot and Hérissé, destroyed by the action of the soluble ferment pectinase, which converts it into a reducing sugar. This change can take place even when the pectine has been coagulated, but coagulation by pectase cannot occur after the decomposition produced by pectinase.

CHAPTER III.


CHANGES IN THE TRISACCHARIDES AND DISACCHARIDES.

CHEMICAL changes as the result of fermentation are produced in the trisaccharides raffinose, melizitose, and gentianose.

Raffinose, melitose, or melitriose, $C_{18}H_{32}O_{16}$, was discovered in 1876 by Loiseau, and identified with melitose by Tollens. It is hydrolysed by an extract of yeast or of *Aspergillus niger* (Bourquelot) into fructose, glucose, and galactose, thus—




This change occurs in reality in two stages, in the first of which fructose is split off, and melibiose, a disaccharide, is left; this substance is then hydrolysed into glucose and galactose, and the following two equations represent these changes:—



This hydrolysis was formerly attributed to the action of the enzyme invertin, but Fischer and Niebel state that an extract of the small intestines of the horse or dog, which contain invertin, cannot produce this conversion.


The enzyme which produces this change has been termed raffinase, and it occurs in bottom fermentation yeasts. Fischer and Lindner have extracted an enzyme from these, which hydrolyses melibiose, as also has Bau, who attributes the first stage in the conversion to invertin, and the second stage to this enzyme, which he terms melibiase. Scheibler states that these two changes can also be produced by very dilute acids at a low temperature.

Melzitose, discovered by Berthelot in 1859, in Brançon manna, is resolved, according to Bourquelot, by an extract of *Aspergillus niger* into glucose and a disaccharide, touranose :—




Touranose is isomeric with maltose; its products of hydrolysis by dilute acids being two molecules of glucose. The enzyme termed melizitase, which produces the change, apparently is unable to produce this latter conversion.

Gentianose, a trisaccharide extracted from the fresh gentian root by Bourquelot and Hérissé, is converted by invertin into fructose, and a new disaccharide, gentiobiose :—



An extract of *Aspergillus niger* is capable of completely hydrolysing this trisaccharide into one molecule of fructose and two molecules of glucose :—




The changes which the disaccharides undergo as the result of fermentation have been very thoroughly

investigated; they are hydrolytic in their character, one molecule of water being always taken up.

Cane sugar is the oldest known representative of this group of compounds, on account of the fact that the products obtained from it subsequently undergo alcoholic fermentation. The change which it undergoes in the process of fermentation was shown by Berthelot, in 1860, to be due to the action of the enzyme invertin. This ferment converts cane sugar into d-glucose and d-fructose, and the change can be demonstrated in two ways: (1) the solution before the action of the ferment is dextro-rotatory, but it becomes lævo-rotatory when the change is completed; (2) it is originally unable to reduce Fehling's solution, but afterwards it becomes capable of doing so. The products formed, d-glucose and d-fructose, are both capable of reducing cupric oxide in solution; the lævo-rotation of the solution is due to the fact that d-fructose has a stronger rotatory power to the left than d-glucose to the right.

The equation which represents this change is:—



The rate of this change is at first very rapid, but as the hydrolysis proceeds, it gradually diminishes; according to O'Sullivan and Tompson, this change follows Harcourt's rule, *i.e.* the conditions remaining the same, the rate of change is proportional to the amount of the substance undergoing the change, and is represented by the equation:—

$$\frac{1}{\theta} \log \frac{1}{1-x} = C$$

where C is a constant, θ the time in minutes, and x the quantity changed in unit time.

The rate of the change is influenced by the reaction of the solution, the temperature (optimum at 55°), and the concentration of the cane sugar, 20 per cent. being the most favourable strength. O'Sullivan and Thompson state that the products of the reaction exert no influence, but according to Müller they are inhibitory. Quite recently A. J. Brown has taken up this subject, and he finds that the change does not follow the law of mass action, and that the products have an inhibitory effect. He considers that time is a factor in this change, on account of molecular combination of the enzyme with the cane sugar, the combination having a resistance for a definite period of time before final disruption and change take place. Molecular combination between the invertin and the cane sugar was shown to be very probable, since a solution of invertin and cane sugar can withstand a higher temperature than a solution of invertin alone (O'Sullivan and Thompson).


Henri has also found that the products of the hydrolysis of cane sugar lessen the rapidity of its inversion by invertin, and that if more cane sugar be added during the process, acceleration is produced.

Brown and Pickering, in 1897, worked upon the thermo-chemistry of carbohydrates; both in the hydrolysis of cane sugar by invertin, and of starch by diastase, it was found that heat was evolved.


According to Scharinger, cane sugar is decomposed by a mucus-forming bacillus, with the result that hydrogen, alcohol, acetic, succinic, and lactic acids are formed, the

last acid being the principal product. A carbohydrate of the composition $C_6H_{10}O_5$ was isolated from the mucus formed; this is a galactan, and yields on oxidation oxalic and mucic acids.


Maltose, which has been seen to be a product of the diastatic hydrolysis of starch, becomes changed, under the influence of the ferment maltase or glucase, into two molecules of glucose :—



This hydrolysis has been investigated lately by Croft Hill, whose results are very important; he found that when maltase acts on maltose, the hydrolysis proceeds up to a certain point and then ceases, and that this point depends on the temperature, and on the concentration of the maltose. If the glucose be now removed from the solution, further hydrolysis of the remaining maltose takes place until the same point is again reached. If, however, at this point more glucose be added, a re-conversion into maltose takes place. It appears, therefore, as if a state of equilibrium were set up during the process, just as in the case of esterification, which can be represented by the equation :—



That for maltose would be :—




From these experiments it appears that the enzyme maltase can exert a synthetical action as well as an analytical.

Emmerling has lately confirmed these results of Croft Hill's, as regards the synthetic action of maltase, but he states that isomaltose, not maltose, is the synthetical product. Hill has denied this, saying that the conditions of the experiments were different in the two cases, and that the osazone of Emmerling's was probably impure, and, as has already been shown, it is very difficult to purify. Emmerling has replied, saying that isomaltose was the sugar obtained in his experiments.

Croft Hill has since notified another case of reversed, or synthetical, ferment action. A 60 per cent. solution of glucose, when treated with taka-diastrase containing maltase, was changed into a solution containing 58 per cent. of glucose and 2 per cent. of maltose hydrate; he had observed, before making this experiment, that equilibrium was obtained in a solution of glucose and maltose in the presence of taka-diastrase and maltase, when the solution contained 2 per cent. of maltose: he thinks that other polysaccharides are also formed under the same conditions. Using pancreatic ferment and dextrose, he has also found that a reversible reaction occurs. Similar results of synthetic action by ferments have been lately observed in the case of lipase, lactase, and emulsin.


Lactose, another member of this group, has also been known for a long time. It is owing to its presence in milk that the transformation known as lactic acid fermentation is due. This change takes place in two stages; in the first stage a molecule of water is taken up under the action of the enzyme called lactase, and two sugars, glucose and galactose, are the products:—




These sugars can then undergo the second stage in the transformation into lactic acid.

Bourquelot has stated that lactose can be converted into lactic acid without previous hydrolysis into glucose and galactose, but von Freudenreich has shown that the lactic acid bacteria can produce these two sugars, which are then converted further into lactic acid.

The disaccharides lactose and maltose are acted upon by the enzymes maltase and emulsin in a way similar to the α - and β -glucosides. They can therefore be regarded as glucosides: lactose as the β -galactoside of glucose, and maltose as the α -glucoside of glucose. The following formulæ demonstrate the above facts:—




Trehalose, or mycose, a disaccharide occurring in Syrian manna, is decomposed by trehalase into two molecules of glucose:—




The enzyme trehalase, which produces this decomposition, was discovered by Bourquelot in *Aspergillus niger* and other moulds.

Melibiose, obtained from melitriose as already mentioned, is converted, according to Fischer and Lindner, and Bau, by a bottom fermentation yeast extract, but not by a top fermentation yeast extract, which contains the enzyme melibiase, into glucose and galactose:—



It is also hydrolysed by lactase (Fischer and Armstrong).

Gentiobiose was lately obtained from gentianose by Bourquelot and Hérissé, but not in a crystalline form; according to these observers, it is a definite compound resembling maltose, and is hydrolysed by an extract of *Aspergillus niger* into two molecules of glucose:—



The osones of the disaccharides maltose and melibiose have quite recently been shown by Fischer and Armstrong to be capable of undergoing hydrolysis by ferments in the same way as these disaccharides themselves, with the formation of the same products; these compounds can thus be used in the place of their disaccharides for testing their behaviour with regard to the action of ferments.

These investigators have also synthesised three new disaccharides, which have been called glucosidogalactose, galactosidoglucose, and galactosidogalactose, since they think that their constitution is very similar to that of the glucosides. These three disaccharides are decomposed by emulsin into their constituents, but only two of them—galactosidoglucose and glucosidogalactose—are fermented by bottom yeasts; top yeasts have no action on them at all. Galactosidoglucose in properties very closely resembles melibiose, and it is very probable that these two disaccharides are identical.

Fischer and Armstrong have further obtained a fourth disaccharide, isolactose, by the action of lactase upon a mixture of glucose and galactose. This is another case of synthetical ferment action. As regards its behaviour to ferments, this disaccharide occupies a position between lactose and melibiose. Another disaccharide was formed when lactase acted upon a solution of glucose alone, the synthesising power of this enzyme being here again shown.

These investigators have shown that emulsin also is capable of exerting a synthetic action, since they obtained a disaccharide as the result of its action upon a mixture of glucose and galactase, as in the case of lactase.

CHAPTER IV.

CHANGES IN THE MONOSACCHARIDES.

THE members of the first group of monosaccharides undergo various changes as the result of fermentation. The oldest known change is that called alcoholic fermentation, which glucose especially undergoes, together with fructose, both of which are formed, as before stated, by fermentation from cane sugar. The history of this fermentation process is very interesting, as the whole subject begins with it.


From the very earliest times it has been known that sweet liquids, if exposed to the air and allowed to stand, become cloudy and turbid, and bubbles of gas rise to the surface, which give the appearance of boiling to the liquid. The term "fermentation" was first applied to this change, on account of this appearance of boiling, but it was soon also applied to all changes where any effervescence took place, especially to the process of putrefaction, the terms *putrefactio* and *fermentatio* being used synonymously. A froth forms on the liquid, and a sediment, or "lees," in it; later, the whole settles to the bottom, leaving the liquid clear. After the process of apparent boiling is over, the liquid will no longer be sweet, but will have a spirituous smell and taste, owing to the presence of the

substance now known as alcohol. The earliest idea of the nature of this process is found in the writings of Basil Valentine, about the end of the fifteenth century, and was to the following effect: The scum, which is formed in the process, imparted an internal inflammation to the liquid, so that a purification was thereby determined; from this it is seen that he supposed that the alcohol was present in the liquid before the fermentation began. The gas evolved, which gives the boiling appearance, was thought to be the same as ordinary air; but van Helmont, in the following century, pointed out that this gas resembled that sometimes found in caves and wells, and he termed it "gas sylvestre." In 1682 Becher observed that this change occurred only in saccharine liquids, and he considered the alcohol to be a product of the reaction. He thought that the presence of air was necessary to start the process, which, in his opinion, was analogous to combustion; a distinction was also made between fermentation and putrefaction, and between true alcoholic fermentation and acid fermentation, *e.g.* the production of lactic acid. Some twenty years before this observation, Willis had put forward a new view of this change. It had already been found that the scum was necessary for the process, and this scum had been called a ferment; this ferment was supposed to be possessed of a motion peculiar to itself, which it transmitted to the fermentable matter. Stahl, a little later, also held this view, but considered, in addition, that the fermentable matter consisted of an unstable union of salt, oil, and earth; by the internal motion caused by the ferment, these particles were separated from one another, only to combine together

again to form more stable molecules, consisting of the same principles, but in other proportions.

Another century elapsed before any real light was cast upon the nature of this process; then Lavoisier took up the investigation of the changes, which was the first work of a really scientific character upon the subject. Instead of using the juices of fruits which had been employed before, he worked with a solution of cane sugar. This substance he regarded as a vegetable oxide made up of two bases, carbon and hydrogen, brought into the state of an oxide by a certain proportion of oxygen; these three elements were in such a feeble state of equilibrium that only a slight force was necessary to destroy their connection. He made analyses of the sugar, the alcohol, and the carbon dioxide, the results of which are to be found in his "Elements of Chemistry." The change is explained as a mere separation of the elements into two portions, the one being oxygenated at the expense of the other, in order to form carbon dioxide, whilst the other is deoxygenated and converted into the combustible substance called alcohol. The conclusion is that, if it were possible to reunite the alcohol and the carbon dioxide, sugar would be again formed. Though his analyses were far from correct, yet he approached very nearly to the truth.

Twenty years later, Gay Lussac and Thénard, and De Saussure, made more accurate analyses of cane sugar, which supported Lavoisier's view. The formula $C_{12}H_{12}O_{12}$ was given to sugar, and the following equation expressed the change:—



It was pointed out in 1828, by Dumas and Boullay,


that Gay Lussac's results were not quite accurate; this was acknowledged by Gay Lussac, who stated that he modified them to agree with the equation. If, however, it was considered that a molecule of water took part in the process, the results would be accurate, and so Dumas and Boullay rewrote Gay Lussac's equation thus:—



In 1832 Dubrunfaut discovered that cane sugar, before it could undergo fermentation, was transformed into an uncrystallisable sugar. Berthelot showed that this change is produced by a soluble ferment which is present in yeast, and that alcoholic fermentation takes place only after the transformation; he also stated that the above equation is experimentally realised.

In 1847 Schmidt discovered that succinic acid was also a product of this fermentation; but this fact was scarcely noted at the time, and became practically forgotten.

The most important work upon the subject is that of Pasteur, who, about 1860, published the results of his early work on the question, and showed that succinic acid and glycerine, besides alcohol and carbon dioxide, are products of this fermentation; that the glycerine and succinic acid are formed at the expense of the sugar; and that a portion of the sugar is used up by the ferment: the amounts of succinic acid and glycerine produced varied slightly, according to the temperature, the acidity, and the rate of the fermentation. He gave the following equation to represent the formation of succinic acid, glycerine, and carbon dioxide:—



but considered it to be only a very approximate expression of the numerical results which he obtained by analysis.

Acetic acid is another product of alcoholic fermentation. It was first noticed by Béchamp, and attributed by Pasteur to a subsequent fermentation; its production will be considered later.

Higher alcohols, collectively known as fusel oil, are formed also in alcoholic fermentation, but the process of their formation is not yet known. This fusel oil consists of a mixture of propyl, isobutyl, amyl, caproyl, cœnanthyl, and capryl alcohols in various proportions; amyl alcohol occurs sometimes in somewhat large quantities.

It is upon the exciting cause of fermentation that the most varied and interesting theories have been held. In 1680 Lœwenhoek examined the scum on fermenting liquids with his microscope, and found that it consisted of small spherical or ovoid globules. Fabroni, in 1787, considered these particles to be what he called a "vegeto-animal" substance; and Thénard, three years later, thought that the substance was of an animal nature from its nitrogenous chemical composition, but as to its organisation, he made no allusion. Astier, in 1813, still considered these particles to be of animal origin, and thought that they could only live at the expense of the sugar. Desmazières, in 1826, stated the same view. In 1810 Gay Lussac concluded from his experiments that oxygen was necessary to start fermentation, but was not required for its continuation; in his "*Mémoire sur la Fermentation*" he alludes to Appert's method of preserving articles of food for an indefinite time by boiling the vessel in which they are contained, and then closing it up; if air

were allowed to enter, fermentation was set up, and hence his idea that oxygen was the cause of fermentation.

In 1837 Cagniard de Latour discovered the true nature of yeast, that it was composed of living organised cells, consisting of a cell envelope and granular contents; this discovery was made about the same time by two other investigators, Schwann and Kützing, independently of one another. Kützing further discovered the cause of the acetic acid fermentation which takes place at the same time in alcoholic solutions. Each of these investigators put forward a theory, which was formulated in very much the same words—that the process of alcoholic fermentation depended upon the presence of the yeast, or sugar fungus, which extracted from the sugar and the nitrogenous matter also present the materials necessary for its nutrition, and that the elements not taken up in this way united together to form alcohol.

This theory was strongly opposed by Liebig, whose views were not essentially different from those put forward by Willis and Stahl; the yeast, as well as all animal and vegetable matter, was supposed to be in state of internal motion, which was transmitted to the bodies with which it was in contact. Pasteur, on the other hand, by a long series of experiments, definitely proved that the process of fermentation was dependent on the life of the yeast-cell, and that all fermentation was dependent upon life. Pasteur's results have been summed up by Professor Frankland as follows: "No fermentation without organisms; for each kind of fermentation a particular organism."

Besides this and similar fermentations, other changes

were known which took place without the presence of living organisms, such as the conversion of starch into sugar, and of amygdalin into benzaldehyde, prussic acid, and glucose by emulsin. Liebig thought that the yeast contained a ferment which brought about the decomposition of the sugar, but would not acknowledge that the fermentation was brought about by the life-processes of the yeast-cell. Quite recently this has been shown to be the case by Buchner, who has been able to extract from yeast a liquid which can set up alcoholic fermentation. Liebig's idea was expounded in a slightly different manner by Traube in 1858. Fermentation, according to him, is instigated by the enzymes or soluble ferments formed by the vitality of organisms, which are either excreted by them or retained within them to effect intracellular digestion. This view was accepted by Berthelot and Hoppe-Seyler, and it is the one which is most generally held at the present time. Traube's theory thus combines both the vitalistic theory of Latour, Schwann, and Kützing, and the mechanical theory of Liebig.

Another view was suggested by Berzelius, that, namely, of catalytic forces; the process being due to an act of contact, and yeast being considered to be an amorphous principle. Mitscherlich supported this view, but he recognised the organised nature of yeast.

Naegeli was an advocate of Liebig's mechanical theory, holding that the molecules in the enzyme possessed a considerable vibration; these had the power of giving up part of their movement to the slower movement of the other molecules, and thus caused a decomposition.

Since Pasteur, no further important investigations

have been made upon the chemical changes which sugar undergoes as the result of fermentation; Buchner's extract containing zymase is the only recent discovery of any importance. This discovery of Buchner's is challenged by Schunck and Manassein; the former observer showed, in 1854, that erythrozyme, the enzyme of madder, could set up alcoholic fermentation; and the latter, in 1871, found that the presence of yeast cells was not necessary for the process of fermentation, which must therefore be regarded as due to an enzyme.

In 1892 O'Sullivan showed that the hydrolytic functions of yeast followed the same course as a simple chemical interchange, the time curve being like that for invertin, and a straight line; in the case of maltose and dextrose, the rate of alcoholic fermentation was found to be proportional to the amount of these sugars, and different to the rate of the hydrolytic action of yeast upon cane sugar. A. J. Brown has found that there is no direct proportion between the weight of yeast formed and the amount of sugar fermented, and that the fermentation proceeds better with oxygen than without. This is the reverse of the view held by Pasteur, that fermentation was a starvation phenomenon, *i.e.* a phenomenon of life without air.


Quantitative experiments upon the fermentation of mixtures of dextrose and lævulose, and of maltose and dextrose, have been made by Prior and Schulze, who find that the process takes place within the yeast-cell, and is dependent on the diffusion of the sugar solution through the cell-wall; the rate of diffusion varies for the different species of yeast, and the amount of each sugar

fermented varies with its osmotic pressure. Knecht has found that two varieties of yeast ferment dextrose more quickly than *lævulose*, and he concludes that the rapidity of the fermentation of the different sugars is dependent on physical laws as stated by Prior.


During the process of alcoholic fermentation heat is evolved; its amount has been measured by Bouffard, who found that for 180 grams of sugar 23·5 Cals. were evolved, whereas the calculated amount is 32·07 Cals.

By the far-reaching investigations of E. Fischer, the number of sugars belonging to the monosaccharides has been very largely increased; these contain three, four, five, six, seven, eight, or nine atoms of carbon in their molecules, but only those containing carbon atoms which are a multiple of three, the hexoses and the nonoses, have been shown by him to be capable of undergoing alcoholic fermentation. A mixture of the trioses glyceraldehyde and dihydroxyacetone, which are obtained from glycerol by oxidation, was stated by some authors to be fermented, but Emmerling has shown that the pure substances are not acted upon, and he suggested that during the oxidation with platinum black the glycerose is converted into acrose, or d- + l-fructose. Of the aldohexoses, twelve out of a possible sixteen of which have been synthesised, only three can be fermented, and of the ketohexoses only one. These are d-glucose, d-mannose, d-galactose from the first; and d-fructose from the last. The corresponding *lævo*- compounds are not attacked, and d-galactose is fermented only with difficulty. From these facts Fischer concluded that the structure of these bodies played a very important part with regard to their capability of

fermentation; if the configuration of these four sugars be written down side by side, the small differences in their configurations can be readily seen:—



d-glucose, d-mannose, and d-fructose are thus very similar in their structure, at least three of their asymmetric carbon atoms having the same configuration; but d-galactose has only two asymmetric carbon atoms with a configuration like that in the other three sugars, which corresponds with the greater difficulty of fermentation. If the structure of the sugar is still further removed, as in d-talose, only one carbon atom having the same configuration, no fermentation at all takes place:—



That structural differences are, in all probability, the cause of the behaviour of these sugars with regard to fermentation, is seen again amongst the class of bodies known as glucosides, which behave in a very similar manner towards the enzymes maltase and emulsin.

CHAPTER V.


CHANGES IN THE GLUCOSIDES.

SINCE the majority of the substances mentioned at the end of Chapter IV. are derivatives of glucose, the changes which take place in them as the result of fermentation are most appropriately described here, before those which take place in the monosaccharides themselves, and which result in the formation of various products.


The glucosides are very widely distributed in nature, and a large number, which have not been found as naturally occurring, have been prepared artificially. These latter compounds have been very thoroughly investigated by Fischer, who has found that some very interesting relations exist with regard to their decomposition by the ferments known as emulsin and maltase, thus emphasizing his theory of the importance of configuration as being a factor of chemical change by fermentation.

The glucoside amygdalin was the first to be prepared in a crystalline form by Robiquet and Boutron in 1830, from the seeds of the bitter almond. The change which it undergoes was found by Liebig and Woehler to be due to the action of the enzyme emulsin, which exists in the kernel of the almond, and was termed synaptase by


Robiquet. These observers discovered that prussic acid and sugar were amongst the products. It is now known that benzaldehyde, prussic acid, and glucose are the products of this change, which is brought about by emulsin, and the following equation shows their formation :—



It has been discovered by Fischer that amygdalin undergoes a different transformation when it is acted upon by the ferment maltase; in this reaction only one molecule of glucose is split off, and a new glucoside is formed which may be regarded as an artificial product. Fischer named this new compound "mandelnitrilglucosid," and gave it the formula :—




Prussic acid does not appear in this transformation, but if this new glucoside be treated with emulsin, prussic acid, benzaldehyde, and glucose are formed :—




From these results Fischer concluded that the sugar group in amygdalin was present in the form of maltose, and amygdalin is therefore a maltoside.


Emmerling has quite recently obtained amygdalin from mandelnitrilglucosid and glucose, by the action of yeast maltase, which is another instance of synthetic action by enzymes; he obtained, however, only a small



7. Populin into glucose and benzoylsaligenin, only, however, by certain extracts which most probably contain emulsin, *e.g.* of *Aspergillus niger* (Hérissey):—



8. Glucovanillin into glucose and vanillin:—



and glucovanillic acid into glucose and vanillic acid.

9. Indican, the glucoside of indigo, is converted by a soluble ferment, according to Bréaudat, into indigo white and indigluclin; the latter, especially in alkaline solution, is oxidised by another ferment to indigotin; in woad, however, according to Beyerinck, isatan is present, which is decomposed by isatase into free indoxyl, which was formerly thought to be present. Indoxyl is considered by Hazewinkel to be an intermediate product in the formation of indigo, and this explains how in some processes indigo red is formed.


10. Daphnin is converted into daphnetin, and picein into piceol, and in each case glucose.

Hérissey states that the emulsin of almonds decomposes gentiopierin from *Gentia lutea*, syringin and phyllirin, and has a slight action upon ononin and

helleboreïn; the emulsin from aspergillus, however, acts easily upon these two last glucosides.

Among other glucosides which are chemically changed by ferments other than emulsin, the following may be mentioned :—


1. Sinigrin, or potassium myronate, the constituent of black mustard seeds, is converted by the enzyme myrosin into glucose, potassium hydrogen sulphate, and allyl sulphocyanate, or mustard oil :—



Apparently a molecule of water is not required for this change, but Gadamer states that the glucoside contains a molecule of water with which it crystallises and loses at 100° in a vacuum. This observer finds that allylsulphide, allylcyanide, and carbon bisulphide are produced as secondary products in the fermentation. He considers that sinigrin is the potassium salt of an ethereal sulphate derived from the hypothetical imidoxythio-carbonic acid, in which the three hydrogen atoms are replaced by SO_2OK , $\text{C}_6\text{H}_{11}\text{O}_5$, and C_3H_5 , thus :—



2. Sinalbin, contained in white mustard, is decomposed also by myrosin into glucose, sinapin hydrogen sulphate, and oxybenzyl mustard oil :—




Gadamer gives the formula $\text{C}_{30}\text{H}_{42}\text{N}_2\text{S}_2\text{O}_{15} + 5\text{H}_2\text{O}$ to this glucoside, and the change is hydrolytic as in the case of sinigrin.

3. Rubian, discovered by Schunck in 1852, is said by

him to be converted into alizarin and other compounds by an enzyme which he has called erythrozyme.


This observer thinks that three molecules break down, one giving alizarin; the second giving rubiretin, verantin, and water; whilst the third gives glucose, rubiafin, and rubiagin.

4. Xanthorhamnin is, according to Marshall Ward and Dunlop, decomposed by the enzyme rhamnase into rhamnose, probably also galactose and rhamnin or rhamnetin—




C. and G. Tanret have obtained a trisaccharide, rhamninoze, by the action of rhamnase on xanthorhamnin. This carbohydrate yields, on hydrolysis with acids, two molecules of rhamnose and one molecule of galactose, and the glucoside therefore apparently does not contain glucose in its molecule.

5. Gaultherin is hydrolysed by gaultherase, or betulase, into glucose and methylsalicylic acid:—



6. Tannin is hydrolysed by tannase into glucose and gallic acid:—




The compound of tannin with gelatine, tannates, methyl- and phenyl-salicylates, are also said to be changed by this enzyme.

7. A glucoside in *Ecballium elaterium* is, according to Berg, decomposed by an enzyme (which he has named elaterase) into elaterin.

8. Lotusin, a glucoside obtained by Dunstan and

Henry in 1900, is said by these observers to be transformed by an enzyme, lotase, into glucose, hydrogen cyanide, and lotoflavin. Acids convert it into two molecules of glucose, one molecule of hydrogen cyanide, and one molecule of lotoflavin.

Lotusin is a lotoflavin ester of maltose cyanohydrin:—



and the observers point out that amygdalin and lotusin are the only glucosides which yield hydrogen cyanide on hydrolysis.

9. Kolanin, the glucoside of caffeine, yields under the action of a ferment, glucose, caffeine, theobromine, and kola-red; the same products are given by cacaonin. These glucosides are considered by Schweitzer as made up of: (a) kolanin: one molecule of kola-red, three molecules of glucose, and one molecule of caffeine with the elimination of four molecules of water, with the formula $\text{C}_{40}\text{H}_{56}\text{N}_4\text{O}_{21}$; (b) cacaonin: one molecule of cacao-red, $\text{C}_{17}\text{H}_{12}(\text{OH})_{10}$, six molecules of glucose, and one molecule of theobromine, with the loss of eight molecules of water, with the formula $\text{C}_{60}\text{H}_{86}\text{O}_{15}\text{N}_4$.


Knox and Prescott, however, do not think that caffeine and kola red are products of one glucoside.

We now come to the artificial glucosides, with regard to which so many important facts have been discovered by Emil Fischer. By this investigator not only have derivatives of glucose been prepared, but also of mannose, galactose, maltose, etc. In general, two derivatives are



obtained by the action of hydrochloric acid upon the sugar and alcohol, and these are termed α - and β -glucosides. It is among the methyl-glucosides that the majority of observations have been made with regard to their behaviour towards ferments, especially emulsin and maltase.

It has been found that the α - compounds are only hydrolysed by maltase, and not at all by emulsin, which attacks only the β - compounds. The ease with which these compounds are attacked also varies; whereas the glucosides are easily hydrolysed, the galactosides are attacked with difficulty, and the mannosides not at all. Fischer, noting these differences, concluded that the structure of these compounds played a very important part with regard to their behaviour to these enzymes, and that the enzymes themselves must be built up in a manner correlative to that of these glucosides, like lock and key.

If the formulæ of these compounds be written down, it is easily seen that one of the asymmetric carbon atoms is different in the α - and β - compounds:—





α - and β - glucosides.

 α - and β -galactosides.

This difference in all probability accounts for the different action of these enzymes.

Very remarkable evidence of this configurative difference, as being the cause of the enzyme action, exists in the case of the xylosides, whose structure is so very similar to that of the glucosides:—



From the formulæ it will be seen that one of the asymmetric carbon atoms fails in the d-xyloside, and apparently it is this carbon atom in the d-glucoside (which is marked with an asterisk) which defines the action of the enzyme, and which influences the entire configuration of the d-glucosides.

The disaccharides maltose and lactose, as already mentioned, also show similar differences in their susceptibility to these enzymes, and the idea of

configuration therefore gains further support from these examples.

Quite lately Fischer and Armstrong have shown that the β -methyl-, β -ethyl-, and β -phenyl-galactosides are also hydrolysed by kefir-lactase.

CHAPTER VI.

CHANGES IN THE MONOSACCHARIDES.—Continued.

THE chief of the chemical changes other than alcoholic fermentation, which occurs in the monosaccharides, is known as lactic acid fermentation, and it takes place principally in the hexoses, glucose, fructose, and galactose.


Pasteur was the first to definitely show that this fermentation was produced by micro-organisms, though Blondeau, in 1847, first observed them without discovering their significance. It is very probable that an enzyme is the cause of this change, but as yet it has not been isolated from these micro-organisms. In the organs of animals, especially in muscles, lactic acid is always present, and Nasse has pointed out the analogy of its formation to enzyme actions. Oppenheimer has recently investigated the subject with regard to the presence of a lactic acid-forming enzyme, but his experiments are not conclusive as to its existence.

It was formerly supposed that lactose and other polysaccharides could be directly converted into lactic acid, sometimes, perhaps, with difficulty; but it is now known that hydrolysis of these substances into monosaccharides, as already described, takes place before lactic acid can be produced.

This change can be expressed by the equation:—



Apparently no water is required for this process, but, as in the case of alcoholic fermentation, Baeyer considers that two molecules of water may take part in the transformation, thus:—



The reaction, however, is not so simple as expressed in the equation, as acetic acid and other products are also generally formed at the same time.

This method of producing lactic acid has been known for a very long time, in fact, it is probably the oldest acid known, and was first prepared in this way by Scheele. In order to prepare it, calcium carbonate must be added to the solution to neutralise the free acid as fast as it is formed, as the specific micro-organisms cannot grow in an acid medium, and the production of lactic acid, therefore, necessarily would cease. If the acidity is confined to 0·02—0·5 per cent., so as to prevent the secondary formation of butyric acid, 98 per cent. of sugar can be converted into lactic acid (Claffin).

The lactic acid produced in this way is generally inactive, but certain micro-organisms form d-lactic acid, and

Schardinger has obtained l-lactic acid in one instance. It is very probable that the inactive lactic acid is formed first, and is then converted into one of its active varieties, as shown by Frankland and MacGregor in the case of the formation of d-lactic acid.

β -hydroxypropionic acid, $\text{CH}_2\text{OH}-\text{CH}_2-\text{COOH}$, the isomer of ordinary lactic acid or α -hydroxypropionic acid, was obtained by Hilger, by the fermentation of inosite; it was identified by oxidation to malonic acid. Vohl has repeated this experiment, but he obtained α -hydroxypropionic acid; this acid (generally termed lactic acid of fermentation) is thus always produced in fermentation.

Lactic acid is also a product of the fermentation of certain polyhydric alcohols, *e.g.* sorbite, inosite, mannite, and dulcite; also of rhamnose (Tate) and other pentoses; it can also be formed by fermentation from malic acid, carbon dioxide being evolved during the process:—




Wehmer, in 1892, showed that certain monosaccharides, but especially glucose, could undergo yet another change as the result of fermentation, citric acid being the product formed. Two species of micro-organisms were found capable of producing this result, carbon dioxide being evolved during the process. The citric acid obtained is identical with that found in lemon juice, and is the principal product. For this process air is a necessity, the carbon dioxide evolved during the fermentation even hindering the action of these micro-organisms, and consequently it must be allowed to escape. Over 50 per cent. of the glucose can be transformed into citric acid; but if the acidity of the solution reaches 20 per

cent., it becomes deleterious to these micro-organisms, and must be removed by neutralisation, in order that the total quantity can be obtained; if the fermentation is allowed to go too far, the citric acid itself is decomposed. Wehmer has since obtained three other micro-organisms which can produce this fermentation.

Oxalic acid is another product obtained by the fermentation of sugar solutions, which may consist of glucose, galactose, cane sugar, lactose, maltose, dulcitol, mannitol; glycerol can also be converted into oxalic acid in the same way. The micro-organisms which produce this change are hindered in their action by excess of the acid, and a good yield of oxalic acid is only obtained when it is neutralised as fast as it is formed by calcium carbonate, which can be added to the solution. According to Banning, fifteen species of micro-organisms are now known which can produce oxalic acid, all of them from glucose, and the majority from arabinose; glycerol and erythritol by about half of them, as also pyrotartaric acid and glycollic acid. This investigator gives a list of what substances and by how many of these micro-organisms each of the substances is converted into oxalic acid.

The curious phenomenon, observed in wines and saccharine liquids, known as "ropiness," is the result of a fermentation, and is due, according to Pasteur, to a particular organism. It is observed sometimes in the manufacture of cane sugar, and the result is a peculiar viscosity. The change that occurs is probably caused by two organisms, since Pasteur has observed larger and smaller particles of a different kind in the liquid; one of these produces the peculiar gum which resembles dextrin,

as on oxidation it yields oxalic, not mucic acid, and the other produces mannite. During the process, carbon dioxide is evolved; that the process is in all probability due to two organisms is further seen by the varying quantities of gum and mannite produced, that of the gum being very often much larger. Monoyer has proposed the following equations for these two changes:—




The gelatinous matter, called by Scheibler dextran, is probably a derivative of cellulose, and it can be formed by several micro-organisms, *e.g.* *Streptococcus hornensis*, which produces it from media containing 20 per cent. of glucose (Boekhout). A gum, resembling in properties a galactan, since it yields mucic acid on oxidation, is produced by *Bacillus lactis aërogenes* in solutions of lactose or galactose, but not glucose (Emmerling).

These changes, which sugar can undergo, are thus very various. Nef has compared the action of enzymes with the action of those reagents which cause the decomposition of methylene compounds, and regards the enzymes as substances exerting a directive influence on the decomposition of the sugars. Very many other theories have been advanced to explain the processes of the metamorphosis of sugar into alcohol and carbon dioxide, but the suggestion brought forward by Baeyer, in 1870, that water is the active agent, seems to be the most reasonable. He considered that addition of the elements of water took place at certain points, and that by alternate separation and addition in a different order, an accumulation of oxygen atoms at certain parts of the molecule might take

place; at these points there would be weakness, and disruption would occur.

The following formulæ of Baeyer's make this clear—



where the separating elements are indicated by dots.

In 1895 Armstrong drew attention to the fact that accumulation of oxygen atoms might take place at non-contiguous carbon atoms, and he considers that this is a special feature; different effects would thus be produced by different enzymes, since they would affect different systems, and the direction of the attack would depend on the asymmetry of both hydrolyte and hydrolyst. This accounts for the different action of enzymes, as in the case of the glucosides.

CHAPTER VII.

CHANGES RESULTING IN THE FORMATION OF OPTICALLY ACTIVE PRODUCTS.

As has already been pointed out, enzymes and living ferments are very selective in their action upon the carbohydrates and organic acids. Pasteur, in his classical experiments upon the tartrates and racemates, in 1848-1853, was the first to point out the selective action of micro-organisms. He grew a penicillium upon the ammonium salt of racemic acid, and obtained lævo-tartaric acid, the dextro- variety being assimilated by the organism.

This method of Pasteur's has been employed in many cases for the separation of optically active compounds; the separation of inactive mandelic acid by Lewkowitsch, in 1882, into its constituents, and the resolution of glyceric acid by Frankland and Frew, in 1891, by the *Bacillus ethaceticus*, which was grown upon its calcium salts, are two very good examples of this selective action.

Frankland has investigated the action of his *Bacillus ethaceticus*, and of the pneumococcus of Friedländer, upon several carbohydrates, *e.g.* maltose, lactose, raffinose, dextrin, mannitol, glycerol, glucose, and arabinose; the chief products were alcohol and acetic acid. Upon

erythritol, glycol, and dulcitol, *Bacillus ethaceticus* had no action; the author emphasises this curious behaviour with regard to mannitol and dulcitol, and considers this to be a case of the selective action of these organisms. He has obtained only one organism which can ferment both mannitol and dulcitol; the same products are obtained by the action of this *Bacillus ethacetosuccinus*, but rather more succinic acid is formed.

Grimbert has shown that, by the action of Friedländer's pneumococcus, mannitol gives lactic acid, and dulcitol gives succinic acid. In conjunction with Ficquet, he has found that *Bacillus tartricus* acts upon mannitol, and not dulcitol.

Lactic acid of fermentation is inactive, but by some organisms, e.g. *Bacillus acidi lævolactici*, the lævo-acid is produced from dextrose, cane sugar, lactose, and glycerine. Schardinger has shown that alcohol, acetic acid, and lævo-lactic acid are the principal products.

Tate has obtained lævo-lactic acid from dextrose and mannitol, but rhamnose gave inactive lactic acid when acted upon by the lævo-lactic ferment.

Linossier has resolved inactive lactic acid by *Penicillium glaucum*, and has obtained dextro-lactic acid similar to that obtained from muscles.

Frankland and MacGregor have also obtained d-lactic acid from calcium lactate, by the action of a particular organism.

In fermentation, the inactive and the two active varieties of lactic acid are therefore produced; according to some observers the variety obtained depends on the micro-organism, but according to Kayser, the medium

upon which the micro-organism is cultivated is also concerned. Péré's results show that the lævo variety is in most cases produced by two colon bacteria from various carbohydrates; and Harden has obtained the same result with the *Bacillus coli communis*, which attacks mannitol, but not dulcitol. The amount of alcohol produced from mannitol is twice as great as from glucose, and this result is ascribed by Harden to the presence of the group $\text{CH}_2\text{OH}-\text{CHOH}$, which is contained once in glucose, $\text{CH}_2\text{OH}-(\text{CHOH})_4-\text{CHO}$, but twice in mannitol, $\text{CH}_2\text{OH}-(\text{CHOH})_4-\text{CH}_2\text{OH}$. This view, that the alcohol produced originates from this particular group, is confirmed by the action of this organism on glycerol, $\text{CH}_2\text{OH}-\text{CHOH}-\text{CH}_2\text{OH}$, the chief products of this action being alcohol and formic acid. As regards the formation of the lactic acid, two molecules of mannitol or glucose break down in the fermentation, and the middle groups of $\text{CHOH}-\text{CHOH}-\text{CHOH}$ are supposed to give rise to it by the interchange of a hydrogen and an oxygen atom between the two terminal radicles of one group or between those of the two groups, derived from the two molecules of the sugar. The unaltered asymmetric group, $-\text{CHOH}-$, in the centre would remain asymmetric in the new molecule of lactic acid, but the change, being brought about in the presence of the asymmetric agent—whether the organism itself or possibly an enzyme—would take place entirely in one direction or more rapidly in one direction than the other, and an active acid, independent of the previous configuration of the group, but dependent on the nature of the organism, would result. This view is based upon the theories of Fischer and of

Marckwald regarding the influence of configuration of the agent causing the fermentation.

In 1892 Buchner investigated the action of *Penicillium glaucum* and *Aspergillus niger* on fumaric and maleic acids. Although these acids are so closely allied to each other, only the acid ammonium salt of fumaric acid was affected, that of maleic acid being entirely unaffected.

In the same year Purdie and Walker grew *Penicillium glaucum* upon the acid ammonium salt of inactive ethoxysuccinic acid obtained by the action of alcohol on fumaric acid, and found that the lævo- variety was consumed, the dextro- variety being left unaltered.

These are two other cases of selective fermentation by these moulds which were first shown by Pasteur to have this selective action.

Emmerling has very recently investigated certain of the amino-acids with regard to their use as food-material by micro-organisms, and he has found that only the α -amino-acids can be employed, and only by certain moulds; γ -aminobutyric acid was assimilated by all, whereas α -aminobutyric acid was of little value. Most remarkable was the action upon serin and isoserin; the former was very good, and the latter useless as a food-material.


Carbon compounds, as food-material for bacteria, have been studied by Bokorny, and his results are of interest with regard to the assimilation of various alcohols; whereas methyl, ethyl, and amyl alcohols could be employed, propyl alcohol was good for fungi, but not for bacteria, and benzyl alcohol was a strong poison. Acids were also investigated; of these acetic, lactic, succinic, tartaric were good, and oxalic, and valerianic acids were of no use.

CHAPTER VIII.

CHANGES IN ESTERS.


THE decomposition of fats and oils by the enzyme known as lipase or steapsin, is another hydrolytic process ; it may conveniently be described here before the other processes produced by oxidation, etc., are considered.

It was discovered by Claude Bernard that fats taken as food by animals were not assimilated in that form, but that they were transformed, by an enzyme present in pancreatic juice, into free fatty acid and glycerine. This change that the fats, or the esters of glycerine with palmitic, stearic, and oleic acids, generally known as palmitin, stearin, oleïn, undergo, may be represented by the equation :—



An enzyme of this nature is also present in plants, which can produce a similar change in oils. These are transformed by the plant into starch, and were in 1859

considered by Sachs to be converted directly. In 1871 Müntz showed that free fatty acid was formed from the oil; and in 1876 Schützenberger proved that glycerine was a product of this change as well as fatty acid. It was then supposed by Detmer, in 1880, that fatty acid gave rise to starch, and he suggested the equation—



to represent this change.

It was then supposed that the glycerine gave rise to the starch, but it has been since shown by Mazé that sugar can be formed from the oil, and that this is consequently the antecedent of starch. These changes are probably all produced by the action of ferments in the plants.

The rancidity of butter, caused by the action of micro-organisms, is a common example of this decomposition, and an enzyme has been obtained by Laxa from moulds which produces this change; it also decomposes monobutyrim and glycerides of insoluble acids, those of the highest molecular weight being first attacked.

Recent work upon the action of the enzyme known as lipase shows that it is also capable of hydrolysing esters other than those of glycerine; *e.g.* acetic ester, succinic ester, benzoic ester, salicylic ester. This power of lipase of being able to hydrolyse esters was in 1900 confirmed by Kastle and Loevenhart, who have also shown that the higher the molecular weight of the acid (in the acetic series), the more readily is it hydrolysed by lipase. The rate of the hydrolysis was shown to be dependent on the concentration of the enzyme, but independent of that of the ester; the velocity was not constant, but gradually

diminished as the hydrolysis proceeded. The hydrolysis has been shown by these observers to be an incomplete and a reversible action, as they have obtained butyric ester from butyric acid and alcohol. Butyric ester was also used to determine the activity of solutions of lipase.

Kastle has also investigated the action of lipase upon the salts of certain acid esters, and has found that they are not hydrolysed, whereas the neutral esters are hydrolysed; *e.g.* diethylfumarate, diethyloxalate, diethylphthalate are hydrolysed, but not sodium ethylfumarate, potassium ethyloxalate, barium ethylphthalate. Apparently, therefore, there is no hydrolysis when the ester is electrolytically dissociated, and he assumes that hydrolysis is preceded by combination of the ester with the enzyme, and that combination cannot take place with ions, but only with undissociated molecules.

Hanriot has also found that the lipase of horse's serum can exert a synthetical action, and he has also shown that the acids of higher molecular weight are hydrolysed more quickly than those of lower. Excess of acid was found to arrest the hydrolysis, and Hanriot supposes that an unstable compound of the enzyme with the acid is formed, as the enzyme can be regenerated when attenuated by chemical action, and can return to its former activity. The hydrolysis is explained by the supposition that the ferment is a weak base which combines with the fatty acid to form a readily dissociated salt, comparable to ferric oxide and alumina, which also form unstable salts with organic acids, and can promote a slight hydrolysis of glyceryl esters.

Together with Camus, Hanriot has invented a method of quantitatively estimating lipase; monobutyrim, which was first shown by Berthelot to be hydrolysed by this enzyme, is used, and the butyric acid liberated is titrated.

CHAPTER IX.

CHANGES IN UREA AND URIC ACID.

THE last of the changes resulting from fermentation, in which the process is hydrolytic and the products are definitely known, is the conversion of urea into ammonium carbonate.


This change has been known for many years, and was recognised by van Helmont as a process of fermentation, but the cause of the change has only of late years been determined. It is a process which takes place whenever urine is allowed to stand; the acid reaction disappears, and it then becomes alkaline, and smells strongly of ammonia.

The equation $\text{CO}(\text{NH}_2)_2 + 2\text{H}_2\text{O} = (\text{NH}_4)_2\text{CO}_3$ represents this change, which was first explained by Dumas in 1830.

Micro-organisms were shown by Pasteur to be the cause, and van Tieghem succeeded in isolating a particular one, known as *Bacterium ureæ*. Since then many other bacteria have been found to possess this power.


Miquel supposed that an enzyme was the cause of this change, and Musculus has obtained an extract from certain bacteria which can produce this conversion. It has been termed urease by Miquel; the enzyme, according to Lea, is excreted at the death of the microbe.

Uric acid is stated to be changed by the same organisms into the same product, with the evolution of carbon dioxide. F. and L. Sestini have suggested the following equation—




to represent the result which they obtained.

E. Gérard, in 1896, found that uric acid, when dissolved in a solution of disodium phosphate, was converted by micro-organisms into urea and ammonium carbonate; he suggested that the ammonium carbonate was a secondary product resulting from the urea by the action of a urophagic micro-organism, the urea being the true product of the fermentation of uric acid. This, a little later, he was able to prove, and he gave the following equation to represent this change:—



It was shown by Magnier de la Source that when uric acid was boiled with water, dialuric acid was formed, which on hydrolysis gave urea and tartronic acid, and hence its up to the present hypothetical formation in the fermentation of uric acid.

According to Burri, hippuric acid in the form of its calcium salt is fermented, but with greater difficulty than uric acid, into the same product, namely, ammonium carbonate. Van Tieghem states that *Bacterium ureæ* converts hippuric acid into its constituents, benzoic acid and glycocoll, thus:—



This acid and also taurocholic acid can be hydrolysed by a ferment excreted by many bacteria. That the decomposition is caused by a ferment is shown by the fact that the same hydrolysis occurs when the bacteria are destroyed with ether (Hoppe-Seyler).

CHAPTER X.

CHANGES OCCURRING AS THE RESULT OF OXIDATION.

THE formation of acetic acid by fermentation is a process differing from those changes already described, by not being hydrolytic, but produced by oxidation. This process is used on the large scale for the production of vinegar, and has been known for a long time. Acetic acid, the principal constituent, was one of the first acids known.

Before the time of Lavoisier, it was not known that oxygen was necessary for this conversion of alcohol into acetic acid, which takes place in wines when they are allowed to stand in the air; it was, however, generally supposed that air was necessary. Lavoisier showed that the oxygen of the air was absorbed in the process.

In 1821 Davy discovered that alcohol could be converted into acetic acid when platinum black was moistened with alcohol; it was subsequently shown, in 1823, by Döbereiner, that oxygen was used up in this change, and that the alcohol was converted into acetic acid and water.

In this process a skin is formed on the surface of the liquid, which was named *Mycoderma aceti*, or, mother-of-vinegar, by Persoon, in 1822. Kützing, in 1837,


re-investigated this scum, and concluded that it was of a living nature. This mycoderma was held by Liebig to be non-living, and porous like platinum black; its action depended on the pores in its substance, in which oxygen was mechanically contained; the formation of acetic acid was due to the movement set up by the decomposing matter, oxidation being thus caused.

The subject was carefully investigated by Pasteur, but he did not at once recognise the *rôle* played by the mycoderma, which was supposed to act like platinum black; he found that the change was only effected during the life of the organism, and that if the action were continued, carbon dioxide and water were produced from the acetic acid. In 1873 it was shown by von Knieriem and Ad. Mayer that these two processes of producing acetic acid were not comparable, since the mycoderma could only decompose weak alcoholic solutions at a temperature not above 40°; the platinum black could effect the transformation of strong alcohol, even at a high temperature. It is now generally supposed that the change is due to the direct action of living organisms, since no enzyme as yet capable of producing this change has been obtained from them.

The most recent and careful observations were carried out by A. J. Brown, in 1886, with two particular organisms which he had isolated. He has confirmed the older observation of Pasteur's, that if the process be continued beyond the complete oxidation of the alcohol to acetic acid, carbon dioxide and water are formed from the latter.

In this process Brown has shown that the alcohol


passes through two stages: aldehyde is formed first, and is subsequently converted into acetic acid, thus:—



Brown has made several other investigations upon the oxidising action of these two micro-organisms; propyl alcohol was converted into propionic acid, but the oxidation of methyl, isoprimary butyl, and amyl alcohols could not be effected. Henneberg has also shown that propylalcohol can be oxidised to propionic acid by the acetic acid micro-organisms.

Glucose was oxidised to gluconic acid in the presence of calcium carbonate, added to neutralise the free acid formed,—a confirmation of Boutroux's result of 1880.

The following equation is given by Brown to represent this change—



the oxidation thus affecting the aldehyde group.

Further, mannitol is oxidised to lævulose; since mannitol can be obtained from d-glucose by reduction with sodium amalgam, and mannitol can be converted by the *Bacterium aceti* into lævulose, a new method of converting d-glucose into lævulose has been discovered by Brown.


In the vinegar plant, a membrane is formed which contains cellulose, and Brown shows that this can be formed from dextrose, mannitol, and lævulose, but not from cane sugar, starch, or alcohol. He proposes the name of *Bacterium xylinum* for this microorganism.

The oxidation of sorbitol to sorbose by the sorbose

bacterium, which is probably identical with *Bacterium xylinum*, has been described by Bertrand, who has also shown that it can oxidise mannitol to lævulose, and glycerol to dihydroxyacetone. Here the carbon atoms in the molecules are a multiple of 3; but this is not a necessity, since the micro-organism can also oxidise erythritol, arabitol, and perseitol; in each case two hydrogen atoms are removed. Bertrand states that only those alcohols seem to be oxidised which contain a CHOH chain where no hydrogen atom exists on the same side as the hydroxyl group, which is attacked in the oxidation. He has also shown that xylose can be oxidised to xylonic acid, aldose to a monobasic acid, arabinose to arabonic acid, glucose to gluconic acid, and galactose to galactonic acid.

Vincent and Délachanal obtained Bertrand's results with regard to the oxidation of mannitol to lævulose a year earlier, and Emmerling has recently confirmed the oxidations of sorbitol and glycerol by the sorbose bacterium.

Boutroux has obtained a hydroxygluconic acid by the oxidation of gluconic acid in the presence of calcium carbonate, and he gives the formula—



to this acid from the results which he obtained by oxidising it with dilute nitric acid.

CHAPTER XI.


CHANGES OCCURRING AS THE RESULT OF OXIDATION.—Continued.

IN nature, both in the animal and the vegetable kingdom, changes are produced by enzymes, known collectively now as oxydases, which are the result of oxidation.

Yoshida, in 1883, noticed that the juice of the lac tree of South-Eastern Asia underwent the peculiar changes of turning black and setting hard, producing the material used in lacquer varnish. He thought that this was due to an enzyme, and he investigated the change from this point of view. The juice was found to consist of four substances—urushic acid, to which he gave the formula $C_{14}H_{18}O_2$, gum, water, and a nitrogenous body which contained an enzyme. The change caused by this enzyme is produced in the urushic acid, which is oxidised, forming oxyurushic acid, $C_{14}H_{18}O_3$. That oxidation was the cause of the change was shown by the transformation of urushic acid into oxyurushic acid by chromic acid. Some ten years later Bertrand reinvestigated this change, and confirmed Yoshida's result. He gave the name laccase to the enzyme, but he called urushic acid laccol, as it resembles in many of its properties the polyatomic phenols. This observer found that an alcoholic solution of laccol turned brown at once, and ultimately became black under the

influence of the ferment. Bertrand has also investigated the action of this enzyme upon various phenols, but especially hydroquinone and pyrogallol. The hydroquinone became rose-red and green crystalline scales, with a metallic lustre separated out, consisting of quinhydrone, the odour of quinone being apparent. When performed in a sealed tube, it was found that oxygen had been absorbed.

The change may be expressed by the equation :—



The pyrogallol was converted into purpurogalline, which sublimed on heating, giving orange-red needles, soluble in alcohol and acetic acid.

An alcoholic solution of gum guaiacum becomes blue in the presence of air and a very small quantity of laccase, but if the proportion of laccase is large, the blue colour becomes green and then yellow; this solution serves as a convenient test for laccase, which has been found to be of very general occurrence in vegetables and fungi, and probably plays a part in the respiration of plants.

Bertrand has investigated the action of laccase upon dihydric and trihydric phenols, amidophenols and diamidobenzenes, and he has found that the ortho- and para- derivatives are readily oxidised by oxygen or air in the presence of laccase, whilst the meta- compounds are not affected, or only with difficulty. The rate of oxidation seems to depend upon the tendency of the compound to form a quinone. The author concludes that laccase acts most readily, if not exclusively, upon compounds of the benzene series which contain not fewer than two

hydroxyl or amidogen groups in the ortho- or para-positions.

Other compounds oxidised by laccase are pyrocatechuic acid, gallic acid, tannin, and hexahydric phenol, $C_6(OH)_6$; carbon dioxide is liberated in the cases of gallic acid and tannin. The simple monatomic phenols are not acted upon.

The activity of this enzyme is curiously associated with the presence of manganese, and Bertrand states that the activity is proportional to the amount of manganese, since the absorption of oxygen in the presence of manganese was found to be considerably greater than when no manganese was present. He has also found that the manganese cannot be replaced by other metals.

Bertrand has suggested that the oxydase may be a special combination of certain proteids containing acid groups with manganese, the proteid group varying for each enzyme. Under such conditions, the manganese can be kept in solution, the oxygen being conveyed by the manganese, and the proteid constituting the enzyme.

Bourquelot and Bertrand have further shown that the blue colours produced in fungi when they are cut open, are formed by the action of an oxidising enzyme, which acts only when the air has direct access to the part; in all probability laccase is the enzyme which produces this colouration. Schönbein considered that the colouration was due to the action of ozone upon a chromogen in the fungus, and that the fungus contained a substance capable of forming ozone from the air. That a definite chromogen is present in fungi was shown by Ludwig, in 1872,

who has thus confirmed Schönbein's supposition of its existence.

Certain other fungi, *e.g.* *Russula nigricans*, produce a red and finally a black colour. This change has also been investigated by Bertrand, who has been able to isolate tyrosine from the juice of this fungus and from the root of the beet, and he has shown that the red colour is produced from this body by another enzyme, isolated from the fungus, which he has called tyrosinase; it is not brought about by laccase nor by oxidising agents. A solution of tyrosine, when acted upon by this ferment, becomes red and then black, and finally an amorphous precipitate is formed in the solution.

Bourquelot has investigated the changes produced by the oxidising enzymes in the extracts of mushrooms, and he has found that all the cresols, resorcinol, guaiacol, and eugenol, which gives a white precipitate of vanillin, are acted upon; also m-toluidine, xyloidine, and aniline "for red," which is converted into a colouring matter like magenta.

Even in solutions containing 50 per cent. of methyl or ethyl alcohol, these extracts are able to readily oxidise tyrosine. He has, therefore, tried the oxidising power of these extracts upon insoluble phenols, with the result that the o-, p-, and m-xylenols, thymol, carvacrol, α - and β -naphthol, are all acted upon; α -naphthol becomes blue-violet, and a dull blue precipitate is formed, whereas β -naphthol gives a white precipitate which turns yellow; phenolic esters are also oxidised.

Since laccase and tyrosinase exist together in the extracts of certain fungi, it is not quite certain whether

the above oxidations are produced by tyrosinase, though in all probability it is this enzyme which is the cause of these changes; this oxydase is also said to occur in animals and in bacteria (Roux, Von Fürth and Schneider, Lehmann).

Sarthou states that the latex of *Schinus Molle*, which is a milky fluid, becomes blue when it is exposed to the air, and that this change is effected by an enzyme termed schinoxydase. This ferment can oxidise quinol, resorcinol, pyrogallol, in the presence of atmospheric oxygen; the ash of the enzyme, which was isolated, contained iron, and its oxidising power is almost proportional to the amount of iron. He supposes that iron plays a part in this change like manganese in the changes produced by laccase.

Certain wines undergo a peculiar disorder known as "casse" or "cassure." This has now been proved to be due to the presence of an oxydase enzyme named cœnoxydase by Cazeneuve, which can decolourise the wine and produce a red-brown precipitate. Bouffard, who has studied this disorder, has found that a temperature of 60° prevents it, and that it can also be prevented by a small amount of sulphurous acid. It is probably upon the tannins in the wine that this enzyme acts, as the change does not take place if they are extracted with ether; perhaps the cœnoxydase is removed together with the tannin.

This enzyme has also been found in the juices of apples, pears, etc., and is the cause of their browning when they are cut open; according to Lindet, it is the tannin in the apple-juice which is oxidised by this enzyme.

The "bouquet" of wines is stated by Tolomei to be caused by an oxidising enzyme in the yeast, and he showed its existence in several species; an oxydase also exists in beer yeasts, and is the cause of an extract becoming brown. According to Grüss, the oxydase of yeast oxidises tetramethyl-p-phenylenediamine, but not guaiacol, and he proposes to divide oxydases into guaiacol-oxydases and amino-oxydases, as he has found oxidising enzymes, similar to that in yeast, in plants.

The odour of valerian is another change caused by an oxydase enzyme (Charles).

In the manufacture of indigo, indigo-white is the product obtained, which on exposure to the air becomes blue; an enzyme present in the original glucoside has, according to Bréaudat, this oxidising action upon the indigo-white.

In the animal organism, several oxidation processes occur as the result of oxidation by fermentation; for instance, phenol and benzoic acid or hippuric acid are the products obtained when benzene and toluene are taken as food.

An alkaline solution of p-phenylenediamine mixed with α -naphthol, is coloured blue when extracts of organs are added to it. Ehrlich first showed that this was due to the formation of indophenol. It serves as a test for the presence of oxidising ferments together with the other colour reactions, with guaiacum tincture and guaiacol. Salicylic aldehyde and benzylalcohol have been used by Schmiedeberg as a means of quantitatively estimating the ferment, the amounts of salicylic acid and benzoic acid produced being determined. Formaldehyde was used by

Pohl, and arsenious acid by Spitzer, for the same purpose. The most important of the oxydases of the animal organism is the one named salicylase by Abelous, which oxidises salicylic aldehyde to salicylic acid, and it occurs in most of the organs of the animal body.

Medvedeff, who has investigated the oxidising action of liver extracts on salicylaldehyde, states that the amount of oxidation is proportional to the square root of the concentration of the aldehyde, since this is the proportion of the substance dissociated in solution, but directly proportional to the concentration of the ferment.

According to Abelous and Biarnès, oxygen is consumed and carbon dioxide is liberated when salicylic aldehyde is oxidised to salicylic acid by liver extracts.

Spitzer has been able to convert the nuclein bases into uric acid by extracts of the liver and the spleen in the presence of air which was bubbled through the solution, putrefaction being carefully excluded; xanthine and hypoxanthine were almost quantitatively converted into uric acid; adenine and guanine, however, were oxidised to a lesser extent.

A third oxidising ferment, according to Abelous and Biarnès, has no action on salicylic aldehyde, but turns guaiacum tincture blue.


Cavazzani infers the existence of an oxidising ferment—cerebro-spinase—in the cerebro-spinal fluid, as it gives with pyrogallol and sulphuric acid a crystalline deposit of purpurogalline; with tannic acid it gives a brown, with quinol a rose colouration, and with o-toluidine a rose-violet precipitate.

The disappearance of sugar from blood which has been

shed, is another case of oxidation in animal tissues. Seegen has shown that it is not caused by micro-organisms, as the same process takes place in the presence of chloroform; it is due to a glycolytic ferment produced by the leucocytes, or white blood corpuscles, and it exists not only in blood, but in various organs. It is not yet definitely known what change the sugar undergoes: the experiments of Oppenheimer have led to no definite conclusions, since he could not detect alcohol or acetone, though he obtained traces of iodoform; nor as yet has he been able to prove the formation of lactic acid, which is very probably produced in the process.

The light-producing organs and the eggs in the ovary of the female glowworm give a blue colour with guaiacum tincture. Dubois considers that the active agent in the production of light is an enzyme which he has named luciferase, but he believes it is not a result of oxidation.


Waters rich in iron in the form of ferrous hydrogen carbonate, contain several species of micro-organisms, which are able to oxidise this compound after it has passed by osmosis through their cell-walls, and convert it into ferric hydroxide with the evolution of carbon dioxide, thus:—



This change can be followed in the deposited layers in the sheaths of these microbes. At first they are pale yellow, and can be decolourised by water containing carbon dioxide; but later on they become brown-red, and are only decolourised by dilute hydrochloric acid; finally this reagent cannot decolourise them. We have to thank Winogradsky for the majority of the investigations upon

these iron bacteria, which can only grow when ferrous carbonate is present, ferric oxide even being useless as a form of iron. Molisch states that manganese can replace the iron. The deposits of ferruginous ochre and bog iron ore are said to be the result of the action of these bacteria.

Hydrogen sulphide in well-waters is oxidised by the sulphur bacteria to sulphur, in accordance with the equation—



oxygen being necessary for this transformation. This change was definitely shown in 1886 by Winogradsky, and further investigations were carried out by Jegunow, who has shown that the sulphur in a higher level of the bacterial cell is oxidised to sulphuric acid, which is excreted. The reverse process of reduction of sulphates will be described in the next chapter.

CHAPTER XII.

CHANGES OCCURRING AS THE RESULT OF REDUCTION.

DE REY-PAILHADE, in 1889, observed that hydrogen sulphide was formed from sulphur, when it was mixed with yeast, and he attributed this change to the action of a reducing enzyme, which he termed philothion. Sostegni and Sannino have obtained the same result; they added flowers of sulphur to a solution of sugar, which was fermented with a pure cultivation of yeast, and found that hydrogen sulphide was evolved. Salkowski has noticed the production of hydrogen sulphide in wine, but he attributes its formation to the action of bacteria on the sulphur compounds present in it.

Philothion, sometimes alone and sometimes with laccase, has been found by De Rey-Pailhade to be contained in germinating seeds; he has shown that it contains a proteid radicle which can enter into loose combination with manganese or hydrogen in the same way as Bertrand showed in the case of laccase.

The power possessed by philothion, of forming hydrogen sulphide from sulphur, is stated by Cossettini to be lost when it is filtered through a Chamberland filter, and Pozzi-Escot finds that philothion does not give the blue colour with guaiacum tincture and hydrogen peroxide,

which is the usual test for these enzymes. Hydrogen peroxide, however, is decomposed with the evolution of oxygen.

This observer identifies philothion with catalase, an enzyme stated by Loew to be of universal occurrence, and to whose presence the power of decomposing hydrogen peroxide, possessed by all enzymes, is due; this observer states that catalase exists in an insoluble and a soluble form, termed α - and β -catalase respectively.

Pozzi-Escot finds that philothion can also cause the hydrogenation of selenium and phosphorus, but not, or only very slightly, of tellurium and arsenic.

The production of hydrogen sulphide by micro-organisms has been investigated by Zelinsky, who isolated a bacterium—*Bacterium hydrosulfureum ponticum*—from the ooze at the bottom of the Black Sea; it can produce hydrogen sulphide not only from proteid matter, but also from inorganic salts, such as sulphates, sulphites, thiosulphates, and even ammonium thiodiglycollate when they are added to the nutrient medium. *Vibrio hydrosulfureus* is shown to be active in the same way. Beyerinck has also obtained a micro-organism which transforms sulphates into hydrogen sulphide; but Saltet finds that this *Bacillus desulfuricans* only partially reduces sulphates to sulphites or other oxygenated sulphur compounds which are then reduced by other bacteria to hydrogen sulphide. The cycle which sulphur compounds undergo in nature is thus completed; for in the previous chapter (Chapter XI.) the oxidation changes which sulphur underwent were briefly described.

Compounds of arsenic, in wall-papers especially, are

acted upon by moulds, and arsenical vapours, formerly thought to be arseniuretted hydrogen, though this was denied by some investigators, are evolved. Biginelli has recently shown that the vapours consist of diethylarsine. Arsenious oxide in small quantities is completely volatilised by *Penicillium brevicaulis*, which produces the arsenical odour even from 0.01 and often from 0.001 milligram of arsenic; this mould serves as a good biological test for arsenic.

These various investigations have been carried out by Gosio, Emmerling, Abel and Bittenberg, and by Galli-Valerio and Strzyzowski.

König has shown that tartaric acid is reduced to succinic acid by *Bacterium termo*, and Emmerling that malic acid is converted into succinic acid almost quantitatively by *Bacillus lactis aërogenes*.

Basile states that glucose is reduced to mannitol in certain Sicilian wines, and thus confirms Malbot, who thinks that it may be caused by the butyric ferment which liberates hydrogen. Gayon and Dubourg find that an enzyme can reduce lævulose to mannitol. This mannitol ferment decomposes the sugars not fermented by yeast, and thus affords a means of detecting lævulose in the presence of fermentable carbohydrates.

Certain colouring matters, *e.g.* indigo-carmin, are bleached by micro-organisms when they are cultivated in solutions to which the colours have been added; this reduction is said not to be due to the action of nascent hydrogen.

In the animal organism processes of reduction, as well as oxidation, also occur. Abelous and Gérard have made

extracts of various organs of the animal body, and have found that nitrates can be reduced to nitrites, butyric acid to butyric aldehyde, and methylene blue decolourised by these extracts; further, nitrobenzene can be reduced to aniline. Pozzi-Escot states that fibrin contains a reducing enzyme, or reductase; and that there is also one present in urine; he uses hydrogen peroxide for detecting their presence.

CHAPTER XIII.

NITRIFICATION AND DENITRIFICATION.

PROCESSES of oxidation and reduction are continually taking place in the soil: these are now known to be the result of the action of several micro-organisms, which are able to bring about these changes.

Formerly it was thought that the change of ammonium salts into nitrates in the soil was a purely chemical process, analogous to that which takes place when air and ammonia are heated together in the presence of spongy platinum; by the porosity of the soil, which mechanically contained the oxygen of the air, the change was supposed to take place—a kind of catalytic action, similar to that of platinum black. In 1877 it was pointed out by Schloesing and Müntz that the process took place very slowly at 0°C ., that it had an optimum temperature at 37°C . and that above 60°C . the change no longer took place. These results strongly suggested the action of micro-organisms, the above observers holding that only one of these was necessary. In 1890 researches were undertaken by several investigators, among whom may be mentioned Frankland, Warington, and especially Winogradsky.

From the results obtained, it is now known that the change takes place in two stages. In the first stage the ammonium salts are converted into nitrites by the action of the nitrous organisms, or nitrose bacteria; only two types of bacteria, of each of which there are several species, are able to bring about this oxidation, and they are known as *Nitrosomonas* (Europe, Asia, Africa) and *Nitrosococci* (South America and Australia). They have no further action upon nitrites which can be produced in solutions of asparagine, urea, etc. In the second stage nitrites are converted into nitrates by the action of the nitric organisms, or nitro-bacteria, of which there is only one genus; this process does not take place in the presence of ammonia, and not until all the ammonium salts are converted into nitrites. Stutzer and Hartleb state that 89–93 per cent. of the nitrogen as nitrites can be converted into nitrates by the nitrate bacteria.

Experiments by Omeliansky show that the nitrogen of organic compounds is not attacked, but must be first converted into ammonia; substituted amines, however, are attacked, according to Demoussy; monomethylamine sulphate is converted into ammonia, and then into nitrite and nitrate; trimethylamine is attacked more slowly; there is no intermediate formation of monomethylamine, the three methyl groups being all attacked at the same time. Aniline, pyridine, and quinoline are attacked still more slowly, and the oxidation takes place apparently with greater difficulty when the molecule is more complex.

These bacteria derive their carbon from carbon dioxide;

they will not grow on gelatine, nor on the ordinary culture media, but require a mineral solution for their cultivation.

The reverse process of reduction of these substances—called denitrification—also takes place; in some cases nitrogen can be liberated from both nitrates and nitrites, *e.g.* by *Bacterium denitrificans* II. (Burri and Stutzer); in other cases nitrates are converted only into nitrites, and Maasen states that eighty-five micro-organisms are able to convert potassium nitrate into potassium nitrite. Loew has observed reduction as far as ammonia.

Ampola and Ulpiani, from the results of their experiments on denitrification by means of *Bacteria denitrificans* V. and VI., conclude that the process can be expressed by the equation:—



Asparagine and nitromethane are not attacked, and ethyl nitrate only very slowly, and the authors conclude that denitrification only takes place with oxygen compounds of nitrogen, and in those which are electrolytes and therefore dissociated in solution. The various micro-organisms are also divided into three classes: (1) those which destroy nitrites, but not nitrates; (2) those which destroy nitrates, but not nitrites; (3) those which destroy both nitrites and nitrates. Denitrification was also found to take place more rapidly with salts in which the metal is more electropositive and of lower atomic weight.

In this way free nitrogen is liberated, but it can again be assimilated by another set of bacteria occurring in the roots of the leguminosæ, and converted into albuminous bodies.

The cycle of changes which nitrogen passes through is,

as we have seen, very complicated : albuminous substances are converted into urea by animals ; urea is converted into ammonium carbonate, which in turn becomes nitrite and nitrate ; finally nitrogen is liberated, and this is assimilated and converted into albuminous substances.

CHAPTER XIV.

VARIOUS CHANGES OCCURRING AS THE RESULT OF FERMENTATIONS.

LACTIC acid, which itself is a product of fermentation, can undergo two changes in the same way. The first of these—butyric acid fermentation—has been known, perhaps, as long as that of lactic acid, since it takes place at the same time. Special organisms, *e.g.* *Bacillus amylobacter*, are the cause of this fermentation, and the change of lactic acid into butyric acid can be expressed by the equation:—



Carbon dioxide and hydrogen are generally produced in the process, and the hydrogen may cause the reduction of other organic substances which are generally present during the fermentation.

Not only, however, can butyric acid be produced from lactic acid, but also from the substances which can give rise to lactic acid; these have already been enumerated, and, in addition, higher acids, such as malic, tartaric, citric, and mucic, can be fermented with the formation of butyric acid as a product. From mucic acid, butyric acid is only slowly formed, and it ought to be considered as a secondary product.

The homologues of butyric acid, *e.g.* propionic,


valerianic, and acetic acids, and also succinic acid, are often produced at the same time ; they are, perhaps, formed by the action of the nascent hydrogen.

Mannite, quercite, and glycerine are decomposed by various micro-organisms, with the production of butyric acid ; from glycerine, methylalcohol, acetic acid and traces of formic acid are also formed in the presence of calcium carbonate.

Schattenfroh and Grassberger have described two very similar micro-organisms which form lactic acid as well as butyric acid, with the evolution of carbon dioxide and hydrogen. One of these micro-organisms is mobile, and almost completely converts lactose into butyric acid ; dextrose, cane sugar, and starch are not so completely converted. The other micro-organism is non-mobile, and produces both butyric and lactic acids. The lactic acid formed is sometimes inactive and sometimes dextro-rotatory.

The second of these changes—propionic acid fermentation—takes place in the calcium salt of lactic acid ; the principal product is propionic acid, but acetic and valerianic acids have also been found. Strecker first noticed this change in 1854, but the most careful study has been made by Fitz, who has isolated the micro-organism which produces it.


The lactic acid splits up into two parts of propionic acid and one of acetic acid, together with carbon dioxide and water. No hydrogen is evolved, and it thus differs from butyric acid fermentation. This change can be represented by the equation :—




Malic acid, according to Fitz, is decomposed by another bacillus, with the production of propionic acid in rather larger quantities; butyric acid and alcohol are formed at the same time, but only in traces.

The production of butyl alcohol is another very important chemical change resulting from fermentation. It can be obtained from glycerol and mannitol by the action of the *Bacillus butylicus* obtained by Fitz; and Emmerling, who has investigated this change, finds that butyl alcohol is not obtained from dextrose, but that the products are ethyl alcohol and a solid fatty acid, probably palmitic acid. Beyerinck's *Granulobacter butylicum*, however, forms butyl alcohol from dextrose, but not from glycerol.

Calcium formate is converted into calcium carbonate and hydrogen by certain bacteria, though probably an enzyme is the cause of this change, since it continues after the micro-organisms have been destroyed. The following equation represents this change:—



Pakes and Jollyman have recently investigated the products obtained from sodium formate by the action of certain bacteria, and they express the reaction as follows:—



the total amount of carbon dioxide, evolved as gas and in combination, being equal in volume to the amount of hydrogen evolved.

Gonnermann has quite recently investigated the hydrolysis of acid amides and anilides by ferments; *e.g.* pepsin, trypsin, and emulsin, and by extracts of organs; and he finds that acetamide, acetanilide, formanilide, and others are hydrolysed; ptyalin, invertin, and maltase had no action.

E. Gérard has described the formation of creatine from creatinine by a kidney extract, and he refers to a similar dehydrating effect obtained by Abelous and Ribaut in the formation of hippuric acid by kidney extract.

Grimbert has obtained acetylmethylcarbinol from dextrose by the action of *Bacillus tartricus*; it was identified by its osazone and osotetrazone; it was also obtained from other sugars.

Various compounds have been obtained by the action of bacteria; *e.g.* esters by *Bacillus esterificans fluorescens* (Maasen); quinone by *Streptothrix chromogena* (Beyerinck). Went describes the formation of carbon disulphide by *Schizophyllum lobatum*.

In this chapter it will be convenient to shortly describe the colouring matters produced by bacteria.

The best known is that produced by *Bacillus cyano-genus*. The result of its action is the production of a blue pigment often seen in milk, especially in warm weather. Erdmann considered this colouring matter to belong to the aniline group, and it is probably triphenylrosaniline. The blue colouration in cheese is due to another micro-organism, named *Bacillus cyaneo-fuscus* by Beyerinck. This organism also gives rise to black glue.

The blue pigment formed by *Bacillus pyocyaneus* has been termed pyocyanine $C_{14}H_{14}N_2O$. It can be extracted

by chloroform, and was first obtained crystalline by Fordos. According to Boland, a reddish-brown colouring matter is found with the pyocyanine, which is insoluble in chloroform but can be extracted by alkalies; this is supposed to be formed from the pyocyanine in the culture of the bacillus. A green fluorescent colouring matter is produced by this microbe when it is cultivated in bouillon.

The oldest known chromogenic micro-organism is *Micrococcus prodigiosus* or *Bacterium prodigiosum*, which is the cause of the so-called "bleeding bread," "red snow," etc. The red pigment, termed prodigiosin, is insoluble in water, but is soluble in alcohol, chloroform, and carbon bisulphide. Griffiths gave it the formula $C_{38}H_{56}NO_5$, but Scheurlen has obtained analyses which do not agree with this. It has been suggested that this colouring matter is identical with fuchsine, but this statement has been contradicted by Helm and Bordoni-Uffreduzzi.

The red, yellow, and rose-coloured pigments were termed lipochromes by Zopf, the red being termed liporhodines, and the yellow lipoxanthines. They are converted by sulphuric acid into deep blue crystals of lipocyanine. They are closely related to the lipochromes of fat, yolk of egg, etc., and to the carotene of carrots.

A violet colouring matter is produced by *Bacterium violaceum*, and an indigo blue one by *Bacterium indigonaceum*. Lankester has named a bacterial colouring matter bacterio-purpurin, which pigment is also found in the sulphur bacteria.

Lastly, green fluorescent pigments are produced by

bacteria, *e.g.* by *Bacilli fluorescens liquefaciens* and *non-liquefaciens*: it has been termed bacterio-fluorescein. Jordan has found that phosphorus and sulphur are essential for its formation, and he has arranged a number of organic compounds in the order of their fluorescegenic value: asparagine, succinic, lactic, citric, tartaric acids, etc.; and he points out that the presence of a methylene group is favourable to fluorescence, but not coincident with dibasicity, nor with the presence of two methylene groups as asserted by Lepierre.

In conclusion, phosphorescence and heat can be produced by bacteria.

CHAPTER XV.

CHANGES IN BLOOD.

AMONG the albumins and allied substances, the transformation which blood undergoes when it is shed, is undoubtedly the change which has been known for the longest time. This change is commonly called coagulation, or "clotting," and it is the result of the action of a ferment.

When blood is drawn and allowed to stand, it quickly becomes gelatinous, but after some hours it is found to consist of a yellowish liquid containing a fibrinous substance, which is red in colour, owing to the mechanical entanglement in it of the red blood corpuscles. If diluted blood be looked at under the microscope, these corpuscles are seen to run together, forming rouleaux, and it was formerly thought that coagulation consisted in the running together of these bodies. In 1772 Hewson showed that these red blood corpuscles took no part in the process, and that the fibrinous substance was formed from the blood plasma, or *liquor sanguinis*. This observer isolated from the plasma "coagulable lymph," which is now termed "fibrinogen." Buchanan, in 1835, made several observations upon coagulation; he found that hydrocele fluid and pericardial fluid could not

coagulate of themselves, but if he added to them a solution which he obtained from lymphatic glands or from blood clot, coagulation at once set in. He compared this action to that of a ferment, and was therefore the first to suggest that this process was due to a ferment. Denis, in 1859, obtained by another method a substance which he called plasmine, and which on solution could undergo coagulation; this substance was no doubt identical with Hewson's coagulable lymph, or fibrinogen, as it is now called.

Much more complete investigations were carried out by Alexander Schmidt of Dorpat, who rediscovered the facts observed by Buchanan, which had been forgotten. From hydrocele fluid he prepared fibrinogen; and from serum, fibrinoplastin. Of themselves these two substances would not coagulate, but on adding one to the other coagulation resulted. He found that fibrinoplastin contained another substance, whose presence was necessary for the coagulation, but only in 1872 was he able to isolate this by precipitation with alcohol; he called it fibrin ferment, and considered that coagulation was caused by the action of fibrinoplastin on fibrinogen under the influence of fibrin ferment. Fibrinoplastin was subsequently named serum-globulin.

The next experiments were performed by Hammarsten, who showed that fibrin could be formed from fibrinogen and fibrin ferment without the presence of serum-globulin, which had no effect when pure, but contained a factor, when impure, which was necessary for the process; this could be replaced by impure casein or by calcium salts. The fibrinogen is split up during the change,

and does not combine with serum-globulin, as was formerly supposed.

Subsequently, Green found that fibrin ferment contained calcium sulphate, and that calcium sulphate was necessary for the change. Ringer and Sainsbury have shown that any calcium salt can replace the sulphate, and that barium and strontium salts can also be used. Experiments by Horne, in 1896, show that these salts, in the proportion of 0·5 per cent. and upwards, retard the coagulation, barium salts being the most powerful, and calcium salts the least; their influence is overcome by dilution with distilled water, or by the addition of a suitable amount of potassium oxalate; solutions of sodium and potassium chloride of 0·07 per cent. and upwards also retarded coagulation.

Calcium salts are therefore necessary for coagulation; this factor is often made use of when blood is wanted in an uncoagulated condition, by precipitating them as oxalates.

Fibrinogen and calcium salts will not coagulate by themselves, but fibrin ferment must also be present as a third factor, in order that coagulation may take place.

Fibrin ferment is a nucleoproteid; and Pekelharing and Halliburton, who have extracted nucleoproteids from various organs, have shown that these can cause the coagulation of fibrinogen in the presence of calcium salts. The former has therefore suggested that the nucleoproteid transfers calcium with which it combines to fibrinogen, whose calcium salt is insoluble, and constitutes fibrin; the nucleoproteid then combines with more calcium and transfers it again to fibrinogen. This

view is urged by Frederikse, who finds that the presence or absence of serum-globulin in such mixtures makes no difference in the amount of fibrin formed. He also confirms Hammarsten's statement, that fibrinogen is split up into fibrin or coagulated proteid and a soluble globulin by the action of fibrin ferment; the same decomposition is said to be brought about by the action of dilute acetic acid.

Lilienfeld regards the active agent in coagulation as a nucleo-albuminous substance, named nucleohiston, which arises from the leucocytes, especially from their nuclei, and from platelets. In this substance the nuclein hastens while the histon hinders the coagulation, the separation into nuclein and histon being effected by calcium salts. From fibrinogen he has obtained a substance, thrombosin, by precipitation with acetic acid and nuclein, or rather nucleic acid, which acts in the same way; thrombosin is coagulated by calcium salts, and the fibrin so formed is regarded as a calcium compound of thrombosin. Schäfer has criticised these assertions of Lilienfeld's, stating that further evidence is necessary to show that thrombosin and fibrinogen are different substances; also that Lilienfeld does not prove that nucleo-albumin does not enter into the reaction, for it is impossible to prepare fibrinogen quite free from nucleo-albumin.

Hammarsten and Cramer also consider that thrombosin is the same as fibrinogen. In the same publication, Hammarsten shows that calcium salts are not essential for the formation of fibrin from fibrinogen, since they both contain the same amount of calcium, and he confirms Pikelharing, that calcium salts act in the genesis


of fibrin ferment, *i.e.* in the change of prothrombin, the precursor of fibrin ferment, into thrombin, the fibrin ferment.

The formation of fibrin from fibrinogen is usually regarded as a process of hydrolysis, the fibrinogen being converted into fibrin and fibrinoglobulin (Arthus). Hammarsten has found that the elementary composition of these substances is very similar, and he brings forward the view that the change may consist in an intramolecular rearrangement in the fibrinogen, part of it appearing as fibrin, and part remaining in solution as fibrinoglobulin.

Under several other conditions, besides that of the removal of calcium salts, blood can be prevented from clotting, *e.g.* by leech extract, albumoses, peptones; and many experiments have been made in order to elucidate the manner in which fibrin is formed, but these do not throw any more light upon this important change.

The following facts, however, still remain proved: nucleoproteid (or prothrombin) combines with calcium to form fibrin ferment, which then acts upon fibrinogen, and causes the formation of fibrin, which constitutes the coagulum.

The facts can be tabulated thus:—



CHAPTER XVI.

CHANGES IN MILK, MUSCLE, AND IN THE LIQUID OF THE PROSTATE GLAND.

A TRANSFORMATION, very similar in many ways to the coagulation of blood, is carried out daily in the manufacture of cheese from milk. This change is produced by the action of rennin, a soluble ferment or enzyme contained in calves' stomachs, from which it can be extracted to form rennet.

Milk contains three proteids—caseinogen, lactalbumin, lactoglobulin, the two latter only in small quantities; also milk, sugar, fat, calcium phosphate, and other salts. Cheese is obtained from the caseinogen, which forms the curd, which contains the fat mechanically; the other proteids, the sugar, and the salts are contained in the liquid which constitutes the whey.

The action of the rennin was supposed in 1859, by Liebig, to be the conversion of lactose into lactic acid; this neutralised the alkali which kept the caseinogen in solution, thus causing its precipitation. This idea evidently arose from the fact that acids, when added to milk, produce a precipitation of the caseinogen. Soxhlet held a very similar view, but he took into account the presence of the calcium phosphate; this was converted by the lactic acid,


obtained from lactose by the action of rennin, into acid calcium phosphate, which precipitated the caseinogen.

The most important work upon the coagulation of milk is that of Hammarsten, who has shown that the lactose takes no part in the change, and that the clot can even be formed in an alkaline medium, as first pointed out by Selmi. The presence of calcium was found essential for the conversion, and Ringer states also that other calcium salts than the phosphate can be used.

The production of casein is a double process; in the first instance the caseinogen is converted into soluble casein by the rennin, and secondly the casein is precipitated by calcium, probably as calcium salt or caseate of lime (Halliburton).

Hammarsten considers that in the process the caseinogen is split into two parts—the insoluble casein and the whey proteid, which is soluble.

This change can be tabulated thus—




and is therefore slightly different from that which occurs in blood, where the calcium takes part in the first stage of the change.

The similarity between the formation of fibrin in the clotting of blood and the production of *rigor mortis* in muscle, was first pointed out by Kühne, and he made the first investigations upon this change, about the year 1860. From frog's muscle at a temperature of 0° C., he

obtained a plasma which possessed the power of coagulation; in very much the same way as in blood and milk; this clot contracted and gave a serum, but, instead of being fibrinous, this coagulum remained gelatinous and soft. The term "myosin" was given to it.

Halliburton has obtained the same results with plasma obtained from the muscles of rabbits and other animals, and he has shown that the myosin obtained, if redissolved after coagulation, could be made to clot even three or four times. He states that a ferment, which he has isolated, is the cause of this change, and he has called it myosin ferment. From the plasma he has obtained four proteids, which he has named paramyosinogen, myosinogen, myoglobulin, and myoalbumin; and he has shown that the coagulation is caused in the myosinogen, which carries down with it on coagulation the paramyosinogen. He considers that the myosinogen does not split up into two proteids, as is supposed to happen in the case of fibrinogen; in the coagulation, however, sarcolactic acid is formed and carbon dioxide is evolved, thus:—



Von Fürth has also carried out investigations upon the subject, and has obtained two proteids from muscle plasma:—(1) Paramyosinogen, constituting 17–22 per cent., and (2) myosinogen or myogen, constituting 77–83 per cent. of the plasma; this latter corresponds to Halliburton's myosinogen. These two proteids enter into the formation of the clot; paramyosinogen passes directly into myosin fibrin, but myosinogen or myogen passes

through an intermediate soluble stage into myogen fibrin, thus:—

Paramyosinogen —————→ myosin fibrin

Myosinogen → soluble myogen → myogen fibrin

He thus confirms Halliburton, that two proteids enter into the formation of the clot, and he regards Halliburton's myoglobulin as part of the myosinogen which is not coagulated.

Stewart and Sollman have obtained the same results, but they regard Von Fürth's soluble myogen as part of the paramyosinogen. Both the proteids are converted into insoluble myosin, the myosinogen passing first into a modification similar or identical to paramyosinogen, thus:—

Paramyosinogen —————→ myosin

Myosinogen → paramyosinogen → myosin

They confirm Halliburton also, that coagulation is accompanied by an increase of acidity.

Another case of coagulation is said by Camus and Gley to take place in the fluid of the vesiculæ seminales, which is somewhat viscid. If a drop of the liquid from the prostate gland of the same animal, guinea-pig, rat, or hedgehog, is added to the first liquid, coagulation sets in. This change is stated by the observers to be due to an enzyme which they have named vesiculase, on account of the fact that the property is lost when the prostatic liquid is heated to 70° C. for fifteen minutes. Beyond these observations nothing further is known of this transformation.

CHAPTER XVII.

CHANGES OCCURRING IN ALBUMINS AS THE RESULT OF THE ACTION OF PEPSIN.

By far the most important changes which albumins and allied substances undergo as the result of fermentation are those produced by the digestive ferments of the animal organism, pepsin and trypsin. In plants similar ferments, bromelin and papain, have been found, which resemble trypsin in their action very closely.

The change which takes place in the albumins is known as proteolysis, and it consists in a breaking up of the very complex albumin molecule into smaller ones, the size and constitution of some of which, especially those obtained from trypsin digestion, are known. Since some of these later changes are known to take place by hydrolysis, it is very probable that the complex molecule is broken up in a similar way; especially so as the same results can be obtained with mineral acids or superheated steam. The changes are thus quite comparable to those produced in the carbohydrates, smaller molecules being obtained at each stage; here, however, the end-products are very much more various, and consist chiefly of amino-acids.


In the animal organism, albumins taken as food are first acted upon by the gastric juice of the stomach, which contains pepsin; and it is therefore most convenient to describe the changes produced by this ferment before going on to those produced by trypsin.

Pepsin, in order to produce proteolytic changes, must always be accompanied by an acid; in the gastric juice hydrochloric acid is always present in about 0·2–0·5 per cent. In this connection, observations have of late years been made by Mayer, Hahn, and Wroblewsky in particular, upon which acids are the most favourable to peptic action; and it has been found that oxalic acid produces the most speedy solution of the albumin. Next in order comes hydrochloric acid, then nitric, phosphoric, tartaric, lactic, citric, malic, sarcolactic or d-lactic, sulphuric, and acetic acids, in the order mentioned. A dilution of 0·01–0·5 per cent. was the most favourable; pepsin will not act in an alkaline solution, and even neutral salts retard its action.

Very many investigations have been made upon the products of the action of pepsin upon albumins, fibrin and egg-albumin especially. The earliest were those of Meissner and his pupils, in 1859–62, who obtained the following substances: parapeptone, metapeptone, dyspeptone, α -peptone, β -peptone, γ -peptone. Brücke, who made a determination of the products about the same time, could, however, only obtain two products, which corresponded to parapeptone and γ -peptone.


Kühne, some years later, studied the action both of pepsin and trypsin, and found that the only real difference was that trypsin produced a more complete hydrolysis.

He obtained, by using pepsin, albumose and peptone; this he called amphopeptone on account of the fact that trypsin only converts half the peptone further into amino-acids. He considered the albumin molecule to be split up into two halves, antialbumose and hemialbumose, and devised the following diagram to represent these changes:—




Fränkel and Langstein have recently shown that Kühne's amphopeptone consists of two substances; (1) a peptone; (2) a substance identical with the carbohydrate complex of albumin—albamin.

Chittenden, Neumeister, and others have somewhat recently studied the products of pepsin digestion very carefully, and they have come to the following conclusions: fibrin or egg-albumin is converted firstly into syntonin, chiefly by the action of the acid, and then this is split up into two primary albumoses, which are called protoalbumose and heteroalbumose. Heteroalbumose, on standing under water, or on drying, is converted into dysalbumose, which can be reconverted by acids into heteroalbumose. These two albumoses then give deuteroalbumose, and subsequently peptone. The diagram shows these results more plainly:—



Some interesting conclusions from the results of molecular weight determinations of peptone and protoalbumose by Sabanejeff as given by Neumeister seem appropriate here. Peptone was found to possess a molecular weight of about 400, and protoalbumose of 2400; thus on decomposition six peptone molecules would be obtained from one protoalbumose. Taking the albumin molecule as 15,000, then albumin would give altogether 40, and heteroalbumose would give 34 molecules of peptone.

The other albumins, when subjected to pepsin digestion, give very similar products; these may be tabulated as follows:—



Chittenden and Mendel, who have investigated the changes produced in this substance, find that there is a gradual falling off of carbon in the percentage composition of these substances formed in the digestion; they suppose

that the four substances above named are only the main steps in the hydration process, and that there are still other links in the chain which cannot as yet be separated, and which show the same general chemical behaviour.


Globulin $\rightarrow$ globuloses $\rightarrow$ peptone

Egg-albumin $\rightarrow$ ovalbumoses $\rightarrow$ peptone


Myosin $\rightarrow$ myosinoses $\rightarrow$ peptone

The compound proteids are firstly split up into their constituent parts, one of which only undergoes peptic transformation.

The behaviour of casein in peptic digestion, especially with regard to the phosphorus, has been studied carefully by Salkowsky; though frequently stated that the phosphorus is not dissolved, but is separated as insoluble paranuclein, he finds that much of it is dissolved, only about 15 per cent. being contained in the paranuclein; the more favourable the conditions of digestion, the less paranuclein is obtained, and paranucleic acid is obtained from the paranuclein. The phosphorus in solution is in the form of a phosphorised albumose, in which it is not combined, however, as phosphoric acid; if dry hard lumps of casein are avoided, it is completely dissolved, and this is done by making a solution of casein before adding the digestive fluid, which acts best in the proportion of 500 to 1 of casein. Lindemann confirms this, and states that caseinogen is more digestible than casein. The first product of the digestion of casein is caseoproteose, which contains all the phosphorus of the original proteid; paranucleic acid, together with paranuclein, are formed in the digestion, and Salkowski has prepared the iron salt of this acid; thus:—



The decomposition of the albuminoids has been carefully studied by Chittenden, with the following results:—



No peptone is formed from elastin, and keratin is not attacked at all.

Umber has obtained from crystallised egg-albumin and serum-albumin, the primary proteoses, three deutero-proteoses, A, B, C, and peptone; and Zunz finds, as primary products of peptic proteolysis, acid albumin, proto- and heteroproteose, and deuteroproteose B; as secondary products three other deuteroproteoses, two peptones, and unknown substances which do not give the biuret reaction, ammonia and amino-substances.

Pick has investigated the proto- and hetero- albumose obtained from fibrin, and finds that they differ from one another in solubility and in the amount of nitrogen in a basic form which they contain, heteroalbumose containing 39 per cent. and protoalbumose 25 per cent.; hetero-albumose yields a large quantity of leucine and glycocoll,

whereas protoalbumose yields only a little leucine and no glycocoll. They arise together from fibrin, but not from one another, and they both yield, on further peptic digestion, deutoalbumoses A and B and peptone B. This confirms the results of Zunz.

As regards the nature of the end-products of gastric digestion, a great deal of work has lately been done. Pfaundler obtained from serum-albumin leucine and a diamino-acid, probably histidine, but from fibrin there was no leucine; he finds that the principal end-products are apparently substances intermediate between peptone and amino-acids; they do not give the biuret reaction, and contain more than one carbon nucleus, in the case of serum-albumin at least there being a leucine and diamino-nucleus present. Langstein has obtained from crystallised egg-albumin the following substances: leucine, tyrosine, glutamic acid, aspartic acid, cystine, lysine (in small amount), pentamethylenediamine, hydroxyphenylethylamine, and "a polymeric carbohydrate containing nitrogen;" also a base giving skatole and two unknown acids which give the biuret reaction; he did not find the intermediate substances of Pfaundler, perhaps because the digestion had gone too far. Malfatti has found that tryptophane is formed by the prolonged action of pepsin-hydrochloric acid on Witte's peptone, and also small quantities of leucine, tyrosine, and hexone bases.

V. Harlay has found a very interesting substance among the products of pepsin digestion: it is a chromogen, as it gives, on treatment with the enzyme tyrosinase, a red colour; on decomposition with trypsin it yields tyrosine and tryptophane, a body first observed by Neumeister

among the products of trypsin digestion. According to Lawroff, pepsin breaks up proteids as much as trypsin does.

Sawjaloff has recently advanced the theory that peptonisation of proteid matter is not to make it more easy for absorption, as there is a reverse process of building up of proteids of larger molecular weight from the products of peptonisation, and this action is attributed to the rennet ferment; a substance termed plastein is thus synthesised, which appears to be the same, whatever the proteid may be which is digested. It resembles Kühne's anti-albumid in properties; Krüger's results, that the digestive products inhibit the action of pepsin, help to support this view.


CHAPTER XVIII.

CHANGES OCCURRING IN ALBUMINS AS THE RESULT OF THE ACTION OF TRYPSIN.

CLAUDE BERNARD, and in 1857 Corvisart, who refers to Purkinje and Pappenheim as having made experiments with pancreatic juice in 1836, were really the first to begin the scientific study of the digestion of albumins with pancreatic juice. The former considered bile to be necessary, but Corvisart showed that the juice by itself could dissolve albumins; and that they were converted into substances similar to those of gastric digestion. These results were at first denied, but later they were confirmed by Meissner, Danilewsky, and Kühne; this last observer showed that the action was due to an enzyme to which he gave the name of trypsin. Until Kolbe discovered that disinfectants could prevent the action of those micro-organisms which are always present in the intestine, the products of digestion by trypsin were very much confused and complicated; Kühne, in his experiments, made use of this fact, and was therefore the first to investigate the real products of trypsin digestion.

Kühne discovered that the peptone which is formed is only partially further broken up by trypsin, and he gave

the name of amphopeptone to the total quantity, anti-peptone to that part not attacked, and hemipeptone to that part which was further broken down. He surmised that the splitting of the albumin molecule took place earlier in the process, and he made attempts to isolate intermediate substances, which he called amphoalbumoses. That part of the albumin not attacked was called antialbumid; very strong trypsin converts it into anti-peptone, as was observed by Schützenberger in 1875. It is really, however, converted into antialbumose or deuteroalbumose, and Kühne gave the following diagram to show these results :—



Now, however, it is known that no antialbumose is formed, but only antipeptone.

The characteristic changes produced by trypsin have been more carefully studied with fibrin. As with pepsin, this substance has also the property of taking up the ferment, and a method of obtaining trypsin in a relatively pure condition is thus obtained.

The fibrin first dissolves, but does not swell up as it does in peptic digestion, being rather eroded from the outside. The solution has the property of a globulin, but no globulin is formed from serum-albumin, etc., also no syntonin nor a product corresponding to that formed in peptic digestion. Primary albumoses are not formed, but

deuteroalbumose is obtained directly, then amphopeptone, and finally antipeptone and hemipeptone, of which the former constitutes rather more than half of the total quantity of the peptone. The further action of trypsin consists in the conversion of hemipeptone only into amino-acids, and a chromogen, which was first observed by Tiedemann and Gmelin in 1831, and called tryptophane by Neumeister. This chromogen forms a very suitable substance for determining the progress of tryptic digestion.

According to Nencki, tryptophane, called proteinchromogen by Stadelmann, can be brominated, when it gives rise to two substances. These, on removal of the bromine, have similarities on the one hand to hæmatoporphyrin and to bilirubin, and on the other hand to melanin; he therefore considers them the mother substances of the animal colouring matters, even of hæmoglobin.

Hopkins and Cole have quite recently isolated tryptophane, and have given it the formula $C_{11}H_{12}O_2N_2$; it gives the Adamkiewicz reaction, and yields skatole and indole when heated; it is probably indoleaminopropionic acid or skatoleaminoacetic acid.

The chief acids obtained are leucine, tyrosine or p-oxyphenylaminopropionic acid, aspartic acid, and glutamic acid; besides these, Drechsel and Hedin have found lysine or α , ϵ -diaminocaproic acid, and lysatinine, which is now known to be a mixture of lysine and arginine or δ -guanidine- α -aminovalerianic acid; Kutscher has obtained histidine, $C_6H_9O_2N_3$.

Glycocoll, long known as a decomposition product of gelatine, has been found by Spiro among the decomposition products of albumins.


Emerson has obtained p-hydroxyphenylethylamine, which is formed from the tyrosine, which loses carbon dioxide, its amount decreasing as the quantity of this substance increases.

Ammonia, according to Hirschler and Stadelmann, is given off in small quantities when trypsin acts upon albumins. Its amount has been determined by Dziergowski and Salaskin, who obtained it both in peptic and tryptic digestion. For the different albumins its amount varies, but in every case it is greater than that liberated by dilute acid or alkali, and it never corresponds to the total amino-nitrogen of the albumin.

Lea states that the removal of the products has an influence upon the digestion by trypsin, which is more rapid and complete than that of pepsin.

Nencki has stated that he has obtained xanthine, guanine, hypoxanthine, and adenine in the digestion of fibrin by trypsin; but Kossel denies this, saying that they are products of nucleins resulting from the entanglement of leucocytes in the fibrin.

The following diagram shows the formation of the principal products:—



Lately, investigations have been carried out as to the nature of antipeptone. Siegfried and Balke considered it to be a chemical individual, and the former investigator regarded it as identical with his carnic acid, $C_{10}H_{15}O_5N_3$. Kutscher, on the other hand, maintains that antipeptone is not a chemical entity, because he has obtained from it a mixture of substances which can be separated by phosphotungstic acid into an acid part and a basic part; the basic part consists of histidine, arginine, and another unknown base, and from the acid part aspartic acid was obtained. Siegfried states that antipeptone can be obtained pure from impure antipeptone by precipitation with alcohol, and he has since isolated two acids from antipeptone which he designates α -antipeptone, $C_{10}H_{17}O_5N_3$, and β -antipeptone, $C_{11}H_{19}O_5N_3$; they were not equal in amount, and they do not give quite the same products on hydrolysis with stannous chloride and hydrochloric acid; neither of them is identical with his carnic acid; he thus shows that antipeptone is not a mixture of substances, as stated by Kutscher, though his two acids can give those substances on hydrolysis.

The proteids are generally quickly decomposed by trypsin into their components, and the albumin digested like fibrin; *e.g.* casein is almost completely digested, giving tyrosine, casein albumose, and casein antipeptone. Biffi, who investigated this decomposition, found that the phosphorus was partially converted into phosphoric acid, and partially into another compound not precipitated by magnesia mixture.

The nuclein of nucleoproteids is peptonised with the separation of nucleic acids, and then of nuclein bases,

which Bokay states resist the action of trypsin, though Kutscher, as above mentioned, found xanthine, hypoxanthine, and guanine.

Of the albuminoids, only collagen and elastin are attacked by trypsin; the native collagen, however, is only changed when it has been first transformed into gelatine, or gluten, by the action of steam or acids, by the latter of which it is changed in peptic digestion.

Chittenden and Solley state that gelatine is converted into protogelatose and deuterogelatose, and that no gelatine peptone is formed when putrefactive changes are very carefully prevented from occurring. Chittenden and Hart state that no peptone is formed from elastin, but that protoelastose and deuterogelatose are both formed.

From gelatine Nencki states that he has obtained gelatine peptone, leucine, glycocoll, and ammonia; and from elastin Wälchi has obtained glycocoll, leucine, aminovalerianic acid, and ammonia.

The lecithines are also decomposed by pancreatic juice like fats, and are converted into their constituents, viz. choline, glycerophosphoric acid, and fatty acid. The choline, according to Hasebroek, is converted into carbon dioxide, marsh gas, and ammonia, the same products being also produced by bacteria when no oxygen is allowed to be present.

Kossel and Matthews have investigated the action of pepsin and trypsin on protamines (salmine and sturine), and have found that they are only attacked by trypsin, which converts them firstly into protones, and then partially only into hexone bases (lysine, arginine, and

histidine); the existence of an anti-group is said by them to be present on account of this fact, and also an intermediate stage between protones and hexones is probable.

Gulewitsch finds that only p-diacetylamidophenol, among many other less complex chemical compounds, is acted upon by trypsin, with the formation of acetic acid. He cannot confirm Nencki and Blank, who stated that hippuric acid was decomposed, and considers that their trypsin contained the fat-splitting ferment of the pancreas.

Baldwin and Levene find that the diphtheria and tetanus toxines are digested by trypsin, and thus rendered inert; the activity of tuberculine is only destroyed by the prolonged action of trypsin.

According to Cohnheim, the passage of peptone through the intestinal wall depends on its further decomposition into simpler compounds caused by a ferment which he has termed erepsin; it acts rapidly on proteoses and peptones, producing ammonia, leucine, tyrosine, lysine, histidine, arginine; it acts feebly upon histones, and not upon proteids except caseinogen; protamines are, however, acted upon.

Matthews even suggests that the centrosome in the unfertilized eggs of *Arbacia* plays the part of a proteolytic enzyme, as he has observed localized areas of liquefaction in the protoplasm.

CHAPTER XIX.

PROTEOLYSIS BY FERMENTS OTHER THAN PEPSIN AND TRYPSIN.

A PROTEOLYTIC ferment has been discovered by Babcock and Russell in the ripening of cheese, which they have named galactase. It has an action like trypsin, alkaline media being the best, but it is not identical with it, and it is found in the milk of all animals. Albumoses, peptones, amino-acids, and ammonia are formed, these nitrogen compounds being fairly evenly divided. It readily decomposes hydrogen peroxide, and it liquefies gelatine. These results have been confirmed by von Freudenreich.

A phenomenon, first described by Salkowski and termed autodigestion, occurs in the organs after death; it is not due to putrefaction, as it takes place in the presence of chloroform and sodium fluoride. Biondi has confirmed Salkowski in his observations, and shows that the enzyme which produces this action is not trypsin, as considered by Neumeister, in the following way: (1) xanthine-like substances are formed which in tryptic digestion are present in a latent condition; (2) autodigestion is not inhibited by an acid medium; (3) tryptophane is not a product, whereas it occurs in tryptic digestions.

Magnus-Levy has obtained i-lactic acid, d-lactic acid, and succinic acid; also formic, acetic, and butyric acids, hydrogen, carbon dioxide, and hydrogen sulphide from livers which were subjected to autolysis; further, glycocoll was obtained by Jacoby in autodigestion of the liver.

In yeast there is also a proteolytic enzyme, as Hahn found that an appreciable quantity of gelatine was dissolved in twenty-four hours by it, and he thus confirms Will as regards its presence in yeast; in conjunction with Geret he has shown that the albumin in yeast extract is decomposed with the formation of tyrosine and leucine. These investigators have further shown that it can digest added albumin in addition to that in the extract, with the formation of the same products; hypoxanthine is also found, its amount increasing during the digestion. They have termed this enzyme endotrypsin, and it resembles pepsin in requiring an acid medium, and trypsin in the nature of its products. In addition to the above products, Kutscher has isolated histidine, arginine, lysine, aspartic acid, and a compound $C_8H_6O_4N_4$.

A similar ferment has also been found in *Aspergillus niger* and other fungi by Bourquelot and Hérissé; this enzyme of *Aspergillus niger* has been found by Malfitano to act upon gelatine, nucleo-proteids, globulins, and albuminates, but not on egg-albumin. Emmerling and Reiser have investigated the action of *Bacillus fluorescens liquefaciens* on gelatine. Twenty-five per cent. of the nitrogen was eliminated as ammonia; and methylamine, trimethylamine, choline, and betaine were identified in the solution. Fibrin gave peptone, arginine, leucine and aspartic acid when acted upon by this micro-organism.

Peptonising bacteria are also found in milk which produce an enzyme very similar to trypsin; in the fermentation of casein aromatic acids are formed as well as peptone, leucine, and tyrosine; and cane sugar can be inverted, but not lactose (Kalischer).

Proteolytic enzymes are also probably the cause of bacteriolysis, the term used for their autodigestion.

The vegetable ferment bromelin was discovered by Marcano in 1891, and the products of its action have been examined by Chittenden and his pupils. Except for a small residue, fibrin is completely dissolved, and is converted into the various albumoses and peptones, and finally into amino-acids. The small residue is called anti-albumid. The other albumins behave similarly.

Papain, the proteolytic enzyme of the papaw tree, was first investigated by Wurtz in 1879, who said it converted albumins into peptones and leucine. Then, in 1883 and 1884, Martin made a complete examination, and found a milk-curdling ferment as well as a proteolytic one in the juice of this tree. The products of the digestion of fibrin were a globulin-like body, peptone, leucine, and tyrosine, more of the former than of the latter being formed.

Vegetable proteids are converted in a similar way, a small anti-albumid residue being left. Halliburton states that no peptone is formed from vegetable albumins by this enzyme, the process stopping short at the proteoses.

Harlay considers it intermediate in power between pepsin and trypsin, and he has obtained the same chromogen by its use as he obtained with pepsin.


Chittenden, who made very complete observations,

states that the nitrogen in the albumoses produced by vegetable ferments is less than in those produced by trypsin.

Emmerling has quite recently published the results he has obtained by the digestion of fibrin with papain; the fibrin dissolved giving peptone, which, by prolonged action, was converted into tyrosine, arginine, glycocoll, alanine, leucine, aspartic acid, and phenylalanine. Mendel and Underhill find that papain can act in both alkaline and slightly acid media, giving products corresponding closely to those obtained with pepsin; leucine, tyrosine, and tryptophane do not occur if bacteria are excluded; bromelin, however, readily forms leucine, tyrosine, and tryptophane, even in acid media.

A proteolytic ferment is also contained in the germinating seeds of the Lupin and other plants; it acts upon conglutin, giving leucine and tyrosine, but not asparagine, which is not regarded by Schulze as a primary product of the decomposition of albumin; Bokorny has observed the same decomposition by malt enzymes.

A theory to account for the action of these enzymes, especially papain, was published by Wurtz in 1881; he has compared the process with that which takes place in the formation of ether from alcohol and sulphuric acid. In this process, ethylsulphuric acid is first formed; this is then acted upon by alcohol, giving ether and sulphuric acid; a small quantity of sulphuric acid can thus transform large quantities of alcohol into ether. The enzyme is supposed to have the property of combining with the albumin, forming a compound which is decomposed by a further quantity of albumin, thus:—



This theory evidently arose from the fact that pepsin, trypsin, etc., can be taken up by fibrin as before mentioned.

CHAPTER XX.

THE CHANGES AND PRODUCTS OCCURRING AS THE RESULT OF PUTREFACTION.

THE last of the changes resulting from fermentation is known as putrefaction; here several different changes are without doubt taking place, but only in a very few instances have they been definitely determined.

Putrefaction is a process which has been known from early times to take place in organic substances when they are exposed to the air, and it was considered by the earliest observers to be a kind of fermentation, similar to the production of alcohol from sugar solutions. It was Pasteur, again, who showed that micro-organisms were the cause of these changes, and he found that air was not necessary for the majority of the organisms producing putrefaction, which are now known to be anærobic; it is now thought to be very probable that enzymes are secreted by those organisms which produce many of these changes.

Analogous processes also take place in the intestines of animals both normally and abnormally, and the products resulting are very similar to those occurring in putrefaction; it is, therefore, convenient to describe the changes occurring in the intestines and in the air together.

The first stages in the decomposition of albumins consist in their solution and conversion into peptone; this, according to Salkowski, is an enzyme action; but Kühne has denied that this enzyme is trypsin. The quantity of albumoses and peptones obtained is always small, as they are immediately further broken down. In this connection Neumeister has made some very interesting experiments; he finds that native albumins are not attacked until peptone is no longer present, or until it is completely removed from the mass of decomposing matter; but gelatine is an exception, as he has found peptone to be formed from it under the same conditions of putrefaction, the quantity increasing instead of decreasing as in the former cases. In the intestine it is found that bacteria are not necessary for digestion, but also that they do no harm (Nuttall and Thierfelder).

In tryptic digestion, the aromatic radicles in the albumin molecule yield chiefly tyrosine, but in putrefaction various aromatic bodies are formed. These are produced partly by the action of bacteria on the tyrosine, and partly by their action on the albumin directly.


The chief aromatic compounds obtained by putrefaction are :—

(a) Tyrosine and its derivatives—


1. Hydroparacumaric acid, or p-oxyphenylpropionic acid.
2. p-oxyphenylacetic acid.
3. Phenylpropionic acid.
4. Phenylacetic acid.
5. p-cresol.
6. Phenol.

According to Baumann, these are produced from tyrosine in the following way:—


1. Reduction of the tyrosine with nascent hydrogen—




2. Oxidation of the hydroparacumaric acid—



3. Splitting off of carbon dioxide—



4. Oxidation of the p-cresol—



- (b) Those produced from albumin directly—

1. Indole (Nencki and Kühne).
2. Skatole.
3. Skatole-carboxylic acid (Salkowski).
4. Skatole-acetic acid (Nencki, Salkowski).

These products, according to Baumann, are not primary products of bacterial action on albumin, but they are formed in the decomposition of an intermediate substance which is not known.

From these products it is seen that there must be

several aromatic nuclei in the albumin molecule, the majority of which are split off as tyrosine by fermentation.

From the fatty series no aromatic body as yet has been found as a product of bacterial action, although this would, according to Neumeister, be quite possible, since it is not known why bacteria give indole and skatole, and other decomposing agents only tyrosine, and *Proteus vulgaris* practically only indole. But the selective action of these organisms, on the other hand, as has been shown in several other cases, may be sufficient to account for these results.

In the body these various products combine chiefly with sulphuric acid, and are excreted as sulphuric acid esters; hippuric acid is formed from phenylpropionic acid and glycocoll, the phenylpropionic acid first undergoing oxidation; phenaceturic acid $\text{C}_6\text{H}_5\text{—CH}_2\text{CO—NH—CH}_2\text{COOH}$ is formed from phenylacetic acid and glycocoll.

The aliphatic products of intestinal putrefaction are leucine, aspartic, and glutamic acids, also θ -amino-valerianic acid (Salkowski); the ammonium salts of caproic, valerianic, and butyric acids; marsh gas and hydrogen; in addition to these succinic and glutaric acids, probably from the aspartic and glutamic acids. From the sulphur in the albumin molecule, sulphuretted hydrogen and methylmercaptan are formed.

Elastin and gelatine give the same products, but in addition glycocoll.

These changes are very probably hydrolytic in character.

Besides these products, outside the organism certain basic substances are formed, which in their reactions


closely resemble the vegetable alkaloids; they belong chiefly to the fatty series, whereas the true plant alkaloids are pyridine derivatives. In the living body these bases are seldom formed, oxygen being necessary for their formation. Dupré and Bence Jones in 1866, and Selmi in 1873, who suggested the name of Ptomaine for the whole group, first hinted at their existence, but they were unable to isolate them.

The recognition of these substances has been chiefly studied by Brieger, who has isolated the following bases:—

1. Monamines: methylamine, ethylamine, dimethylamine, diethylamine, trimethylamine, triethylamine.
2. Diamines: putrescine, cadaverine.

A diamine of the constitution $C_6H_{16}N_2$, which is probably hexamethylenediamine, was obtained by Garcia in 1893 from putrefying horseflesh; also saprine, $C_5H_{16}N_2$, and neuridine, $C_5H_{14}N_2$ (Ehrenberg).

According to Baumann and Udransky, the diamines are produced synthetically from monamines, thus:—



Collidine, $C_8H_{11}N$, has been obtained by Nencki from gelatine; and hydrocollidine, $C_8H_{15}N$, and parvoline, $C_9H_{13}N$, are generally produced from horse and cattle flesh which are undergoing putrefaction.


Choline also is found, which most probably originates from lecithine, and from which, perhaps, dimethyl-, trimethyl-, and triethylamine are formed.

Mydatoxine and mydine have been obtained by Brieger from putrefying human corpses.

All these substances are not poisonous, or only poisonous in large doses, and for the following bases, which are very poisonous, Brieger suggested the name Toxine.

1. Peptotoxine, found by Brieger in many peptones.
2. Neurine, $C_5H_{14}NO$, found in putrefying flesh.
3. Muscarine, $C_5H_{13}NO_2$, found by Brieger in putrefying fish; the poison of the fly agaric.

These two bodies very probably originate from choline, thus:—



4. A base, $C_2H_4(NH_2)_2$, isomeric with ethylenediamine, found in putrefying fish.


5. Mydaleine, from human corpses (Brieger).

6. Mytilotoxine, $C_6H_{15}NO_2$, obtained by Brieger from edible mussels.

7. Tyrotoxicone, found in milk and cheese by Vaughan.

8. Tyrotoxine, found in cheese by Dokkum.

9. Methylguanidine, obtained by Brieger from putrefying horseflesh; it is probably produced by oxidation from the creatine of muscles, thus:—



The creatinine group of compounds (xanthocreatinine, $C_5H_{10}N_4O$, crusocreatinine, $C_5H_8N_4O$, and amphocreatinine,

$C_9H_{19}N_7O_4$) and the uric acid group of compounds (adenine, xanthine, hypoxanthine) were called Leucomaines by Gautier in 1888; they are poisons produced from lecithine, or proteids, by the metabolic activity of the body.

Brieger has also investigated the poisonous substances produced in certain diseases; in cystinuria, cystine is formed in the organism; and in cholera, typhoid, tetanus, etc., several poisonous substances are formed. Using pure cultures of the micro-organisms of these latter diseases, Brieger isolated from cholera, putrescine, cadaverine, methylguanidine, indole, and nitrites; from typhoid, typhotoxine, $C_7H_{17}NO_2$, and the non-poisonous neuridine; and from tetanus, tetanine, $C_{13}H_{22}N_2O_4$, putrescine, tetanotoxine, spasmotoxine, and an unknown base.

Oméliansky has found that their toxins can be produced from amino-acids by the bacteria of cholera and diphtheria.

The real poisons in these diseases, however, are bodies of albumose properties, the constitution of which is not yet known; similar bodies constitute the deadly poisons of snakes, fishes, spiders, insects, plants, from which, *e.g.*, abrine, ricine, lupinotoxine have been obtained.

THE principal deductions from the mass of work on the chemical changes occurring in fermentation, which has been summarised above are as follows:—

1. That the active agent in all cases is a living organism.

2. That this organism sets up either by its own metabolism, or by bodies secreted by, or excreted from it, changes in substances of different constitution, *e.g.*, carbohydrates and albumins.

3. That these changes are principally hydrolytic, and that some are oxidative, and a few reductive.

4. That the results of these changes are generally simplifications of the original body.

5. In certain cases synthesis occurs.

LONDON, *October*, 1902.

BIBLIOGRAPHY

- R. Abel und P. Buttenberg.** Ueber die Einwirkung von Schimmelpilzen auf Arsen und seine Verbindungen. (Chem. Centr., 1900, i, 428.)
- J. E. Abelous et G. Biarnes.** Sur le pouvoir oxydant du sang. (Compt. rend. Soc. Biol., 1894, 536.)
- J. E. Abelous et G. Biarnes.** Sur le pouvoir oxydant du sang et des organes. (Compt. rend. Soc. Biol., 1894, 799.)
- J. E. Abelous et G. Biarnes.** Mécanisme d'oxydation organique. (Compt. rend. Soc. Biol., 1896, [x], 3, 94.)
- J. E. Abelous et E. Gérard.** Sur la présence, dans l'organisme animal, d'un ferment soluble réduisant les nitrates. (Compt. rend., 1899, 129, 56.)
- J. E. Abelous et E. Gérard.** Sur la présence, dans l'organisme animal, d'un ferment soluble réducteur. Pouvoir réducteur des extraits d'organes. (Compt. rend., 1899, 129, 164.)
- J. E. Abelous et E. Gérard.** Sur la coexistence d'une diastase réductrice et d'une diastase oxydante dans les organes animaux. (Compt. rend., 1899, 129, 1023.)
- J. E. Abelous et E. Gérard.** Transformation de la nitrobenzine en phénylamine ou aniline par un ferment réducteur et hydrogénant de l'organisme. (Compt. rend., 1900, 130, 420.)
- G. Ampola and C. Ulpiani.** Sulla denitrificazione. (Gazzetta, 1898, 28, i, 410.)
- G. Ampola and C. Ulpiani.** Sull'azione reductrice dei batteri denitrificanti. (Gazzetta, 1899, 29, i, 49.)
- H. E. Armstrong.** Presidential Address. (Journ. Chem. Soc., 1895, 1136.)
- E. F. Armstrong.** The Synthetical Action of Enzymes. (British Association, 1902.)

- M. Arthus.** Fibrinogène et Fibrine. (Compt. rend. Soc. Biol., 1894, 306.)
- M. Arthus.** Glycolyse dans le sang. (Compt. rend., 1892, 114, 605.)
- M. Arthus et Pagès.** Nouvelle theorie chimique de la coagulation du sang. (Arch. de physiol. norm. et path., 1890, [v], 2, 739.)
- S. M. Babcock and H. L. Russell.** Unorganised Ferments in Milk: a New Factor in the Ripening of Cheese. (14th Ann. Rep. Agr. Expt. Stat. Univ. Wisconsin, 1897, 161.)
- S. M. Babcock, H. L. Russell, and A. Vivian.** Properties of Galactase: a Digestive Ferment of Milk. (15th Ann. Rep. Agr. Expt. Stat. Univ. Wisconsin, 1897-98, 77.)
- S. M. Babcock, H. L. Russell, and A. Vivian.** Distribution of Galactase. (15th Ann. Rep. Agr. Expt. Stat. Univ. Wisconsin, 1898, 87, 93.)
- S. M. Babcock, H. L. Russell, A. Vivian, and Hastings.** The Action of Galactase and other Proteolytic Ferments on Milk and Cheese. (16th Ann. Rep. Agr. Expt. Stat. Univ. Wisconsin, 1899.)
- S. M. Babcock and H. L. Russell.** Unorganised Ferments of Milk: a new Factor in the Ripening of Cheese. (Centr. Bakt. Par., 1897, [ii], 3, 615.)
- S. M. Babcock and H. L. Russell.** Galaktase, das der Milch eigentümliche proteolytische Ferment, seine Eigenschaften und seine Wirkung auf die Proteide der Milch. (Centr. Bakt. Par., 1900, [ii], 6, 17, 45, 79.)
- Adolf Baeyer.** Ueber die Wasserentziehung und Ihre Bedeutung für das Pflanzenleben und die Gährung. (Ber. d. deutsch. chem. Ges., 1870, 3, 63.)
- J. L. Baker.** The Action of Ungerminated Barley Diastase on Starch. Part I. (Journ. Chem. Soc. Trans., 1902, 81, 1177.)
- E. R. Baldwin and P. A. Levene.** Action of Proteolytic Enzymes on Toxins. (J. Med. Research, 1901, 6, 120.)
- F. Banning.** Zur Kenntniss der Oxalsäurebildung durch Bakterien. (Centr. Bakt. Par., 1902, [ii], 8, 395, 425, 453, 520, 556.)
- G. Basile.** Fermentazione mannitica dei vini in Sicilia. (Staz. Sper. Agrar., 1894, 26, 451.)

- A. Bau.** Ueber ein neues Enzym der Hefe. (Chem. Zeit., 1895, 19, 1873.)
- A. Bau.** Ueber die Vergährbarkeit der Galaktose. (Bied. Centr., 1897, 26, 213.)
- A. Bau.** Ueber krystallisierte Melibiose. (Chem. Centr., 1899, ii, 526.)
- R. W. Bauer.** Ueber eine aus Aepfelpektin entstehende Zuckerart. (Landw. Versuchs-Stat., 43, 191.)
- E. Baumann.** Zur Kenntniss der aromatischen Substanzen des Thierkörpers. (Zeit. physiol. Chem., 1877-78, 1, 60.)
- E. Baumann.** Weitere Beiträge zur Kenntniss der aromatischen Substanzen des Thierkörpers. (Zeit. physiol. Chem., 1880, 4, 304.)
- E. Baumann.** Zur Kenntniss der aromatischen Produkte des Thierkörpers. (Ber. d. deutsch. chem. Ges., 1880, 13, 279.)
- E. Baumann.** Ueber die Bildung von Hydroparacumarsäure aus Tyrosin. (Ber. d. deutsch. chem. Ges., 1879, 12, 1450.)
- E. Baumann.** Die aromatische Verbindungen im Harn und die Darmfäulniss. (Zeit. physiol. Chem., 1886, 10, 123.)
- E. Baumann und L. Brieger.** Ueber die Entstehung von Kresolen bei der Fäulniss. (Zeit. physiol. Chem., 1879, 3, 149.)
- R. Benjamin.** Beiträge zur Lehre von der Labgerinnung. (Virchow's Archiv, 1896, 145, 30.)
- A. Berg.** Sur le mode de formation de l'elatérine dans l'*Ecballium elaterium*. (Bull. Soc. Chim., 1897, [iii], 17, 85.)
- C. Bernard.** Recherches sur les usages du suc pancréatique dans la digestion. (Compt. rend., 1849, 28, 249.)
- C. Bernard.** Leçons sur les phénomènes de la vie. Paris, 1878-79.
- G. Bertrand.** Sur le latex de l'arbre à laque. (Compt. rend., 1894, 118, 1215.)
- G. Bertrand.** Recherches sur le latex de l'arbre à laque du Tonkin. (Bull. Soc. Chim., 1894, [iii], 11, 717.)
- G. Bertrand.** Sur la laccase et sur le pouvoir oxydant de cette diastase. (Compt. rend., 1895, 120, 266.)
- G. Bertrand.** Sur la recherche et la présence de la laccase dans les végétaux. (Compt. rend., 1895, 121, 166.)
- G. Bertrand.** Sur une nouvelle oxydase ou ferment soluble oxydant d'origine végétale. (Compt. rend., 1896, 122, 1215. Bull. Soc. Chim., 1896, [iii], 15, 793.)

- G. Bertrand.** Sur les rapports qui existent entre la constitution chimique des composés organiques, et leur oxidabilité sous l'influence de la laccase. (Compt. rend., 1896, **122**, 1132.)
- G. Bertrand.** Sur la présence simultanée de la laccase et de la tyrosinase dans le suc de quelques champignons. (Compt. rend., 1896, **123**, 463.)
- G. Bertrand.** Sur l'intervention du manganèse dans les oxydations provoquées par la laccase. (Compt. rend., 1897, **124**, 1032.)
- G. Bertrand.** Sur l'action oxydante des sels manganeux et sur la constitution chimiques des oxydases. (Compt. rend., 1897, **124**, 1355.)
- G. Bertrand.** Nouvelles recherches sur les ferments oxydants ou oxydases. (Ann. Agron., 1897, **23**, 285.)
- G. Bertrand.** Recherches sur la laccase, nouveau ferment soluble à propriétés oxydantes. (Ann. Chim. Phys., 1897, [vii], **12**, 115.)
- G. Bertrand.** Sur la préparation biochimique de Sorbose. (Compt. rend., 1896, **122**, 900; Bull. Soc. Chim., 1896, [iii], **15**, 627.)
- G. Bertrand.** Action de la fleur du vin sur la sorbite. (Compt. rend., 1898, **126**, 653.)
- G. Bertrand.** Action de la bactérie du Sorbose sur les alcools plurivalents. (Compt. rend., 1898, **126**, 762.)
- G. Bertrand.** Sur le produit d'oxydation de la glycerine par la bactérie du Sorbose. (Compt. rend., 1898, **126**, 842.)
- G. Bertrand.** Préparation biochimique de la dioxyacetone cristallisée. (Compt. rend., 1898, **126**, 984.)
- G. Bertrand.** Action de la bactérie du Sorbose sur le sucre de bois. (Compt. rend., 1898, **127**, 124.)
- G. Bertrand.** Action de la bactérie du Sorbose sur les sucres aldéhydiques. (Compt. rend., 1898, **127**, 728.)
- G. Bertrand et A. Mallèvre.** Sur la pectase et sur la fermentation pectique. (Compt. rend., 1894, **119**, 1012.)
- G. Bertrand et A. Mallèvre.** Nouvelles recherches sur la pectase et la fermentation pectique. (Compt. rend., 1895, **120**, 110.)
- G. Bertrand et A. Mallèvre.** Sur la diffusion de la pectase dans le règne végétal et sur la préparation de cette diastase. (Compt. rend., 1895, **121**, 726.)
- M. W. Beyerinck.** Ueber Glucoside und Enzyme in den Wurzeln einiger Spiräaarten. (Chem. Centr., 1899, ii, 259.)
- M. W. Beyerinck.** On Indigo-fermentation. (Proc. K. Akad. Wetensch. Amsterdam, 1900, **2**, 495.)

- M. W. Beyerinck.** Further Researches on the Formation of Indigotin from Woad (*Isatis tinctoria*). (Proc. K. Akad. Wetensch. Amsterdam, 1900, **3**, 101.)
- M. W. Beyerinck.** Sur la production de Quinone par le *Streptothrix chromogena* et la biologie de ce microbe. (Arch. Néerl., 1900, iii, **2**, 327.)
- M. W. Beyerinck.** Sur la formation de l'hydrogène sulfuré dans les canaux, et le genre nouveau aérobacter. (Arch. Néerland. Sci. Exact. Nat., 1900, [ii], **4**, 1.)
- U. Biffi.** Zur Kenntniss der Spaltungsprodukte des Caseïns bei der Pankreasverdauung. (Virchow's Archiv, 1898, **152**, 130.)
- P. Biginelli.** Composizione e Costituzione chimica del gas arsenicale delle tapezzerie. (Atti Real. Accad. Lincei, 1900, [v], **9**, ii, 210, 242.)
- B. Binz.** Die Reduktion der Arsensäure durch Organsäfte. (Chem. Centr., 1897, i, 394.)
- C. Biondi.** Beiträge zur Lehre der Fermentativen Prozesse in den Organen. (Virchow's Archiv, 1896, **144**, 373.)
- F. W. J. Boekhout.** Ueber Dextranbildner. (Centr. Bakt. Par., 1900, [ii], **6**, 161.)
- T. Bokorny.** Das Verhalten nitrirter Kohlehydrate gegen Pilze. (Chem. Zeit., 1896, **20**, 985.)
- T. Bokorny.** Ernährbarkeit der Spaltpilze durch verschiedene Kohlenstoffverbindungen. (Pflüger's Archiv, 1897, **68**, 114.)
- T. Bokorny.** Enthalten die keimenden Samen peptonisierende oder andere proteolytische Enzyme? (Pflüger's Archiv, 1902, **90**, 94.)
- G. W. Boland.** Ueber Pyocyamin, den blauen Farbstoff des *Bacillus pyocyaneus*. (Chem. Centr., 1899, ii, 265.)
- C. Bouchard.** Sur les poisons qui existent normalement dans l'organisme et en particulier sur la toxicité urinaire. (Compt. rend., 1886, **102**, 669.)
- C. Bouchard.** Sur les variations de la toxicité urinaire pendant la veille et pendant le sommeil. (Compt. rend., 1886, **102**, 727.)
- C. Bouchard.** Influence de l'abstinence, du travail musculaire et de l'air comprimé sur les variations de la toxicité urinaire. (Compt. rend., 1886, **102**, 1127.)
- A. Bouffard.** Sur la cassage des vins. (Compt. rend., 1894, **118**, 827.)
- A. Bouffard.** Détermination de la chaleur dégagée dans la fermentation alcoolique. (Compt. rend., 1895, **121**, 357.)

- A. Bouffard.** Observations sur quelques propriétés de l'oxydase des raisins. (Compt. rend., 1897, **124**, 706.)
- A. Bouffard et L. Semichon.** Contribution à l'étude de l'oxydase des raisins. (Compt. rend., 1898, **126**, 423.)
- E. E. Bourquelot.** Ferments solubles sécrétés par l'*Aspergillus niger* et le *Penicillium glaucum*. (Bull. Soc. Biol., 1893, 653.)
- E. E. Bourquelot.** Sur un ferment soluble nouveau dédoublant le tréhalose en glucose. (Compt. rend., 1893, **116**, 826.)
- E. E. Bourquelot.** Inulase et fermentation alcoolique indirecte de l'inuline. (Compt. rend., 1893, **116**, 1143.)
- E. E. Bourquelot.** Présence d'un ferment analogue à l'émulsine dans les champignons. (Compt. rend., 1894, **117**, 383.)
- E. E. Bourquelot.** Sur la présence de l'éther méthylsalicique dans quelques plantes indigènes. (J. pharm. et chim., 1894, [v], **30**, 433.)
- E. E. Bourquelot.** Maltase et fermentation alcoolique du maltose. (J. pharm. et chim., 1895, [vi], **2**, 97.)
- E. E. Bourquelot.** Sur l'hydrolyse de raffinose (mélitose) par les ferments solubles. (J. pharm. et chim., 1896, [vi], **3**, 390.)
- E. E. Bourquelot.** Sur la présence dans le *Monotropa hypopythis* d'un glucoside et sur le ferment soluble hydrolysant de ce glucoside. (Compt. rend., 1896, **122**, 1002, and J. pharm. et chim., 1896, **3**, 577.)
- E. E. Bourquelot.** Des composés oxydables sous l'influence du ferment oxydant des champignons. (Compt. rend., 1896, **123**, 315.)
- E. E. Bourquelot.** Action du ferment soluble oxydant des champignons sur les phénols insolubles dans l'eau. (Compt. rend., 1896, **123**, 423.)
- E. E. Bourquelot.** Nouvelles recherches sur le ferment oxydant des champignons. II. Son action sur les phénols. III. Son action sur quelques dérivés éthers des phénols. (J. pharm. et chim., 1896, [vi], **4**, 241 and 440.)
- E. E. Bourquelot.** Sur la physiologie du gentianose : son dédoublement par les ferments solubles. (Compt. rend., 1898, **126**, 1045.)
- E. E. Bourquelot.** Sur les pectines. (Compt. rend., 1899, **128**, 1241, and J. pharm. et chim., 1899, [vi], **9**, 563.)
- E. E. Bourquelot et G. Bertrand.** La laccase dans les champignons. (Compt. rend., 1895, **121**, 783.)
- E. E. Bourquelot et G. Bertrand.** Sur la coloration des tissus

- et du suc de certains champignons au contact de l'air. (J. pharm. et chim., 1896, [vi], 3, 177.)
- E. E. Bourquelot et E. Gley.** Action du sérum sanguin sur la matière glycogène et sur le maltose. (Compt. rend. Soc. Biol., 1895, 247.)
- E. E. Bourquelot et E. Gley.** Digestion du tréhalose. (Compt. rend. Soc. Biol., 1895, 555.)
- E. E. Bourquelot et H. Hérissey.** Sur les propriétés de l'émulsine des champignons. (Compt. rend., 1895, 121, 693.)
- E. E. Bourquelot et H. Hérissey.** Sur l'hydrolyse du mélézitose par les ferments solubles. (J. pharm. et chim., 1896, [vi], 4, 385.)
- E. E. Bourquelot et H. Hérissey.** Sur l'existence, dans l'orge germée, d'un ferment soluble, agissant sur la pectine. (Compt. rend., 1898, 127, 191.)
- E. E. Bourquelot et H. Hérissey.** Sur la matière gélatineuse (pectine) de la racine de gentiane. (J. pharm. et chim., 1898, [vi], 7, 473.)
- E. E. Bourquelot et H. Hérissey.** Sur l'hydrolyse de la pectine de gentiane. (J. pharm. et chim., 1898, [vi], 8, 49.)
- E. E. Bourquelot et H. Hérissey.** De l'action des ferments solubles sur les produits pectiques de la racine de gentiane. (J. pharm. et chim., 1898, [vi], 8, 145.)
- E. E. Bourquelot et H. Hérissey.** Sur la pectine de *Cynorrhodon*. (J. pharm. et chim., 1899, [vi], 10, 5.)
- E. E. Bourquelot et H. Hérissey.** Sur la constitution du gentianose. (Compt. rend., 1901, 132, 571.)
- E. E. Bourquelot et H. Hérissey.** Recherche et présence d'un ferment soluble protéolytique dans les champignons. (Compt. rend., 1898, 127, 666.)
- E. E. Bourquelot et H. Hérissey.** Sur la composition de l'albumen de la graine de caroubier; production de galactose et de mannose par hydrolyse. (Compt. rend., 1899, 129, 228, 391.)
- E. E. Bourquelot et H. Hérissey.** Germination de la graine de caroubier; production de mannose par un ferment soluble. (Compt. rend., 1899, 129, 614.)
- E. E. Bourquelot et H. Hérissey.** Sur les ferments solubles produits pendant la germination par les graines à albumen corné. (Compt. rend., 1900, 130, 42.)

- E. E. Bourquelot et H. Hérissé.** Sur l'individualité de la séminase, ferment soluble sécrété par les graines de légumineuses à albumen corné pendant la germination. (Compt. rend., 1900, 130, 340.)
- E. E. Bourquelot et H. Hérissé.** Les hydrates de carbone de réserve des graines de Luzerne et de Fenugrec. (Compt. rend., 1900, 130, 731.)
- E. E. Bourquelot et H. Hérissé.** Sur la présence de séminase dans les graines à albumen corné au repos. (Compt. rend., 1900, 131, 903.)
- E. E. Bourquelot et H. Hérissé.** Sur la composition de l'albumen de la graine de *Phoenix canariensis* et sur les phénomènes chimiques qui accompagnent la germination de cette graine. (Compt. rend., 1901, 133, 302.)
- E. E. Bourquelot et L. Nardin.** Sur la préparation du gentianose. (Compt. rend., 1898, 126, 280.)
- L. Bréaudat.** Sur le mode de formation de l'indigo dans les procédés d'extraction industriels. Fonctions diastatiques des plantes indigofères. (Compt. rend., 1898, 127, 769.)
- L. Bréaudat.** Nouvelles recherches sur les fonctions diastatiques des plantes indigofères. (Compt. rend., 1899, 128, 1478.)
- L. Brieger.** Die Ptomaine. Berlin, 1885 and 1886.
- A. J. Brown.** The Chemical Action of Pure Cultivations of *Bacterium aceti*. (Journ. Chem. Soc. Trans., 1886, 172.)
- A. J. Brown.** An Acetic Ferment which forms Cellulose. (Journ. Chem. Soc. Trans., 1886, 432.)
- A. J. Brown.** On the Influence of Oxygen and Concentration on Fermentation. (Journ. Chem. Soc. Trans., 1892, 369.)
- A. J. Brown.** The Specific Character of the Fermentative Functions of Yeast Cells. (Journ. Chem. Soc. Trans., 1894, 911.)
- A. J. Brown.** Enzyme Action. (Journ. Chem. Soc. Trans., 1902, 373.)
- H. T. Brown.** A Search for a Cellulose-dissolving (cytohydrolytic) Enzyme in the Digestive Tract of Certain Grain-feeding Animals. (Journ. Chem. Soc. Trans., 1892, 352.)
- H. T. Brown and J. Heron.** Contributions to the History of Starch and its Transformations. (Journ. Chem. Soc. Trans., 1879, 596.)
- H. T. Brown and J. Heron.** Some Observations on the Hydrolytic Ferments of the Pancreas and Small Intestine. (Proc. Roy. Soc., 1880, 393.)

- H. T. Brown and G. H. Morris.** The Non-crystallisable Products of the Action of Diastase on Starch. (*Journ. Chem. Soc. Trans.*, 1885, 527.)
- H. T. Brown and G. H. Morris.** The Amylodextrin of W. Nägeli and its Relation to Soluble Starch. (*Journ. Chem. Soc. Trans.*, 1889, 449.)
- H. T. Brown and G. H. Morris.** The Determination of the Molecular Weights of the Carbohydrates. Part II. (*Journ. Chem. Soc. Trans.*, 1889, 55, 462.)
- H. T. Brown and G. H. Morris.** On the Action of Diastase in the Cold on Starch-paste. (*Journ. Chem. Soc. Trans.*, 1895, 309.)
- H. T. Brown and G. H. Morris.** On the Isomaltose of C. J. Lintner. (*Journ. Chem. Soc. Trans.*, 1895, 709.)
- H. T. Brown, G. H. Morris, and J. H. Millar.** Experimental Methods employed in the Determination of the Products of Starch-hydrolysis by Diastase. (*Journ. Chem. Soc. Trans.*, 1897, 72.)
- H. T. Brown, G. H. Morris, and J. H. Millar.** Relations of Specific Rotatory Powers and Cupric-reducing Powers of Starch-hydrolysis by Diastase. (*Journ. Chem. Soc. Trans.*, 1897, 115.)
- H. T. Brown and J. H. Millar.** Maltodextrin: its Oxidation Products and Constitution. (*Journ. Chem. Soc. Trans.*, 1899, 75, 286.)
- H. T. Brown and J. H. Millar.** The Stable Dextrin of Starch Transformations, and its Relation to the Maltodextrins and Soluble Starch. (*Journ. Chem. Soc. Trans.*, 1899, 75, 315.)
- H. T. Brown and T. A. Glendinning.** The Velocity of Starch Hydrolysis by Diastase, with Some Remarks on Enzyme Action. (*Journ. Chem. Soc. Trans.*, 1902, 81, 388.)
- H. T. Brown and S. Pickering.** Thermal phenomena attending the change in rotatory power of freshly prepared solutions of certain carbohydrates. (*Journ. Chem. Soc. Trans.*, 1897, 71, 758.)
- H. T. Brown and S. Pickering.** Thermochemistry of Carbohydrate Hydrolysis. (*Journ. Chem. Soc. Trans.*, 1897, 71, 783.)
- E. Brücke.** Studien über die Kohlehydrate und über die Art, wie sie verdaut und aufgesaugt werden. (*Wien. Acad. Ber.*, 1872, ii, 65, 126.)
- E. Buchner.** Alcoholische Gährung ohne Hefezellen. (*Ber. d. deutsch. chem. Ges.*, 1897, 30, 117 and 1110.)

- E. Buchner.** Ueber zellenfreie Gährung. (Ber. d. deutsch. chem. Ges., 1898, **31**, 568.)
- E. Buchner und R. Rapp.** Alcoholische Gährung ohne Hefezellen. (Ber. d. deutsch. chem. Ges., 1897, **30**, 2668; 1898, **31**, 209, 1084, 1090, 1581; 1899, **32**, 127.)
- K. Bülow.** Ueber die dextrinartigen Abbauproducte der Stärke. (Pflüger's Archiv, 1896, **62**, 131.)
- R. Burri et A. Stutzer.** Sur les bactéries destructives de nitrates et sur la perte d'azote qui résulte de leur action. (Ann. Agron., 1896, **22**, 491.)
- W. Butkewitsch.** Ueber das Vorkommen proteolytischer Enzyme in gekeimten Samen und über ihre Wirkung. (Chem. Centr., 1901, i, 190.)
- L. Camus.** Formation de lipase par le *Penicillium glaucum*. (Compt. rend. Soc. Biol., 1897, 192.)
- L. Camus.** De la lipase dans les cultures l'*Aspergillus niger*. (Compt. rend. Soc. Biol., 1897, 230.)
- L. Camus et E. Gley.** Action coagulante du liquide prostatique sur le contenu des vésicules séminales. (Compt. rend., 1896, **123**, 194.)
- L. Camus et E. Gley.** Action coagulante du liquide de la prostate externe du Hérisson sur le contenu des vésicules séminales. (Compt. rend., 1899, **128**, 1417.)
- P. Carles.** Valériane et oxydase. (J. pharm. et chim., 1900, [vi], **12**, 148.)
- E. Cavazzani.** Versuche über die Anwesenheit eines Oxydationsfermentes in der Cerebrospinalflüssigkeit. (Chem. Centr., 1901, i, 333.)
- P. Cazeneuve.** Sur un ferment soluble oxydant de la casse des vins. (Compt. rend., 1897, **124**, 406.)
- P. Cazeneuve.** Sur quelques propriétés du ferment de la casse des vins. (Compt. rend., 1897, **124**, 781.)
- R. Chevastolon.** Sur l'inuline de l'ail, de la jacinthe, de l'asphodèle et de la tubérose. (J. pharm. et chim., 1895, [vi], **2**, 83.)
- R. H. Chittenden.** On the Proteolytic Action of Bromelin, the Ferment of Pine-apple Juice. (J. Physiol., 1894, **15**, 249.)
- R. H. Chittenden and Amerman.** A Comparison of Natural and Artificial Gastric Digestion. (J. Physiol., 1893, **14**, 483.)

- R. H. Chittenden and Hartwell.** The Relative Formation of Proteoses and Peptones in Gastric Digestion. (*J. Physiol.*, 1891, **12**, 12.)
- R. H. Chittenden and L. B. Mendel.** Proteolysis of Crystallised Globulin. (*J. Physiol.*, 1894, **17**, 48.)
- R. H. Chittenden, L. B. Mendel, and McDermott.** Papain-Proteolysis. (*Amer. J. Physiol.*, 1898, **1**, 255.)
- R. H. Chittenden and Smith.** On the Primary Cleavage Products formed in the Digestion of Gluten-casein of Wheat by Pepsin-hydrochloric Acid. (*J. Physiol.*, 1890, **9**, 410.)
- R. H. Chittenden and Solley.** The Primary Cleavage Products formed in the Digestion of Gelatine. (*J. Physiol.*, 1891, **12**, 23.)
- A. A. Claflin.** Manufacture and Applications of Lactic Acid. (*J. Soc. Chem. Ind.*, 1897, **16**, 516.)
- O. Cohnheim.** Die Umwandlung des Eiweiss durch die Darmwand. (*Zeit. physiol. Chem.*, 1901, **33**, 451.)
- O. Cohnheim.** Weitere Mittheilungen über das Erepsin. (*Zeit. physiol. Chem.*, 1902, **35**, 134.)
- Cornu.** Sur les ferments oxydants de la vigne. (*J. pharm. et chim.*, 1899, [vi], **10**, 342.)
- G. Cossetini.** Ueber das Philothion. (*Chem. Centr.*, 1901, i, 789.)
- C. D. Cramer.** Ueber das sogenannte Thrombosin Lilienfelds. (*Zeit. physiol. Chem.*, 1897, **23**, 74.)
- Cross and Bevan.** Cellulose. London, 1895.
- F. Czapek.** Zur Biologie der holzbewohnenden Pilze. (*Ber. d. deutsch. bot. Ges.*, 1899, **17**, 166.)
- E. Demoussy.** Sur l'oxydation des ammoniacques composées par les ferments du sol. (*Compt. rend.*, 1898, **126**, 253.)
- R. Dubois.** Sur la luciférase ou zymase photogène des animaux et des vegetaux. (*Compt. rend.*, 1896, **123**, 653.)
- G. Düll.** Ueber die Einwirkung von Oxalsäure auf Inulin. (*Chem. Zeit.*, 1895, **19**, 166 and 216.)
- W. R. Dunstan and T. A. Henry.** Nature and Origin of the Poison of *Lotus Arabicus*. (*Proc. Roy. Soc.*, 1900, **67**, 224, and 1901, **68**, 374.)
- S. Dzierzgowski and S. Salaskin.** Ueber die Ammoniakabspaltung bei der Einwirkung von Trypsin und Pepsin auf Eiweisskörper. (*Chem. Centr.*, 1901, ii, 645.)

- J. Effront.** Sur la formation de l'acide succinique et de la glycerine dans la fermentation alcoolique. (Compt. rend., 1894, **119**, 169.)
- J. Effront.** Sur un nouvel hydrate de carbone, la caroubine. (Compt. rend., 1897, **125**, 38.)
- J. Effront.** Sur une nouvelle enzyme hydrolytique "la caroubinase." (Compt. rend., 1897, **125**, 116.)
- J. Effront.** Sur la caroubinose. (Compt. rend., 1897, **125**, 309.)
- J. Effront.** Les enzymes et leurs applications. Paris, 1899.
- A. Ehrenberg.** Ueber einige in einem Falle von sogenannter "Wurstvergiftung" aus dem schädlichen Materiale dargestellte Fäulnisbasen, sowie über einige, durch die Thätigkeit eines besonderen, im gleichen Materiale aufgefundenen, Bacillus gebildete Zersetzungs-Produkte. (Zeit. physiol. Chem., 1887, **11**, 239.)
- W. A. van Ekenstein.** Sur la caroubinose et sur la d-mannose. (Compt. rend., 1897, **125**, 719.)
- R. L. Emerson.** Ueber das Auftreten von Oxyphenyläthylamin bei Pankreasverdauung und über fermentative CO₂-Abspaltung. (Beitr. chem. Physiol. Path., 1902, **1**, 501.)
- R. Emmerich and O. Loew.** Bakteriologische Enzyme als Ursache der erworbenen Immunität und die Heilung von Infektionskrankheiten durch dieselben. (Chem. Centr., 1899, ii, 135.)
- O. Emmerling.** Ueber einen neuen aus Glycerin Buttersäure erzeugenden Bacillus. (Ber. d. deutsch. chem. Ges., 1896, **29**, 2726.)
- O. Emmerling.** Zur Frage, wodurch die Giftigkeit arsenhaltiger Tapeten bewirkt sind. (Ber. d. deutsch. chem. Ges., 1896, **29**, 2728.)
- O. Emmerling.** Butylalkoholische Gährung. (Ber. d. deutsch. chem. Ges., 1897, **30**, 451.)
- O. Emmerling.** Ueber Schimmelpilzgährung. (Ber. d. deutsch. chem. Ges., 1897, **30**, 454, and 1899, **32**, 1915.)
- O. Emmerling.** Bemerkung zu vorstehender Entgegnung des Herrn Gosio. (Ber. d. deutsch. chem. Ges., 1897, **30**, 1026.)
- O. Emmerling.** Zur Kenntniss des Sorbosebacteriums. (Ber. d. deutsch. chem. Ges., 1899, **32**, 541.)
- O. Emmerling.** Das Verhalten von Glycerinaldehyd und Dioxyacetone gegen Hefe. (Ber. d. deutsch. chem. Ges., 1899, **32**, 542.)
- O. Emmerling.** Ueber Spaltpilzgährungen. (Ber. d. deutsch. chem. Ges., 1900, **33**, 2477.)

- O. Emmerling.** Synthetische Wirkung der Hefenmaltase. (Ber. d. deutsch. chem. Ges., 1901, **34**, 600, 2206, 3810.)
- O. Emmerling.** Ueber die Eiweisspaltung durch Papayotin. (Ber. d. deutsch. chem. Ges., 1902, **35**, 695 and 1012.)
- O. Emmerling.** Aminosäuren als Nährstoffe für niedere Pflanzen. (Ber. d. deutsch. chem. Ges., 1902, **35**, 2289.)
- O. Emmerling.** Die Zersetzung Stickstofffreier organischer Substanzen durch Bakterien. Braunschweig, 1902.
- O. Emmerling und O. Reiser.** Zur Kenntniss eiweisspaltender Bakterien. (Ber. d. deutsch. chem. Ges., 1902, **35**, 700.)
- A. Fernbach.** Sur la tannase. (Compt. rend., 1901, **131**, 1214.)
- A. Fernbach et Hubert.** Sur la diastase protéolytique du malt. (Compt. rend., 1900, **130**, 1783.)
- E. Fischer.** Synthese einer neuen Glucobiose. (Ber. d. deutsch. chem. Ges., 1890, **23**, 3689.)
- E. Fischer.** Einfluss der Configuration auf die Wirkung der Enzyme. (Ber. d. deutsch. chem. Ges., 1894, **27**, 2985, 3479, and 1895, **28**, 1429.)
- E. Fischer.** Spaltung von Trehalose. (Ber. d. deutsch. chem. Ges., 1895, **28**, 1433.)
- E. Fischer.** Ueber ein neues dem Amygdalin ähnliches Glucosid. (Ber. d. deutsch. chem. Ges., 1895, **28**, 1508.)
- E. Fischer.** Ueber die Isomaltose. (Ber. d. deutsch. chem. Ges., 1895, **28**, 3024.)
- E. Fischer.** Bedeutung der Stereochemie für die Physiologie. (Zeit. physiol. Chem., 1898–99, **26**, 60.)
- E. Fischer und E. F. Armstrong.** Synthese einiger neuen Disaccharide. (Ber. Akad. Wiss. Berlin, 1901, **7**, 123, and Ber. d. deutsch. chem. Ges., 1902, **35**, 3144.)
- E. Fischer und E. F. Armstrong.** Darstellung der Osone aus der Osazonen der Zucker. (Ber. d. deutsch. chem. Ges., 1902, **35**, 3141.)
- E. Fischer und E. F. Armstrong.** Ueber die isomeren Acetohalogen-Derivate der Zucker und die Synthese der Glucoside. (Ber. d. deutsch. chem. Ges., 1902, **35**, 3153.)
- E. Fischer und P. Lindner.** Ueber die Enzyme von *Schizosaccharomyces octosporus* und *S. Marxianus*. (Ber. d. deutsch. chem. Ges., 1895, **28**, 984.)

- E. Fischer und P. Lindner.** Ueber die Enzyme einiger Hefen. (Ber. d. deutsch. chem. Ges., 1895, **28**, 3034.)
- E. Fischer und P. Lindner.** Verhalten der Enzyme gegen Melibiose, Rohrzucker und Maltose. (Ber. d. deutsch. chem. Ges., 1895, **28**, 3035.)
- E. Fischer und W. Niebel.** Ueber das Verhalten der Polysaccharide gegen einige thierische Secrete und Organe. (Chem. Centr., 1896, i, 499; Ber. d. deutsch. chem. Ges., 1896, **29**, iv, 1156.)
- A. Fitz.** Ueber die Gährung des Glycerins. (Ber. d. deutsch. chem. Ges., 1876, **9**, 1348.)
- A. Fitz.** Ueber alkoholische Gährung. (Ber. d. deutsch. chem. Ges., 1876, **9**, 1352.)
- A. Fitz.** Ueber Schizomycetgährungen. (Ber. d. deutsch. chem. Ges., 1877, **10**, 276.)
- A. Fitz.** Ueber Spaltpilzgährungen. (Ber. d. deutsch. chem. Ges., 1878, **11**, 1898; 1879, **12**, 479; 1880, **13**, 1309.)
- C. Flügge.** Die Microorganismen. Leipzig, 1896.
- S. Fränkel und L. Langstein.** Ueber die Spaltungsprodukte des Eiweisses bei der Verdauung. (Monatsh., 1901, **22**, 335.)
- P. F. and G. C. Frankland.** The Nitrifying Process and its Specific Ferment. (Phil. Trans. Roy. Soc. B., 1890, 107.)
- P. F. Frankland.** Pasteur Memorial Lecture. (J. Chem. Soc. Trans., 1897, 683.)
- P. F. Frankland and J. J. Fox.** Fermentation of Mannitol and Glycerol. (Proc. Roy. Soc., **46**, 345.)
- P. F. Frankland and W. Frew.** The Fermentation of Calcium Glycerate by *Bacillus ethaceticus*. (J. Chem. Soc. Trans., 1891, **59**, 81.)
- P. F. Frankland and W. Frew.** An Optically Active Glyceric Acid. (J. Chem. Soc. Trans., 1891, **59**, 96.)
- P. F. Frankland and W. Frew.** A Pure Fermentation of Mannitol and Dulcitol. (J. Chem. Soc. Trans., 1892, **61**, 254.)
- P. F. Frankland, A. Stanley, and W. Frew.** Fermentations induced by the Pneumococcus of Friedländer. (J. Chem. Soc. Trans., 1891, **59**, 253.)
- P. F. Frankland and J. S. Lumsden.** The Decomposition of Mannitol and Dextrose by the *Bacillus ethaceticus*. (J. Chem. Soc. Trans., 1892, **61**, 432.)

- P. F. Frankland and J. MacGregor.** The Fermentation of Arabinose by *Bacillus ethaceticus*. (J. Chem. Soc. Trans., 1892, **61**, 737.)
- P. F. Frankland and J. MacGregor.** Sarcolactic Acid obtained by Fermentation of Inactive Lactic Acid. (J. Chem. Soc. Trans., 1893, **63**, 1028.)
- J. J. Frederikse.** Einiges über Fibrin und Fibrinogen. (Zeit. physiol. Chem., 1894, **19**, 143.)
- Fremy.** Recherches sur les fermentations. (Compt. rend., 1872, **75**, 976 and 1060.)
- H. Friedenthal.** Das Molekulargewicht der löslichen Stärke. (Chem. Centr., 1899, **i**, 924.)
- E. von Freudenreich.** Ueber das in der Milch vorhandene unorganisierte Ferment, die sogenannte Galaktase. (Centr. Bakt. Par., 1900, [ii], **6**, 332.)
- O. von Fürth.** Ueber die Eiweisskörper des Muskelplasmas. (Arch. exp. Path. Pharm., 1895, **36**, 231.)
- O. von Fürth und H. Schneider.** Ueber tierische Tyrosinasen und Ihre Beziehung zur Pigmentbildung. (Beitr. chem. Physiol. Path., 1901, **1**, 229.)
- J. Gadamer.** Les glucosides des moutardes noire et blanche. (J. Pharm. 1896, [vi], **4**, 462.)
- B. Galli-Valerio und C. Strzyzowski.** Ueber den biologischen Arsennachweis. (Chem. Centr., 1901, **i**, 63.)
- S. A. Garcia.** Ueber Ptomaine, welche bei der Fäulniss von Pferdefleisch und Pankreas entstehen. (Zeit. physiol. Chem., 1893, **17**, 543.)
- U. Gayon et E. Dubourg.** Nouvelles recherches sur le ferment mannitique. (Ann. Inst. Pasteur, 1901, **15**, 527.)
- E. Gérard.** Présence dans le *Penicillium glaucum* d'un ferment agissant comme l'émulsine. (Compt. rend. Soc. Biol., 1893, 651, and J. Pharm., 28, [v], 11.)
- E. Gérard.** Sur le dédoublement de l'amydaline dans l'économie. (J. Pharm. et Chim., 1896, [vi], **3**, 233.)
- E. Gérard.** Sur une lipase végétale extraite du *Penicillium glaucum*. (Compt. rend., 1897, **124**, 370.)
- E. Gérard.** Fermentation de l'acide urique par les microorganismes. (Compt. rend., 1896, **122**, 1019, and **123**, 185.)

- E. Gérard.** Transformation de la créatine en créatinine par un ferment soluble déshydratant de l'organisme. (Compt. rend., 1901, **132**, 153.)
- L. Geret und M. Hahn.** Zum Nachweis des im Hefepresssaft enthaltenen proteolytischen Enzyms. (Ber. d. deutsch. chem. Ges., 1898, **31**, 206 and 2335.)
- Gerlach und Vogel.** Ueber eiweissbildende Bakterien. (Centr. Bakt. Par., 1901, [ii], **7**, 609.)
- Gessard.** Sur la tyrosinase. (Compt. rend., 1900, **130**, 1327.)
- H. Gillot.** Recherches experimentales sur l'hydrolyse et l'utilisation de la raffinose par le *Penicillium glaucum*. (Bull. Acad. Roy. Belg., 1900, 99.)
- M. Gonnermann.** Ueber die dextrinartigen Abbauprodukte der Stärke. (Pflüger's Archiv, 1902, **89**, 493.)
- B. Gosio.** Zur Frage, wodurch die Giftigkeit arsenhaltiger Tapeten bedingt sind. (Ber. d. deutsch. chem. Ges., 1897, **30**, 1024.)
- B. Gosio.** Ulteriori ricerche sulla biologia e sul chimismo delle arseniomuffe. (L'Orosi, 1900, **23**, 361.)
- Gouirand.** Sur la présence d'une diastase dans les vins cassés. (Compt. rend., 1895, **120**, 887.)
- J. Reynolds Green.** The Soluble Ferments and Fermentation. Cambridge, 1901.
- V. Griessmayer.** Ueber das Verhalten von Stärke und Dextrin gegen Jod und Gerbsäure. (Liebig's Annalen, 1871, **160**, 40.)
- L. Grimbert.** Action du pneumocoque de Friedländer sur les sucres. (J. Pharm., 1895, [vi], **2**, 529.)
- L. Grimbert.** Production d'acétylméthylcarbinol par le *Bacillus tartricus*. (Compt. rend., 1901, **132**, 706.)
- L. Grimbert et L. Ficquet.** Sur un nouveau ferment des tartrates, le *Bacillus tartricus*. (J. Pharm., 1898, [vi], **7**, 97.)
- J. Grüss.** Ueber Lösung von Cellulose durch Enzyme (Cytase). (Chem. Centr., 1896, i. 313.)
- J. Grüss.** Ueber Oxydasen und die Guajakreaction. (Chem. Centr., 1898, ii, 366.)
- J. Grüss.** Ueber Oxydaseerscheinungen der Hefe. (Chem. Centr., 1901, ii, 364 and 436.)
- W. Gulewitch.** Ueber das Verhalten des Trypsins gegen einfachere chemische Verbindungen. (Zeit. physiol. Chem., 1899, **27**, 540.)

- M. Hahn.** Das proteolytische Enzym des Hefepresssaftes. (Ber. d. deutsch. chem. Ges., 1898, **31**, 200.)
- M. Hahn und L. Geret.** Ueber das Hefe-Endotrypsin. (Zeit. Biol., 1900, **40**, 117.)
- W. D. Halliburton.** On Muscle Plasma. (J. Physiol., 1887, **8**, 159.)
- W. D. Halliburton.** On the Nature of Fibrin Ferment. (J. Physiol., 1888, **9**, 265.)
- W. D. Halliburton.** Nucleo-proteids. (J. Physiol., 1895, **18**, 306.)
- O. Hammarsten.** Ueber die Milchgerinnung und die dabei wirkenden Fermente der Magenschleimhaut. (Maly's Ber., 1872, 118.)
- O. Hammarsten.** Ueber den chemischen Verlauf bei der Gerinnung des Caseïns mit Lab. (Maly's Ber., 1874, **4**, 135.)
- O. Hammarsten.** Zur Kenntniss des Caseïns, und der Wirkung des Labfermentes. (Maly's Ber., 1877, **7**, 166.)
- O. Hammarsten.** Zur Lehre von der Faserstoffgerinnung. (Pflüger's Arch., 1877, **14**, 211.)
- O. Hammarsten.** Ueber das Paraglobulin. (Pflüger's Arch., 1878, **17**, 413; **18**, 38.)
- O. Hammarsten.** Ueber das Fibrinogen. (Pflüger's Arch., 1878, **19**, 563.)
- O. Hammarsten.** Ueber den Faserstoff und seine Entstehung aus dem Fibrinogen. (Pflüger's Arch., 1883, **30**, 457.)
- O. Hammarsten.** Ueber die Bedeutung der löslichen Kalksalze für die Faserstoffgerinnung. (Zeit. physiol. Chem., 1896, **22**, 333.)
- O. Hammarsten.** Weitere Beiträge zur Kenntniss der Fibrinbildung. (Zeit. physiol. Chem., 1899, **28**, 98.)
- M. Hanriot.** Sur un nouveau ferment du sang. (Compt. rend., 1896, **123**, 753.)
- M. Hanriot.** Sur la non-identité des lipases d'origine différente. (Compt. rend., 1897, **124**, 778.)
- M. Hanriot.** Sur le mécanisme des actions diastatiques. (Compt. rend., 1901, **132**, 146, 212.)
- M. Hanriot.** Sur le mécanisme des reactions lipolytiques. (Compt. rend., 1901, **132**, 842.)
- M. Hanriot et L. Camus.** Sur le dosage de la lipase. (Compt. rend., 1897, **124**, 235.)
- A. Vernon Harcourt.** On the Observation of the Course of Chemical Change. (J. Chem. Soc. Trans., **20**, 460.)
- V. Harlay.** Caractères différentiels des produits de la digestion

- pepsique et de la digestion pancreatique de la fibrine. (J. Pharm., 1899, [vi], **9**, 225.)
- V. **Harlay**. Sur quelques observations relatives aux digestions pepsique et pancreatique de l'albumine. (J. Pharm., 1899, [vi], **9**, 424.)
- V. **Harlay**. Sur la digestion de la fibrine et d'albumine par la papaine, et sur une reaction colorée nouvelle des produits obtenus. (J. Pharm., 1900, [vi], **11**, 172.)
- D. F. **Harris**. Coagulation of Milk. (J. Anat. and Physiol., 1894, **29**, 188.)
- G. **Hayem**. Observation à l'occasion du travail de M. Arthus sur le dosage comparatif du fibrinogène et de la fibrine. (Compt. rend. Soc. Biol., 1894, 309.)
- J. J. **Hazewinkel**. Indican—its Hydrolysis and the Enzyme causing the Same. (Proc. K. Akad. Wetensch. Amsterdam, 1900, **2**, 512.)
- A. **Harden**. The Chemical Action of *Bacillus coli communis* and Similar Organisms on Carbohydrates and Allied Compounds. (J. Chem. Soc. Trans., 1901, **79**, 610.)
- S. G. **Hedin**. Zur Kenntniss der Produkte der tryptischen Verdauung des Fibrins. (Du Bois Archiv. f. Physiol., 1891, 273.)
- V. **Henri**. Influence du sucre interverti sur la vitesse d'inversion du saccharose par la sucrase. (Compt. rend. Soc. Biol., 1901, **53**, 288.)
- V. **Henri**. Influence de l'addition, au milieu d'une réaction, de saccharose ou de sucre inverti sur la vitesse d'inversion par la sucrase. (Compt. rend. Soc. Biol., 1901, **53**, 290.)
- H. **Hérissey**. Digestion of the Mannan of Orchid Tubers. (Compt. rend., 1902, **134**, 721.)
- A. **Herzfeld**. Ueber die Einwirkung der Diastase auf Stärkekleister. (Ber. d. deutsch. chem. Ges., 1879, **12**, 2120.)
- Hilger**. Neues Vorkommen des Inosits im Pflanzenreiche und Ueberführung desselben in Paramilchsäure. (Ann. Chem. Pharm., 1871, **160**, 333.)
- A. **Croft Hill**. Reversible Zymohydrolysis. (J. Chem. Soc. Trans., 1898, 634.)
- A. **Croft Hill**. Bemerkungen zur Arbeit von O. Emmerling: Synthetische Wirkung der Hefenmaltase. (Ber. d. deutsch. chem. Ges., 1901, **34**, 1380.)
- A. **Croft Hill**. Taka-diastase and Reversed Ferment Action. (Proc. Chem. Soc., 1901, **17**, 184.)

- A. Croft Hill.** Synthetic Action on Dextrose with Pancreatic Ferment. (Proc. Physiol. Soc. J. Physiol., 1902, **28**, xxvi.)
- S. Hoogewerff and H. ter Meulen.** Contribution to the Knowledge of Indican. (Proc. K. Akad. Wetensch. Amsterdam, 1900, **2**, 520.)
- F. G. Hopkins and S. W. Cole.** Proteid Chemistry, I. A hitherto Undiscovered Product of Tryptic Digestion. (J. Physiol., 1901, **27**, 418.)
- F. Hoppe-Seyler.** Ueber die Prozesse der Gährungen und ihre Beziehung zum Leben der Organismen. (Pflüger's Archiv, 1876, **12**, 1.)
- F. Hoppe-Seyler.** Gährung der Cellulose. (Ber. d. deutsch. chem. Ges., 1883, **16**, 122.)
- R. M. Horne.** Coagulation of Blood. (J. Physiol., 1896, **19**, 356.)
- D. P. Hoyer.** Untersuchungen über die Essigsäurebakterien. (Chem. Centr., 1899, i, 854.)
- M. Jacoby.** Ueber das Aldehyde oxydirende Ferment der Leber und Nebenniere. (Zeit. physiol. Chem., 1900, **30**, 135.)
- E. Jalowetz.** Isomaltose. (Chem. Zeit., 1895, **19**, 2003.)
- H. Jensen.** Studium über das Ranzigwerden der Butter. (Centr. Bakt. Par., 1902, **8**, ii, 11, 42, 74, 107, 140, 171, 211, 248, 278, 309, 342, 367, 406.)
- E. O. Jordan.** Production of Fluorescent Pigment by Bacteria. (Proc. Amer. Physiol. Soc., 1898, 18; Amer. J. Physiol., **2**.)
- O. Kalischer.** Zur Biologie der peptonisierenden Milchbakterien. (Arch. Hygiene, 1900, **37**, 30.)
- J. H. Kastle.** The Inactivity of Lipase towards the Salts of Certain Acid Ethers, considered in the Light of the Theory of Electrolytic Dissociation. (Amer. Chem. J., 1902, **27**, 481.)
- J. H. Kastle and A. S. Loevenhart.** Lipase, the Fat-splitting Enzyme and the Reversibility of its Action. (Amer. Chem. J., 1900, **24**, 491.)
- J. H. Kastle and A. S. Loevenhart.** Nature of Certain of the Oxidising Ferments. (Amer. Chem. J., 1901, **26**, 539.)
- J. H. Kastle and O. M. Shedd.** Phenolphthalin as a Reagent for the Oxidising Ferments. (Amer. Chem. J., 1901, **26**, 526.)

- E. Kayser.** Études sur la fermentation lactique. (Ann. Inst. Pasteur, 1894, **8**, 737.)
- A. Kling.** Oxydation du propylglycol par les ferments oxydants. (Bull. Soc. Chim., 1901, [iii], **25**, 905.)
- W. Knecht.** Auswahl von Kohlehydraten durch verschiedene Hefen bei der alkoholischen Gärung. (Centr. Bakt. 1901, [ii], **7**, 161, 215.)
- J. W. T. Knox and A. B. Prescott.** Caffeine Compound in Kola. (J. Amer. Chem. Soc., 1897, **19**, 63.)
- A. Kossel and A. Matthews.** Zur Kenntniss der Trypsinwirkung. (Zeit. physiol. Chem., 1898, **25**, 190.)
- F. Krüger.** Beobachtungen über die quantitative Pepsinverdauung. (Zeit. Biol., 1901, **41**, 378, 467.)
- Kühne.** Ueber die Verdauung der Eiweissstoffe durch den Pankreassaft. (Virchow's Archiv, 1867, **39**, 130.)
- Kühne und Chittenden.** Ueber die nächsten Spaltungsproducte der Eiweisskörper. (Zeit. Biol., 1883, **19**, 184.)
- F. Kutscher.** Ueber das Antipepton. (Zeit. physiol. Chem., 1898, **25**, 195; 1898, **26**, 110.)
- F. Kutscher.** Chemische Untersuchungen über die Selbstgärung der Hefe. (Zeit. physiol. Chem., 1901, **32**, 59.)
- F. Kutscher.** Ueber das Hefetrypsin. (Zeit. physiol. Chem., 1901, **32**, 419.)
- F. Kutscher und J. Seemann.** Zur Kenntniss der Verdauungsvorgänge im Dunndarm. (Zeit. physiol. Chem., 1902, **34**, 528; 1902, **35**, 432.)
- J. B. V. Laborde.** Sur l'oxydase du *Botrytis cinerea*. (Compt. rend., 1898, **126**, 536.)
- F. Lafar.** Technische Mykologie. (Jena, 1897.)
- L. Langstein.** Zur Kenntniss der Endprodukte der peptischen Verdauung. (Beitr. chem. Physiol. Path., 1902, **1**, 507; 1902, **2**, 229.)
- D. Lawroff.** Zur Kenntniss der Chemismus der peptischen und tryptischen Verdauung der Eiweissstoffe. (Zeit. physiol. Chem., 1899, **26**, 513; 1901, **33**, 312.)
- O. Laxa.** Ueber die Spaltung des Butterfettes durch Mikroorganismen. (Arch. Hygiene, 1901, **41**, 119.)
- Sheridan Lea.** Some Notes on the Isolation of a Soluble Urea-ferment from the *Torula ureæ*. (J. Physiol., 1885, **6**, 136.)

- Sheridan Lea.** A Comparative Study of Artificial and Natural Digestions. (J. Physiol., 1890, **11**, 226.)
- K. B. Lehmann.** Ueber die Bildung von Oxydationsfermenten (Tyrosinase) durch Bakterien. (Chem. Centr., 1902, i, 770.)
- K. B. Lehmann und R. Neumann.** Bakteriologische Diagnostik. (München, 1899.)
- Lepine.** Sur la présence normale dans le chyle d'un ferment destructeur du sucre. (Compt. rend., 1890, **110**, 742.)
- Lepine.** Sur la production du ferment glycolytique. (Compt. rend., 1895, **120**, 139.)
- Lepine et Barral.** Sur le pouvoir glycolytique du sang et du chyle. (Compt. rend., 1890, **110**, 1314.)
- Lepine et Barral.** Sur la destruction du sucre dans le sang *in vitro*. (Compt. rend., 1891, **112**, 146.)
- Lepine et Barral.** Sur l'isolement du ferment glycolytique du sang. (Compt. rend., 1891, **112**, 411.)
- Lepine et Barral.** Sur les variations des pouvoirs glycolytique et saccharifiant du sang, et sur la localisation du ferment saccharifiant dans le sérum. (Compt. rend., 1891, **113**, 1014.)
- Lepinois.** Note sur les ferments oxydants de l'Aconit et de la Belladone. (J. Pharm., 1899, [vi], **9**, 49.)
- J. Lewkowitsch.** Optisch aktive Glycerinsäure und optisch aktive Milchsäure. (Ber. d. deutsch. chem. Ges., 1883, **16**, 2720.)
- T. Liebig.** Ueber die Erscheinungen der Gährung, Fäulniss und Verwesung und ihre Ursachen. (Ann. Chem. Pharm., 1839, **30**, 250, 363.)
- L. Lilienfeld.** Ueber Blutgerinnung. (Zeit. physiol. Chem., 1894, **20**, 89.)
- W. Lindemann.** Ueber die Löslichkeitsverhältnisse des Paracaseins im künstlichen Magensaft. (Virchow's Archiv, 1897, **149**, 51.)
- L. Lindet.** Sur l'oxydation du tannin de la pomme à cidre. (Compt. rend., 1895, **120**, 370.)
- P. Lindner.** Gärversuche mit verschiedenen Hefen- und Zuckerarten. (Chem. Centr., 1901, i, 56, 404.)
- A. R. Ling and J. L. Baker.** The Action of Diastase on Starch. (J. Chem. Soc. Trans., 1895, 702, 739; 1897, 508.)
- G. Linossier.** Sur le dédoublement de l'acide lactique inactif par les moisissures. (Bull. Soc. Chim. 1891, [iii], **6**, 10.)

- C. J. Lintner.** Studien über Diastase. (J. prakt. Chem., 1886, [ii], 34, 378; 1887, 36, 481.)
- C. J. Lintner.** Ueber das Vorkommen von Isomaltose im Biere und in der Wurze. (Zeit. ges. Brauw., 1891, 281.)
- C. J. Lintner.** Ueber die Vergährbarkeit der Isomaltose. (Zeit. ges. Brauw., 1892, 106.)
- C. J. Lintner.** Die Einwirkung von Diastase auf Isomaltose. (Chem. Centr., 1895, i, 91.)
- C. J. Lintner.** Ueber die Invertierung von Maltose und Isomaltose durch Hefe. (Chem. Centr., 1895, i, 271.)
- C. J. Lintner.** Zur Chemie der Stärke. (Chem. Zeit., 1897, 21, 737, 752.)
- C. J. Lintner und G. Düll.** Versuche zur Gewinnung der Isomaltose aus den Produkten der Stärkeumwandlung durch Diastase. (Zeit. angew. Chem., 1892, 363.)
- C. J. Lintner und G. Düll.** Ueber den Abbau der Stärke unter dem Einflusse der Diastasewirkung. (Ber. d. deutsch. chem. Ges., 1893, 26, 2533.)
- C. J. Lintner und G. Düll.** Ein zweites bei der Einwirkung von Diastase auf Stärke entstehendes Achroodextrin. (Chem. Centr., 1894, ii, 864.)
- C. J. Lintner und G. Düll.** Ueber den Abbau der Stärke durch die Wirkung der Oxalsäure. (Ber. d. deutsch. chem. Ges., 1895, 28, 1522.)
- O. Loew.** Catalase, a New Enzyme of General Occurrence. (U.S. Dep. Agric. Report., No. 68, 1901.)
- A. Maassen.** Fruchttätherbildende Bakterien. (Chem. Centr., 1899, ii, 1058.)
- A. Maassen.** Die Zersetzung der Nitrate und Nitrite durch die Bakterien. (Ein Beitrag zum Kreislauf des Stickstoffs in der Natur.) (Chem. Centr., 1901, ii, 820.)
- A. Magnus-Levy.** Ueber die Säurebildung bei der Autolyse der Leber. (Beitr. chem. Physiol. Path., 1902, 2, 261.)
- H. et A. Malbot.** Sur la formation de la mannite dans les vins. (Bull. Soc. Chim., 1894, [iii], 11, 413.)
- H. Malfatti.** Beitrag zur Kenntniss der peptischen Verdauung. (Zeit. physiol. Chem., 1900, 31, 43.)
- G. Malfitano.** La protéolyse chez l'*Aspergillus niger*. (Ann. Inst. Pasteur, 1900, 14, 60.)

- G. Malfitano.** Sur la protéase de *Aspergillus niger*. (Ann. Inst. Pasteur, 1900, **14**, 420.)
- G. Malfitano.** La bactériolyse de la bacteride charbonneuse. (Compt. rend., 1900, **131**, 295.)
- M. von Manassein.** Zur Frage von der alkoholischen Gährung ohne lebende Hefezellen. (Ber. d. deutsch. chem. Ges., 1897, **30**, 3061.)
- L. Mangin.** Eigenschaften und Reaktionen der Pektinsubstanzen. (Bied. Centr., 1894, **23**, 547.)
- W. Marckwald und A. McKenzie.** Ueber die Spaltung racemischer Verbindungen in die activen Componenten. (Ber. d. deutsch. chem. Ges., 1900, **33**, 208.)
- Marlière.** Sur la graine et spécialement sur l'endosperme du *Cerantonia siliqua*. (La Cellule, 1897, **13**, 7.)
- S. H. C. Martin.** Papain Digestion. (J. Physiol., 1884, **5**, 213.)
- S. H. C. Martin.** The Nature of Papain and its Action on Vegetable Proteids. (J. Physiol., 1885, **6**, 336.)
- V. Martinand.** Action de l'air sur le moût de raisin et sur le vin. (Compt. rend., 1895, **121**, 502.)
- V. Martinand.** Sur l'oxydation et la casse des vins. (Compt. rend., 1897, **124**, 512.)
- A. P. Matthews.** Division of Unfertilised Eggs. (Amer. J. Physiol., 1900, **4**, 343.)
- A. Mayer.** Untersuchungen über die Stärkekörner. Wesen und Lebensgeschichte der Stärkekörner der höheren Pflanzen. (Bied. Centr., 1899, **28**, 118.)
- A. Medvedeff.** Ueber die Oxydationskraft der Gewebe. (Pflüger's Archiv, 1896, **65**, 249.)
- A. Medvedeff.** Ueber die oxydativen Leistungen der thierischen Gewebe. (Pflüger's Archiv, 1900, **81**, 540.)
- R. Meissner.** Ueber das Auftreten und Verschwinden des Glykogens in der Hefezelle. (Centr. Bakt. Par., 1900, [ii], **6**, 517, 545.)
- v. Mering.** Ueber den Einfluss diastatische Fermente auf Stärke, Dextrin und Maltose. (Zeit. physiol. chem., 1881, **5**, 185.)
- L. B. Mendel and F. P. Underhill.** Products of Papain and Bromelin Proteolysis. (Proc. Amer. Physiol. Soc., 1900, xiii.)
- Miquel.** Sur le ferment soluble de l'urée. (Compt. rend., 1890, **111**, 397.)
- H. Mittelmeier.** Beiträge zur Kenntniss der diastatischen Zersetzung der Stärke. (Chem. Centr., 1895, ii, 163.)

- H. Mittelmeier.** Beitrag zum Studium der Einwirkung der Diastase auf die Stärke. (Chem. Centr., 1897, ii, 1010.)
- G. H. Morris.** The Hydrolysis of Maltose by Yeast. (J. Chem. Soc. Trans., 1895.)
- F. v. Musculus.** Remarques sur la transformation de la matière amylacée en glucose et dextrine. (Compt. rend., 1860, **50**, 785.)
- F. v. Musculus.** Nouvelle note sur la transformation de l'amidon en dextrine et glucose. (Compt. rend., 1862, **54**, 194.)
- F. v. Musculus.** De la dextrine. (Ann. chim. et phys., 1865, [iv], **6**, 177.)
- F. v. Musculus.** Sur le ferment de l'urée. (Compt. rend., 1876, **82**, 334.)
- F. v. Musculus und D. Gruber.** Ein Beitrag zur Chemie der Stärke. (Zeit. physiol. Chem., 1878, **2**, 177.)
- F. v. Musculus et D. Gruber.** Sur l'amidon. (Compt. rend., 1878, **86**, 1459, and Bull. Soc. Chim., 1878, **30**, 54.)
- F. v. Musculus und v. Mering.** Ueber die Umwandlung von Stärke und Glycogen durch Diastas, Speichel, Pancreas- und Leberferment. (Zeit. physiol. Chem., 1878, **2**, 403.)
- F. v. Musculus et v. Mering.** De l'action de la diastase de la saliva et du suc pancreatique sur l'amidon et sur le glycogène. (Bull. Soc. Chim., **31**, 105.)
- C. v. Naegeli.** Theorie der Gährung. München, 1879.
- W. Naegeli.** Beiträge zur Kenntniss der Stärkegruppe. Leipzig, 1874.
- O. Nasse.** Untersuchungen über die ungeformten Fermente. (Pflüger's Archiv, 1875, **11**, 138.)
- O. Nasse.** Bemerkungen zur Physiologie der Kohlehydrate. (Pflüger's Archiv, 1877, **14**, 477.)
- M. Nencki.** Zur Kenntniss der pankreatischen Verdauungsproducte des Eiweisses. (Ber. d. deutsch. chem. Ges., 1895, **28**, 560.)
- R. Neumeister.** Lehrbuch der physiologischen Chemie. Jena, 1897.
- R. Neumeister.** Bemerkungen zu Edward Buchner's Mittheilungen über "Zymase." (Ber. d. deutsch. chem. Ges., 1897, **30**, 2963.)
- P. Nobécourt et P. Merklin.** Présence d'un ferment dédoublant le salol dans les organes de l'homme et de divers animaux, ainsi que

- dans le lait de femme et de chienne. (Compt. rend. Soc. Biol., 1901, **53**, 148.)
- V. Oméliansky.** Sur la fermentation cellulosique. (Compt. rend., 1895, **121**, 653; 1897, **125**, 1131.)
- V. Oméliansky.** Ueber die Nitrifikation des organischen Stickstoffs. (Chem. Centr., 1899, ii, 347.)
- V. Oméliansky.** Ueber die Fermentation der Cellulose. (Chem. Centr., 1900, i, 918.)
- V. Oméliansky.** Ueber die Gärung der Cellulose. (Centr. Bakt. Par., 1902, [ii], **8**, 193, 225, 231, 257, 289, 321, 353, 385.)
- C. Oppenheimer.** Ferments and their Action. Translated by C. A. Mitchell. London, 1901.
- T. B. Osborne.** The Chemical Nature of Diastase. (J. Amer. Chem. Soc., 1895, **17**, 587.)
- T. B. Osborne and Campbell.** The Chemical Nature of Diastase. (J. Amer. Chem. Soc., 1896, **18**, 536.)
- H. Ost.** Ueber die Isomaltose. (Chem. Zeit., 1896, **20**, 761.)
- C. O'Sullivan.** On the Transformation Products of Starch. (J. Chem. Soc. Trans., 1872, 579; 1879, 770.)
- C. O'Sullivan and Tompson.** Invertase: a Contribution to the History of an Enzyme or Unorganised Ferment. (J. Chem. Soc. Trans., 1890, **57**, 834.)
- C. O'Sullivan.** The Hydrolytic Functions of Yeast. (J. Chem. Soc. Trans., 1892.)
- J. O'Sullivan.** Ueber die Geschwindigkeit der alkoholischen Gärung. (Bied. Centr., 1899, **25**, 863.)
- W. C. C. Pakes and W. Jollyman.** The Bacterial Decomposition of Formic Acid into Carbon Dioxide and Hydrogen. (J. Chem. Soc. Trans., 1901, **79**, 386.)
- L. Pasteur.** Mémoire sur la fermentation appelée lactique. (Ann. de chim. et phys., 1858, [iii], **52**, 407.)
- L. Pasteur.** Mémoire sur la fermentation alcoolique. (Ann. de chim. et phys., 1860, [iii], **58**, 323.)
- L. Pasteur.** Sur la fermentation visqueuse et la fermentation butyrique. (Bull. Soc. Chim., 1861, 30.)
- L. Pasteur.** Note sur un mémoire de M. Liebig, relatif aux fermentations. (Ann. de chim. et phys., 1872, [iv], **25**, 145.)

- L. Pasteur.** Faits nouveaux pour servir à la connaissance de la théorie des fermentations proprement dites. (Compt. rend., 1872, **75**, 784.)
- Payen.** Réaction de la diastase sur la substance amylacée dans différentes conditions. (Ann. chim. et phys., 1865, [iv], **4**, 286.)
- Payen et Persoz.** Mémoire sur la diastase, les principaux produits de ses réactions et leurs applications aux arts industriels. (Ann. chim. et phys., 1833, **53**, 73.)
- C. A. Pekelharing.** Ueber die Beziehung des Fibrinfermentes aus dem Blutserum zum Nucleoproteid, welches aus dem Blutplasma zu erhalten ist. (Chem. Centr., 1895, ii, 53.)
- A. Péré.** Fermentation lactique des corps sucrés. (Ann. Inst. Pasteur, 188, **12**, 63.)
- P. Petit.** Produits de saccharification de l'amidon par la diastase. (Compt. rend., 1897, **125**, 355.)
- P. Petit.** Sur les dextrines de saccharification. (Compt. rend., 1899, **128**, 1176.)
- M. Pfaundler.** Zur Kenntniss der Endprodukte der Pepsinverdauung. (Zeit. physiol. Chem., 1900, **30**, 90.)
- E. P. Pick.** Zur Kenntniss der peptischen Spaltungsprodukte des Fibrins. (Zeit. physiol. Chem., 1899, **28**, 219.)
- E. P. Pick und K. Spiro.** Ueber gerinnungshemmende Agentien im Organismus höhere Wirbelthiere. (Zeit. physiol. Chem., 1900, **31**, 235.)
- P. M. Popoff.** Ueber die Einwirkung von eiweissverdauenden Fermenten auf die Nucleinstoffe. (Zeit. physiol. Chem., 1894, **18**, 533.)
- P. Portier.** Sur la glycolyse des différents sucres. (Compt. rend., 1900, **131**, 1217.)
- H. Pottevin.** Sur la saccharification de l'amidon par l'amylase du malt. (Compt. rend., 1898, **126**, 1218.)
- H. Pottevin.** La saccharification de l'amidon. (Ann. Inst. Pasteur, 1899, **13**, 665.)
- H. Pottevin.** Sur la maltodextrine. (Ann. Inst. Pasteur, 1899, **13**, 728.)
- H. Pottevin.** Sur l'isomaltose. (Ann. Inst. Pasteur, 1899, **13**, 796.)
- H. Pottevin.** La Tannase. Diastase dédoublant l'acide gallotannique. (Compt. rend., 1901, **131**, 1215.)

- H. Pottevin.** Sur la constitution du gallotannin. (Compt. rend., 1901, **132**, 704.)
- M. E. Pozzi-Escot.** Sur les hydrogénases du sang et les propriétés catalytique de la fibrine. (Bull. Soc. Chim., 1902, [iii], **27**, 459.)
- M. E. Pozzi-Escot.** Sur une importante cause d'erreur dans la recherche des diastases. (Bull. Soc. Chim., 1902, [iii], **27**, 460, and Compt. rend., 1902, **134**, 479.)
- M. E. Pozzi-Escot.** Contribution à l'étude des hydrogenases; nouveau cas d'hydrogenation diastatique. (Bull. Soc. Chim., 1902, [iii], **27**, 346.)
- M. E. Pozzi-Escot.** Catalytic Properties of the Hydrogenases: Identification of Loew's Catalase with de Rey-Pailhade's Philothion. (Bull. Soc. Chim., 1902, [iii], **27**, 280.)
- E. Prior.** Ueber ein drittes Diastase-Achroodextrin und die Isomaltose. (Chem. Centr., 1896, ii, 86.)
- E. Prior and H. Schulze.** Beiträge zur Physik der Gährung. (Zeit. angew. Chem., 1901, **14**, 208.)
- E. Prior and D. Wiegmann.** Darstellung und Eigenschaften des Diastase-Achroodextrins III. (Zeit. angew. Chem., 1900, 464.)
- T. Purdie and J. W. Walker.** Optically Active Ethoxysuccinic Acid. (J. Chem. Soc. Trans., 1892.)
- F. Reich-Herzberge.** Ueber die Einwirkung von Trypsin auf Leim. (Zeit. physiol. Chem., 1901, **34**, 119.)
- F. Reinitzer.** Ueber die wahre Natur des Gummifermentes. (Zeit. physiol. Chem., 1890, **14**, 453.)
- F. Reinitzer.** Ueber das zellwandlösende Enzym der Gerste. (Zeit. physiol. Chem., 1897, **23**, 175.)
- E. Reiss.** Ueber die in den Samen als Reservestoff abgelagerte Cellulose und eine daraus erhaltene neue Zuckerart, die "Seminose." (Ber. d. deutsch. chem. Ges., 1889, **22**, 610.)
- J. de Rey-Pailhade.** Rôles respectifs du philothion et de la laccase dans les graines en germination. (Compt. rend., 1895, **121**, 1162.)
- J. de Rey-Pailhade.** Existence du corps proteique prévu par M. G. Bertrand dans le constitution des oxydases. (Bull. Soc. Chim., 1897, [iii], **17**, 756.)
- Ringer.** The Influence of Certain Salts upon the Act of Clotting. (J. Physiol., 1890, **11**, 369.)
- Ringer.** Regarding the Action of Lime Salts on Casein and on Milk. (J. Physiol., 1890, **11**, 464.)

- Ringer.** Further Observations on the Behaviour of Caseinogen. (J. Physiol., 1891, **12**, 164.)
- F. Röhmman.** Zur Kenntniss der bei der Trypsinverdauung aus dem Casein entstehende Producte. (Ber. d. deutsch. chem. Ges., 1897, **30**, 1978, and 1898, **31**, 2188.)
- G. W. Rolfe and G. Defreu.** Analytical Investigation of the Hydrolysis of Starch by Acids. (J. Amer. Chem. Soc., 1896, **18**, 869; 1897, **19**, 261.)
- G. Roux.** Sur une oxydase productrice de pigment, secrété par le coli-bacille. (Compt. rend., 1899, **128**, 693.)
- N. Sacharoff.** Die Demonstration der in Band 24, Nr. 18-19 des Centralblattes für Bakteriologie beschriebenen Versuche über Enzyme. (99, i, 131, 941.) (Chem. Centr., 1899, ii, 723.)
- S. Salaskin.** Ueber die Bildung des Leucinimids bei der peptischen und tryptischen Verdauung des Oxyhämoglobins resp. des Globins. (Zeit. physiol. Chem., 1901, **32**, 592.)
- E. u. H. Salkowski.** Ueber die Bildung von Hydrozimmersäure bei der Pankreasverdauung. (Ber. d. deutsch. chem. Ges., 1879, **12**, 107.)
- E. u. H. Salkowski.** Weitere Beiträge zur Kenntniss der Faulnissprodukte des Eiweiss. (Ber. d. deutsch. chem. Ges., 1897, **12**, 648.)
- E. u. H. Salkowski.** Zur Kenntniss der Pankreasverdauung. (Zeit. physiol. Chem., 1878-79, **2**, 420.)
- E. u. H. Salkowski.** Zur Kenntniss der Eiweissfäulniss. Ueber die Bildung des Indols und Skatols. (Zeit. physiol. Chem., 1883-84, **8**, 417.)
- E. u. H. Salkowski.** Ueber das Verhalten der Skatolcarbonsäure im Organismus. (Zeit. physiol. Chem., 1885, **9**, 23.)
- E. Salkowski.** Ueber das eiweisslösende Ferment der Fäulnissbakterien und seine Einwirkung auf Fibrin. (Zeit. Biol., 1889, **25**, 92.)
- E. Salkowski.** Ueber das Verhalten des Caseins zu Pepsinsalzsäure. (Pfüger's Archiv, 1896, **63**, 401.)
- E. Salkowski.** Kleinere Mittheilungen. (Zeit. physiol. Chem., 1897, **27**, 297.)
- E. Salkowski.** Ueber ein phosphorhaltige Säure aus Casein und deren Eisenverbindung. Vorläufige Mittheilung. (Chem. Centr., 1901, i, 225.)

- E. Salkowski.** Behaviour of the Phosphorus in the Digestion of Casein. (Exper. Stat. Record., **5**, 428.)
- E. Salkowski und M. Hahn.** Ueber das Verhalten des Phosphors im Casein bei der Pepsinverdauung. (Pflüger's Archiv, 1894, **59**, 225.)
- E. Salkowski und Y. Yamagiva.** Zur Kenntniss des Oxydationsferments der Gewebe. (Virchow's Archiv, 1897, **147**, 1.)
- R. H. Saltet** (and C. S. Stockois). Ueber Reduktion von Sulfaten in Brackwasser durch Bakterien. (Centr. Bakt. Par. [ii], **6**, 648, 695.)
- J. Sarthou.** Sur une oxydase retirée du *Schinus Molle*. (J. Pharm., 1900, [vi], **11**, 482.)
- J. Sarthou.** Du rôle que parait jouer le fer dans la schinoxydase. (J. Pharm., 1900, [vi], **11**, 583.)
- W. W. Sawjaloff.** Zur Theorie der Eiweissverdauung. (Pflüger's Archiv, 1901, **85**, 171.)
- E. A. Schäfer.** Experiments on the Conditions of Coagulation of Fibrinogen. (J. Physiol., 1895, **17**, 18, Proc. Physiol. Soc.)
- E. A. Schäfer.** A Text Book of Physiology. 1898.
- F. Schardinger.** Ueber eine neue optisch active Modification der Milchsäure, durch bacterielle Spaltung des Rohrzuckers erhalten. (Monatsh., 1890, **11**, 545.)
- F. Schardinger.** Ueber die Gärprodukte eines schleimbildenden Bacillus in Rohrzuckerlösungen und die Zusammensetzung eines aus dem Schleime isolierten Kohlenhydrates. (Centr. Bakt. Par., 1902, [ii], **8**, 144, 175.)
- A. Schattenfroh.** Biologisches Verhalten und Verbreitung des beweglichen Buttersäurebacillus. (Arch. Hygiene, 1902, **42**, 251.)
- A. Schattenfroh und R. Grassberger.** Weitere Mittheilungen über Buttersäuregärung. (Chem. Centr., 1899, ii, 1060.)
- C. Scheibler und H. Mittelmeier.** Studien über die Stärke. (Ber. d. deutsch. chem. Ges., 1893, **26**, 2930, 3060, 3473.)
- A. Schiffer.** Ueber die nicht krystallisierbaren Produkte der Einwirkung der Diastase auf Stärke. (Chem. Centr., 1892, ii, 825.)
- Schloesing et Müntz.** Recherches sur la nitrification par les ferments organisés. (Compt. rend., 1877, **84**, 301; 1877, **85**, 1018; 1878, **86**, 892.)

- Schloesing et Müntz.** Recherches sur la nitrification. (Compt. rend., 1879, **89**, 891, 1074.)
- A. Schmidt.** Weiteres über den Faserstoff und die Ursachen seiner Gerinnung. (Arch. Anat. Phys., 1862, **428**, 533.)
- A. Schmidt.** Neue Untersuchungen über die Faserstoffgerinnung. (Pflüger's Archiv, 1872, **6**, 445, 457.)
- A. Schneegans.** Betulase, ein in *Betula lenta* enthaltenes Ferment. (Chem. Centr., 1897, **i**, 326.)
- C. F. Schoenbein.** Ueber die katalytische Wirksamkeit organischer Materien und deren Verbreitung in der Pflanzen- und Thierwelt. (J. für prakt. Chem., 1863, **89**, 323.)
- P. Schützenberger.** On Fermentation. London, 1891.
- E. Schunck.** On Rubian and its Products of Decomposition. (Phil. Trans., 1851, 433.)
- E. Schunck.** Erythrozyme. (Phil. Trans., 1853, 74.)
- E. Schunck.** Alkoholische Gährung ohne Hefezellen. (Ber. d. deutsch. chem. Ges., 1898, **31**, 309.)
- C. Schweitzer.** Zur Kenntniss der caffen- und theobrominhaltigem Glykoside in den Pflanzen. (Chem. Centr., 1898, **ii**, 217.)
- Seegen.** Ueber die Umwandlung von Glycogen durch Speichel- und Pancreasferment. (Pflüger's Archiv, 1879, **19**, 106.)
- Seegen und Kratschmer.** Zur Kenntniss der saccharifizirenden Fermente. (Pflüger's Archiv, **14**, 593.)
- F. & L. Sestini.** Ammoniacal Fermentation of Uric Acid. (Gazzetta, 1896, **26**, **ii**, 92.)
- G. Sharp.** Papainverdauung: Völlige Abwesenheit von Pepton. Ueber Papain. (Chem. Centr., 1894, **i**, 512.)
- M. Siegfried.** Ueber Antipepton. (Ber. d. deutsch. chem. Ges., 1900, **33**, 2851; Zeit. physiol. Chem., 1902, **35**, 164.)
- M. Siegfried.** Ueber Antipepton. (Zeit. physiol. Chem., 1899, **27**, 335.)
- E. von Sigmund.** Beiträge zur Einwirkung der Diastase auf unverkleisterte Stärke. (Chem. Centr., 1897, **ii**, 614.)
- L. Sostegni und A. Sannino.** Ueber die Erzeugung von Schwefelwasserstoff bei der Alkoholgärung. (Chem. Centr., 1890, **ii**, 112.)
- Soxhlet.** Beiträge zur physiologischen Chemie der Milch. (J. prakt. Chem., 1872, **6**, 1.)
- W. Spitzer.** Die zuckerzerstörende Kraft des Blutes und der Gewebe. (Pflüger's Archiv, 1895, **60**, 303.)

- W. Spitzer.** Die Bedeutung gewisser Nucleoproteide für die oxydative Leistung der Zelle. (Pflüger's Archiv, 1897, **67**, 615.)
- W. Spitzer.** Weitere Beobachtungen über die oxydativen Leistungen thierischer Gewebe. (Pflüger's Archiv, 1898, **71**, 596.)
- W. Spitzer.** Die Ueberführung von Nucleinbasen in Harnsäure durch die sauerstoffübertragende Wirkung von Gewebeauszügen. (Pflüger's Archiv, 1899, **76**, 192.)
- E. Stadelmann.** Bildung von Ammoniak bei Pancreasverdauung von Fibrin. (Zeit. Biol., 1888, **24**, 261.)
- E. Stadelmann.** Ueber das beim tiefen Zerfall der Eiweisskörper entstehende Proteinchromogen. (Zeit. Biol., 1889, **26**, 491.)
- G. N. Stewart and T. Sollman.** Proteids of Muscle. (J. Physiol., 1899, **24**, 427.)
- A. Stutzer und R. Hartleb.** Die Salpeterpilz. (Chem. Centr., 1897, ii, 622.)
- A. Stutzer und R. Hartleb.** Die Zersetzung von Cement unter dem Einflusse von Bakterien. (Zeit. angew. Chem., 1899, 402.)
- W. Syniewski.** Ueber die lösliche Stärke. (Ber. d. deutsch. chem. Ges., 1897, **30**, 2415; 1898, **31**, 1791.)
- W. Syniewski.** Ueber die Constitution der Stärke. (Annalen, 1899, **309**, 282; Chem. Centr., 1902, ii, 984.)
- G. Tammann.** Die Reactionen der ungeformten Fermente. (Zeit. physiol. Chem., 1892, **16**, 271.)
- G. Tammann.** Zur Wirkung ungeformter Fermente. (Zeit. physikal. Chem., 1895, **28**, 426.)
- C. Tanret.** Sur la picéine, glucoside des feuilles du sapin épicéa (*Pinus picea*). (Compt. rend., 1894, **119**, 80.)
- C. et G. Tanret.** Sur le rhamninose. (Compt. rend., 1899, **129**, 725.)
- C. et G. Tanret.** Sur la rhamninase et la xanthorhamnine. (Bull. Soc. Chim., 1899, [iii], **21**, 1073.)
- W. Tappeiner.** Ueber Celluloseverdauung. (Ber. d. deutsch. chem. Ges., 1882, **15**, 999; 1883, **16**, 1734.)
- G. Tate.** The Fermentation of Dextrose, Rhamnose and Mannitol by a Lævolactic Ferment. (J. Chem. Soc. Trans., 1893, 1263.)
- M. C. Tebb.** Hydrolysis of Glycogen. (J. Physiol., 1898, **22**, 423.)
- G. Tolomei.** Soluble Ferment present in Wine. (Real. Accad. Linc., 1896, [v], **5**, i, 52.)

- L. v. Udransky und E. Baumann.** Ueber das Vorkommen von Diaminen, sogenannten Ptomainen bei Cystinurie. (Zeit. physiol. Chem., 1889, **13**, 562.)
- C. Ulrich.** Ueber die Isomaltose. (Chem. Zeit., 1895, **19**, 1523.)
- F. Umber.** Die Spaltung des krystallinischen Eier- und Serumalbumins sowie des Serumglobulins durch Pepsinverdauung. (Zeit. physiol. Chem., 1898, **25**, 258.)
- Van Rijn.** Die Glykoside. Berlin, 1900.
- Van Tieghem.** Sur la fermentation ammoniacale. (Compt. rend., 1864, **58**, 533.)
- C. Vincent et Délachanal.** Preparation biologique du levulose au moyen de la mannite. (Compt. rend., 1897, **125**, 716.)
- H. Vohl.** Ueber die Qualität der aus dem Inosit entstehenden Milchsäure. (Ber. d. deutsch. chem. Ges., 1876, **9**, 984.)
- R. Warington.** On Nitrification. (J. Chem. Soc. Trans., 1878, **44**; 1879, 429; 1884, 637; 1891, 484; 1896; Proc., 1900, **16**, 65.)
- C. Wehmer.** Préparation d'acide citrique de synthèse par la fermentation du glucose. (Compt. rend., 1893, **117**, 332.)
- C. Wehmer.** Ueber zwei weitere freie Citronensäure bildende Pilze. (Chem. Zeit., 1897, **21**, 1022.)
- C. Wehmer.** Ueber säureliebende Pilze. (Bied. Centr., 1898, **27**, 284.)
- C. Wehmer.** Ueber die physiologische Ungleichwertigkeit der Fumar- und Maleinsäure und die antiseptische Wirkung der letzteren. (Bied. Centr., 1898, **27**, 285.)
- R. Weil.** Die Entstehung des Solanins in den Kartoffeln als Product bakterieller Einwirkung. (Arch. Hygiene, 1900, **38**, 330.)
- E. Weinland.** Ueber die Bildung von Glykogen nach Galaktosefütterung. (Zeit. Biol., 1900, **40**, 374.)
- F. A. Went.** Die Schwefelkohlenstoffbildung durch *Schizophyllum lobatum*. (Chem. Centr., 1896, ii, 939.)
- Th. Weyl.** Fäulniss von Fibrin, Amyloid und Leim. (Zeit. physiol. Chem., 1877-78, **1**, 339.)
- Th. Weyl.** Spaltung von Tyrosin durch Fäulniss. (Zeit. physiol. Chem., 1879, **3**, 312.)
- S. Winogradsky.** Recherches sur les organismes de la nitrification. (Ann. Inst. Pasteur, 1890, **4**, 213, 257, 760; 1891, **5**, 92, 577.)

- K. Wolff.** Denitrifikation und Gärung. (Chem. Centr., 1900, i, 52.)
- G. Sims Woodhead.** Bacteria and their Products. London, 1891.
- Wooldridge.** On the Coagulation of the Blood. (J. Physiol., 1883, 4, 226, 367.)
- A. Wroblewski.** Zur Kenntniss des Pepsins. (Zeit. physiol. Chem., 1895, 21, 1.)
- A. Wroblewski.** Ueber die lösliche Stärke. (Ber. d. deutsch. chem. Ges., 1897, 30, 2108.)
- A. Wroblewski.** Yeast Extract. (Bull. Acad. Sci. Cracow., 1901, 94.)
- A. Würtz.** Sur la papaïne; contribution à l'histoire des ferments solubles. (Compt. rend., 1880, 90, 1379.)
- A. Würtz.** Nouvelle contribution à l'histoire des ferments solubles. (Compt. rend., 1880, 91, 787.)
- Yoshida.** Chemistry of lacquer. (J. Chem. Soc. Trans., 1883, 43, 472.)
- N. Zelinsky.** Sulphydic Fermentation in the Black Sea. (J. Russ. Chem. Soc., 1894, 25, 298.)
- E. Zunz.** Die fractionierte Abscheidung der peptischen Verdauungsprodukte mittelst Zinksulfat. (Zeit. physiol. Chem., 1899, 27, 219.)
- E. Zunz.** Ueber den quantitativen Verlauf der peptischen Eiweiss-spaltung. (Zeit. physiol. Chem., 1899, 28, 132.)

INDEX OF AUTHORS

- Abel, 93
 Abelous, 88, 93, 102
 Ampola, 97
 Armstrong, F. F., 2, 38, 39, 60
 Armstrong, H. E., 66
 Arthus, 109
 Astier, 44

 Babcock, 129
 Baeyer, 62, 65, 66
 Baker, 12
 Baldwin, 128
 Balke, 126
 Banning, 64
 Basile, 93
 Bau, 32, 38
 Baumann, 136, 138
 Béchamp, 44
 Becher, 41
 Berg, 56
 Bernard, 71, 122
 Berthelot, 32, 33, 43, 46, 74
 Bertrand, 30, 81, 82, 83, 84, 85, 91
 Berzelius, 46
 Bevan, 29
 Beyerinck, 54, 92, 101, 102
 Biarnès, 88
 Biffi, 126
 Biginelli, 93
 Biondi, 129
 Blank, 128
 Blondeau, 61
 Boekhout, 65
 Bokay, 127
 Bokorny, 70, 132
 Boland, 103
 Bordoni-Uffreduzzi, 103
 Bouffard, 48, 86
 Boullay, 42, 43
 Bourquelot, 26, 28, 30, 31, 32, 37,
 38, 84, 85, 130

 Boutron, 51
 Boutroux, 80, 81
 Bréaudat, 54, 87
 Brieger, 138, 139, 140
 Brown, A. J., 23, 34, 47, 79, 80
 Brown, H. T., 6, 8, 9, 10, 12, 14, 19,
 21, 23, 24, 25, 26, 34
 Brücke, 6, 25, 115
 Buchanan, 105, 106
 Buchner, 2, 46, 47, 70
 Burri, 76, 97
 Buttenberg, 93

 Camus, 74, 113
 Carles, 87
 Cavazzani, 88
 Cazeneuve, 86
 Chittenden, 116, 117, 119, 127, 131
 Claflin, 62
 Cohnheim, 128
 Cole, 124
 Corvisart, 122
 Cossettini, 91
 Cramer, 108
 Cross, 29
 Czapek, 29

 Danilewsky, 122
 Davy, 78
 Delachanal, 81
 Demoussy, 96
 Denis, 106
 Desmazières, 44
 Detmer, 72
 Döbereiner, 78
 Dokkum, 139
 Drechsel, 124
 Dubois, 89
 Dubourg, 93
 Dubrunfaut, 43
 Düll, 10, 11, 25, 27

- Dumas, 42, 43, 75
 Dunlop, 56
 Dunstan, 56
 Dupré, 138
 Dziergowski, 125

 Effront, 27, 28
 Ehrenberg, 138
 Ehrlich, 87
 van Ekenstein, 28
 Emerson, 125
 Emmerling, 2, 36, 48, 52, 65, 70,
 81, 93, 101, 130, 132
 Erdmann, 102

 Fabroni, 44
 Ficquet, 68
 Fischer, E., 2, 10, 12, 15, 16, 31, 32,
 38, 39, 48, 51, 52, 57, 58, 59, 60,
 69
 Fitz, 100, 101
 Fordos, 103
 Fränkel, 116
 Frankland, 45, 63, 67, 68, 95
 Frederikse, 108
 Frémy, 29
 Freudenreich, 37, 129
 Frew, 67
 v. Fürth, 86, 112, 113

 Gadamer, 55
 Galli-Valerio, 93
 Ganz, 29
 Garcia, 138
 Gautier, 140
 Gay-Lussac, 42, 43, 44
 Gayon, 93
 Gérard, 76, 93, 102
 Geret, 130
 Glendinning, 23
 Gley, 113
 Gmelin, 124
 Gonnermann, 102
 Gosio, 93
 Grassberger, 100
 Green, 26, 106
 Griessmayer, 6, 25
 Griffiths, 103
 Grimbert, 68, 102
 Gruber, 6, 7, 25
 Grüss, 87
 Gulewitsch, 128

 Hahn, 115, 130
 Halliburton, 107, 111, 112, 113, 131

 Hammarsten, 106, 108, 109, 111
 Hanriot, 2, 73, 74
 Harcourt, 33
 Harden, 69
 Harlay, 120, 131
 Hart, 127
 Hartleb, 96
 Hasebroek, 127
 Hazewinkel, 54
 Hedin, 124
 Helm, 103
 van Helmont, 41, 75
 Henneberg, 80
 Henri, 34
 Henry, 57
 Hérissé, 28, 30, 32, 38, 53, 54, 130
 Heron, 6, 25
 Herzfeld, 7, 8, 25
 Hewson, 105, 106
 Hilger, 63
 Hill, 2, 35, 36
 Hirschler, 125
 Hopkins, 124
 Hoppe-Seyler, 46, 77
 Horne, 107

 Jackson, 27
 Jacoby, 130
 Jalowetz, 12
 Jegunow, 90
 Jollyman, 101
 Jones, 138
 Jordan, 104

 Kalischer, 131
 Kastle, 2, 72, 73
 Kayser, 68
 Kiliani, 26
 Knecht, 48
 v. Knieriem, 79
 Knox, 57
 König, 93
 Kolbe, 122
 Kossel, 125, 127
 Krüger, 121
 Kühne, 111, 115, 116, 121, 122, 123,
 135, 136
 Kutscher, 124, 126, 127, 130
 Kützing, 45, 46, 78

 Langstein, 116, 120
 Lankester, 103
 Latour, 45, 46
 Lavoisier, 42, 78
 Lawroff, 121

- Laxa, 72
 Lea, 75, 125
 Lehmann, 86
 Lepierre, 104
 Levene, 128
 Lewkowitsch, 67
 Liebig, 45, 46, 51, 79, 110
 Lilienfeld, 108
 Lindemann, 118
 Lindet, 86
 Lindner, 32, 38
 Ling, 12
 Linossier, 68
 Lintner, 10, 11, 12, 25
 Loevenhart, 2, 72
 Loew, 3, 92, 97
 Loewenhoek, 44
 Loiseau, 31
 Ludwig, 84

 Maasen, 97, 102
 MacGregor, 63, 68
 Magnus-Levy, 130
 Malbot, 93
 Malfatti, 120
 Malfitano, 130
 Mallèvre, 30
 Manassein, 47
 Marcano, 131
 Marckwald, 70
 Marlière, 28
 Martin, 131
 Matthews, 127, 128
 Mayer, 79, 115
 Mazé, 72
 Medvedeff, 88
 Meissner, 115, 122
 Mendel, 117, 132
 Meyer, 23
 Millar, 14, 25
 Miquel, 75
 Mitscherlich, 46
 Mittelmeier, 11, 13, 16, 25
 Molisch, 90
 Mouoyer, 65
 Morris, 8, 9, 10, 12, 14, 21, 25, 26
 Müller, 34
 Müntz, 72, 95
 Musculus, 6, 7, 25, 75

 Naegeli, 9, 46
 Nasse, 23, 61
 Nef, 65
 Nencki, 124, 125, 127, 128, 136, 138

 Neumeister, 116, 117, 120, 124, 129, 136, 137
 Niebel, 31
 Nuttall, 135

 Oméliansky, 29, 96, 140
 Oppenheimer, 61, 89
 Ost, 12
 O'Sullivan, 6, 7, 10, 12, 25, 33, 34, 47

 Pakes, 101
 Pappenheim, 122
 Pasteur, 43, 44, 45, 46, 47, 61, 61, 67, 70, 75, 79, 134
 Payen, 6, 25
 Pekelharing, 107, 108
 Péré, 69
 Persoon, 78
 Persoz, 6, 25
 Pfaunder, 120
 Pick, 119
 Pickering, 34
 Pohl, 88
 Pottevin, 12, 13, 25
 Pozzi-Escot, 91, 92, 94
 Prescott, 57
 Prior, 47, 48
 Purdie, 70
 Purkinje, 122

 Reiser, 130
 Reiss, 29
 Rey-Pailhade, 91
 Ribaut, 102
 Ringer, 107, 111
 Robiquet, 51, 52
 Roux, 86
 Russell, 129

 Sabanejeff, 117
 Sachs, 72
 Sainsbury, 107
 Saliskin, 125
 Salkowski, 91, 118, 129, 135, 136, 137
 Saltet, 92
 Sannino, 91
 Sarthou, 86
 de Saussure, 42
 Sawjaloff, 121
 Schäfer, 108
 Schardinger, 34, 63, 68
 Schattenfroh, 100
 Scheele, 62

- Scheibler, 11, 16, 25, 32, 65
Scheurlen, 103
Schiffer, 11
Schloesing, 95
Schmidt, 43, 106
Schmiedeberg, 87
Schneider, 86
Schönbein, 2, 84, 85
Schulze, 47, 132
Schunck, 47, 55
Schützenberger, 72, 123
Schwann, 45, 46
Schweitzer, 57
Seegen, 27, 89
Selmi, 111, 138
Sestini, 76
Siegfried, 126
Solley, 127
Sollmann, 113
Sostegni, 91
Soxhlet, 110
Spiro, 124
Spitzer, 88
Stadelmann, 93, 124, 125
Stahl, 41, 45
Stewart, 113
Strecker, 100
Stryzowski, 93
Stutzer, 96, 97
Syniewski, 20, 21, 25

Tanret, 56
Tate, 63, 68
Tebb, 27
Thénard, 42, 44
Thierfelder, 135

Tiedemann, 124
van Tieghem, 75, 76
Tollens, 29, 31
Tolomei, 87
Tompson, 12, 33, 34
Traube, 46

Udransky, 138
Ulpiani, 97
Ulrich, 12
Umber, 119
Underhill, 132

Valentine, 41
Vaughan, 139
Vincent, 81
Vohl, 63

Wälchi, 127
Walker, 70
Ward, 56
Warington, 95
Wehmer, 63, 64
Went, 102
Will, 130
Willis, 41, 45
Winogradsky, 89, 90, 95
Wöhler, 51
Wroblewsky, 115
Wurtz, 131, 132

Yoshida, 82

Zelinsky, 92
Zopf, 103
Zunz, 119, 120

INDEX OF SUBJECTS

- Acetic acid, 78
 Achroodextrin, 6, ff.
 Acids, acetic, 78, ff.
 —, amino, 70, 114
 —, butyric, 99, ff.
 —, citric, 63
 —, formic, 101
 —, gluconic, 80
 —, lactic, 61-63, 68, ff.
 —, malic, 101
 —, mucic, 99
 —, oxalic, 64
 —, pectic, 30
 —, pectosic, 30
 —, propionic, 99, ff.
 —, urushic, 82
 Acid amides, hydrolysis of, 102
 Acid fermentation, 41, 61, 78, 99
 Æsculin, 53
 Albumin, 114-133
 Albumoses, 116, ff.
 Alcoholic fermentation, 41-48
 Amidulin, 11
 Amino-acids, 70, 114, ff.
 Amino-oxydases, 87
 Amphopeptone, 116, ff.
 Amygdalin, 51-53
 Amylodextrin, 9, ff.
 Amylogen, 21, ff.
 Anærobic organisms, 134
 Antipeptone, 116, ff., 126
 Arabic acid, 29
 Arabinose, 29, 64, 67, 81
 Arbutin, 53
 Aromatic compounds, 135
 Arsenic compounds, 92, 93
Aspergillus niger, 31, 32, 38, 53,
 54, 70, 130
 Autodigestion, 129

Bacillus acidi lævolactici, 68

Bacillus amylobacter, 99
 — *butylicus*, 101
 — *coli communis*, 69
 — *cyanofuscus*, 102
 — *cyanogenus*, 102
 — *desulfuricans*, 92
 — *esterificans fluorescens*, 102
 — *ethaceticus*, 67
 — *ethacetosuccinus*, 68
 — *fluorescens liquefaciens*,
 104, 130
 — *fluorescens non-liquefaciens*,
 104
 — *lactis aërogenes*, 93
 — *pyocyaneus*, 102
 — *tartricus*, 68, 102
 Bacteriofluorescin, 104
 Bacteriolysis, 131
 Bacteriopurpurin, 103
Bacterium aceti, 80
 — *denitrificans*, 97
 — *hydrosulfureum ponticum*,
 92
 — *indigonaceum*, 103
 — *of iron*, 89
 — *prodigiosum*, 103
 — *of sorbose*, 81
 — *termo*, 93
 — *ureæ*, 75
 — *violaceum*, 103
 — *xylinum*, 80
 Betulase, 56
 Blood-clotting, 105-109
 Bromelin, 131
 Butyl alcohol, 101
 Butyric acid, 99, ff.
 — ferment, 93
 — ester, 72, 73

 Cacaonin, 57
 Calcium formate, 101

- Cane sugar, 33, 42, 131
 Caroubin, 27
 Caroubinase, 28
 Casein, 110, 118, 126, 131
 Casse, 86
 Catalase, 92
 Cellulose, 28, 80
 Cerebrospinase, 88
 Chromogenic micro-organisms, 84,
 102
 Coagulation, 105-183
 Coniferin, 53
 Creatine, 102
 Cytase, 28

 Daphnin, 54
 Denitrification, 97
 Dextran, 65
 Dextrin, 6-27, 64
 Dextrinic acid, 17
 Diastase, 5-25, 34
 Diastase hydrolysis. tabulated
 results of, 25
 ———, velocity of, 23
 Dihydroxyacetone, 48, 81
 Disaccharides, 31-39
 Dystropodextrin, 27

 Elaterase, 56
 Emulsin, 16, 36-40, 51-55
 Endotrypsin, 130
 Erepsin, 128
 Erythrodextrin, 6, ff.
 Erythrozyme, 47, 56
 Esters, 71-74

 Fats, 71
 Ferments, butyric, 93
 ———, caroubinase, 28
 ———, cerebro-spinase, 88
 ———, cytase, 28
 ———, diastase, 5-27
 ———, elaterase, 56
 ———, emulsin, 16, 36, 39, 51, 54
 ———, erythrozyme, 47, 56
 ———, fibrin, 94, 105-109
 ———, galactase, 129
 ———, gaultherase, 56
 ———, glucase, 16, 35
 ———, glycolytic, 89
 ———, hadromase, 29
 ———, inulase, 26
 ———, invertin, 31
 ———, isatase, 54
 ———, laccase, 82-86, 91
 Ferments, lactase, 36, 38, 39
 ———, lipase, 36, 71, 72
 ———, luciferase, 89
 ———, maltase, 16, 35
 ———, mannitol, 80
 ———, melibiase, 32, 38
 ———, melizitase, 32
 ———, myosin, 112
 ———, myrosin, 55
 ———, oxoxygenase, 86
 ———, pectase, 29
 ———, pectinase, 30
 ———, philothion, 91
 ———, raffinase, 32
 ———, reductase, 94
 ———, rennin, 110
 ———, schinoxidase, 86
 ———, seminase, 28
 ———, steapsin, 71
 ———, synaptase, 51
 ———, tannase, 56
 ———, trehalase, 38
 ———, tyrosinase, 85
 ———, urease, 75
 ———, vesiculase, 113
 ———, yeast, 44-50
 ———, zymase, 47
 ———, chromogenic action of.
 84, 102
 ———, reducing, 91-94
 ———, selective action of, 67-70
 Fermentation, acid, 41, 61, 99
 ———, acid and alcoholic, 41
 ———, alcoholic, 41-48
 ———, proteolytic, 114-133
 ———, and putrefaction, 40
 ———, synthetic, 35, 36
 Fibrin, 116, 119
 Fibrinogen, 105-109
 Fusel oil, 44

 Galactase, 129
 Galactose, 29, 31, 36, ff., 48, 49
 Galactosides, 58
 Galactosidogalactose, 39
 Galactosidoglucose, 39
 Gastric digestion, 114-121
 Gaultherin, 56
 Gaultherase, 56
 Gelatine, 127, 130
 Gentianose, 32, 38
 Gentiobiose, 32, 38
 Gentiopierin, 54
 Glucase, 16, 35
 Glucodextrin, 21, 25

Gluconic acid, 80
 Glucose, 21, ff., 63
 Glucosides, 51-58; artificial, 57
 Glucosidogalactose, 39
 Glucovanillin, 54
 Glyceric acid, 67
 Glycogen, 27
 Glycolytic ferment, 89
Granulobacter butylicum, 101
 Grenzdextrin, 21, 25

Hadromal, 29
 Hadromase, 29
 Helicin, 53
 Hydroquinone, 83

Indican, 54
 Indigo, 87
 Indophenol, 87
 Indoxyl, 54
 Inosite, 63
 Inulase, 26
 Inulin, 26
 Invertin, 31, ff.
 Iron bacteria, 90
 Isatase, 54
 Isolactose, 39
 Isomaltose, 10, ff.

Kolanin, 57

Laccase, 82-86
 Laccol, 82, ff.
 Lactase, 36, 38, 39
 Lactic acid, 61-63, 68, ff.
 Lactose, 36
 Lecithines, 127
 Leucomaines, 140
 Lignin, 29
 Lipase, 36, 71-74
 Lotase, 57
 Lotusin, 56
 Luciferase, 89

Malic acid, 100
 Maltase, 16, 35-37, 52
 Maltodextrin, 7, 8, 14, 24
 Maltodextrinic acid, 14
 Maltose, 6-25, 35, 36, 52
 Mandelnitrilglucosid, 52
 Mannitol, 80
 Mannose, 28
 Mannosides, 58
 Melibiase, 32, 38
 Melibiose, 31, 38

Melitriose, 31
 Melizitose, 32
 Metamaltose, 13
 Metapectic acid, 28
Micrococcus prodigiosus, 103
 Milk, 110-113
 Molecular weight, starch, 19
 Monobutyryn, 72-74
 Monosaccharides, 40-66
 Mucic acid, 99
 Muscle-clotting, 111-113
Mycoderma aceti, 78
 Mycose, 38
 Myosin ferment, 112
 Myrosin, 55

Nitrification and denitrification,
 95-98
 Nitro-bacteria, 96
 Nitrosococci, 96
 Nitrosomonas, 96
 Nucleoproteids, 126

Enoxydase, 86
 Olein, 71
 Oxalic acid, 64
 Oxidation products, 78-90
 Oxydases, 82-89

Palmitin, 71
 Pancreatic digestion, 122-128
 Papain, 131
 Pectase, 29
 Pectic acid, 30
 Pectinase, 30
 Pectine, 29
 Pectose, 28
 Pectosic acid, 30
Penicillium brevicaulis, 93
Penicillium glaucum, 68, 70
 Pepsin, 114-121
 Peptic digestion, 114-121
 Peptone, 115
 Philothion, 91
 Phloridzin, 53
 Phosphorescent bacteria, 104
 Pigments due to bacterial action,
 102-104
 Polysaccharides, 5-30
 Populin, 54
 Propionic acid, 99, ff.
 Prostate gland, 113
 Protamines, 127
 Proteolytic fermentation, 114-133
 Ptomaines, 138

- Ptyalose, 23
 Putrefaction, 41, 134-140

 Raffinase, 32
 Raffinose, 31
 Reductase, 94
 Reduction products, 91-94
 Rennin, 110
 Rhamnase, 56
 Rhamnose, 56
 Rhamnin, 56
 Rhamninose, 56
 Rubian, 55

 Salicin, 53
 Salicylase, 88
 Schinoxydase, 86
Schizophyllum lobatum, 102
 Seminal vesicles, 113
 Seminase, 28
 Seminose, 29
 Sinalbin, 55
 Sinigrin, 55
 Sorbitol, 80
 Sorbose, 80
 Starch, 5-25, 71
 — molecule, 9, 19
Streptococcus hornensis, 65
Streptothrix chromogena, 102
 Sulphates, oxidation of, 90
 —, reduction of, 91, 92
 Sulphur bacteria, 90
 Synaptase, 51

 Synthetic action of ferments, 2, 35, 36

 Takadiastase, 36
 Tannase, 56
 Tannin, 56
 Touranose, 32
 Toxinea, 128, 139
 Trehalase, 38
 Trehalose, 38
 Trisaccharides, 31-39
 Trypsin, 114
 Tryptic digestion, 122-128
 Tyrosinase, 85

 Urea, 75-77
 Urease, 75
 Uric acid, 75-77
 Urushic acid, 82

 Valerian, 87
 Vanillin, 54
 Velocity of diastase hydrolysis, 23
 Vesiculase, 113
Vibrio hydrosulfureus, 92

 Xanthorhamnin, 56
 Xylose, 81
 Xylosides, 59

 Yeast ferment, 44-48, 130

 Zymase, 47

THE END.

A SELECT LIST OF BOOKS

IN

NATURAL AND PHYSICAL SCIENCE, MATHEMATICS AND TECHNOLOGY

PUBLISHED BY

MESSRS. LONGMANS, GREEN, & CO.

LONDON: 39 PATERNOSTER ROW, E.C.

NEW YORK: 91 & 93 FIFTH AVENUE.

BOMBAY: 32 HORNBY ROAD.

	PAGE		PAGE
<i>ADVANCED SCIENCE MANUALS</i>	- 38	MEDICINE AND SURGERY	- 25
ALGEBRA	- 9	MENSURATION	- 8
AGRICULTURE	- 35	METALLURGY	- 19
ARCHITECTURE	- 14	MINERALOGY	- 19
ASTRONOMY	- 20	MINING	- 19
BACTERIOLOGY	- 33	NATURAL HISTORY AND GENERAL	
BIOLOGY	- 32	SCIENCE	- 23
BOTANY	- 34	NAVAL ARCHITECTURE	- 19
BUILDING CONSTRUCTION	- 14	NAVIGATION	- 20
CALCULUS	- 10	OPTICS	- 12
CHEMISTRY	- 2	PHOTOGRAPHY	- 12
CONIC SECTIONS	- 9	PHYSICS	- 5
DYNAMICS	- 6	PHYSIOGRAPHY	- 22
ELECTRICITY	- 15	PHYSIOLOGY	- 32
<i>ELEMENTARY SCIENCE MANUALS</i>	- 38	<i>PRACTICAL ELEMENTARY SCIENCE</i>	
ENGINEERING	- 17	SERIES	- 40
EUCLID	- 10	<i>PROCTOR'S (R. A.) WORKS</i>	- 21
GARDENING	- 35	SOUND	- 13
GEOLOGY	- 22	STATICS	- 6
GEOMETRY	- 10	STEAM, OIL, AND GAS ENGINES	- 13
HEALTH AND HYGIENE	- 24	STRENGTH OF MATERIALS	- 17
HEAT	- 13	SURVEYING	- 8
HYDROSTATICS	- 6	TECHNOLOGY	- 23
LIGHT	- 13	TELEGRAPHY	- 16
LOGARITHMS	- 10	TELEPHONE	- 16
<i>LONDON SCIENCE CLASS-BOOKS</i>	- 40	<i>TEXT-BOOKS OF SCIENCE</i>	- 37
<i>LONGMANS' CIVIL ENGINEERING</i>		THERMODYNAMICS	- 13
SERIES	- 18	TRIGONOMETRY	- 12
MACHINE DRAWING AND DESIGN	- 18	<i>TYNDALL'S (JOHN) WORKS</i>	- 36
MAGNETISM	- 15	VETERINARY MEDICINE, ETC.	- 31
MANUFACTURES	- 23	WORKSHOP APPLIANCES	- 19
MECHANICS	- 6	ZOOLOGY	- 32

CHEMISTRY.

ARRHENIUS.—A TEXT-BOOK OF ELECTROCHEMISTRY. By SVANTE ARRHENIUS, Professor at the University of Stockholm. Translated from the German Edition by JOHN MCCRAE, Ph.D. With 58 Illustrations. 8vo., 9s. 6d. net.

CROOKES.—SELECT METHODS IN CHEMICAL ANALYSIS, chiefly Inorganic. By Sir WILLIAM CROOKES, F.R.S., etc. Third Edition, Rewritten and Enlarged. With 67 Woodcuts. 8vo., 21s. net.

FURNEAUX.—ELEMENTARY CHEMISTRY, Inorganic and Organic. By W. FURNEAUX, F.R.G.S., Lecturer on Chemistry, London School Board. With 65 Illustrations and 155 Experiments. Crown 8vo., 2s. 6d.

GARRETT AND HARDEN.—AN ELEMENTARY COURSE OF PRACTICAL ORGANIC CHEMISTRY. By F. C. GARRETT, M.Sc. (Vict. et Dunelm.), Assistant Lecturer and Demonstrator in Chemistry, the Durham College of Science, Newcastle-on-Tyne; and ARTHUR HARDEN, M.Sc. (Vict.), Ph.D., Assistant Lecturer and Demonstrator in Chemistry, the Owens College, Manchester. With 14 Illustrations. Crown 8vo., 2s.

JAGO.—Works by W. JAGO, F.C.S., F.I.C.

INORGANIC CHEMISTRY, THEORETICAL AND PRACTICAL. With an Introduction to the Principles of Chemical Analysis, Inorganic and Organic. With 63 Woodcuts and numerous Questions and Exercises. Fcp. 8vo., 2s. 6d.

AN INTRODUCTION TO PRACTICAL INORGANIC CHEMISTRY. Crown 8vo., 1s. 6d.

INORGANIC CHEMISTRY, THEORETICAL AND PRACTICAL. A Manual for Students in Advanced Classes of the Science and Art Department. With Plate of Spectra and 78 Woodcuts. Crown 8vo., 4s. 6d.

MELLOR.—HIGHER MATHEMATICS FOR STUDENTS OF CHEMISTRY AND PHYSICS. With Special Reference to Practical Work. By J. W. MELLOR, D.Sc., late Senior Scholar, and 1851 Exhibition Scholar, New Zealand University; Research Fellow, the Owens College, Manchester. With 142 Diagrams. 8vo., 12s. 6d. net.

MENDELÉEFF.—THE PRINCIPLES OF CHEMISTRY. By D. MENDELÉEFF. Translated from the Russian (Sixth Edition) by GEORGE KAMENSKY, A.R.S.M., of the Imperial Mint, St. Petersburg; and Edited by T. A. LAWSON, B.Sc., Ph.D., Fellow of the Institute of Chemistry. With 96 Diagrams and Illustrations. 2 vols. 8vo., 36s.

MEYER.—OUTLINES OF THEORETICAL CHEMISTRY. By LOTHAR MEYER, Professor of Chemistry in the University of Tübingen. Translated by Professors P. PHILLIPS BEDSON, D.Sc., and W. CARLETON WILLIAMS, B.Sc. 8vo., 9s.

MILLER.—INTRODUCTION TO THE STUDY OF INORGANIC CHEMISTRY. By W. ALLEN MILLER, M.D., LL.D. With 71 Illustrations. Fcp. 8vo., 3s. 6d.

CHEMISTRY—Continued.

MUIR.—A COURSE OF PRACTICAL CHEMISTRY. By M. M. P. MUIR, M.A., Fellow and Prælector in Chemistry of Gonville and Caius College, Cambridge. (3 Parts.)

Part I. Elementary. Crown 8vo., 4s. 6d.

Part II. Intermediate. Crown 8vo., 4s. 6d.

Part III. [In preparation.]

NEWTH.—Works by G. S. NEWTH, F.I.C. F.C.S., Demonstrator in the Royal College of Science, London.

CHEMICAL LECTURE EXPERIMENTS. With 230 Illustrations. Crown 8vo., 6s.

CHEMICAL ANALYSIS, QUANTITATIVE AND QUALITATIVE. With 100 Illustrations. Crown 8vo., 6s. 6d.

A TEXT-BOOK OF INORGANIC CHEMISTRY. With 155 Illustrations. Crown 8vo., 6s. 6d.

ELEMENTARY PRACTICAL CHEMISTRY. With 108 Illustrations and 254 Experiments. Crown 8vo., 2s. 6d.

OSTWALD.—SOLUTIONS. By W. OSTWALD, Professor of Chemistry in the University of Leipzig. Being the Fourth Book, with some additions, of the Second Edition of Oswald's 'Lehrbuch der allgemeinen Chemie'. Translated by M. M. PATTISON MUIR, Fellow and Prælector in Chemistry of Gonville and Caius College, Cambridge. 8vo., 10s. 6d.

PERKIN.—QUALITATIVE CHEMICAL ANALYSIS (ORGANIC AND INORGANIC). By F. MOLLOY PERKIN, Ph.D., Head of the Chemistry Department, Borough Polytechnic Institute, London. With 9 Illustrations and Spectrum Plate. 8vo., 3s. 6d.

REYNOLDS.—EXPERIMENTAL CHEMISTRY FOR JUNIOR STUDENTS. By J. EMERSON REYNOLDS, M.D., F.R.S., Professor of Chemistry, University of Dublin. Fcp. 8vo., with numerous Woodcuts.

Part I. Introductory. Fcp. 8vo., 1s. 6d.

Part II. Non-Metals, with an Appendix on Systematic Testing for Acids. Fcp. 8vo., 2s. 6d.

Part III. Metals, and Allied Bodies. Fcp. 8vo., 3s. 6d.

Part IV. Carbon Compounds. Fcp. 8vo., 4s.

SHENSTONE.—Works by W. A. SHENSTONE, F.R.S., Lecturer on Chemistry in Clifton College.

THE METHODS OF GLASS-BLOWING AND OF WORKING SILICA IN THE OXY-GAS FLAME. For the Use of Physical and Chemical Students. With 43 Illustrations. Crown 8vo., 2s. 6d.

A PRACTICAL INTRODUCTION TO CHEMISTRY. Intended to give a Practical acquaintance with the Elementary Facts and Principles of Chemistry. With 25 Illustrations. Crown 8vo., 2s.

CHEMISTRY—Continued.

SMITH AND HALL.—THE TEACHING OF CHEMISTRY AND PHYSICS IN THE SECONDARY SCHOOL. By ALEXANDER SMITH, B.Sc., Ph.D., Associate Professor of Chemistry in the University of Chicago, and EDWIN H. HALL, Ph.D., Professor of Physics in Harvard University. With 21 Woodcuts, Bibliographies, and Index. Crown 8vo., 6s. net.

THORNTON AND PEARSON.—NOTES ON VOLUMETRIC ANALYSIS. By ARTHUR THORNTON, M.A., and MARCHANT PEARSON, B.A., Assistant Science Master, Bradford Grammar School. Medium 8vo., 2s.

THORPE.—Works by T. E. THORPE, C.B., D.Sc. (Vict.), Ph.D., F.R.S., Principal of the Government Laboratory, London. Assisted by Eminent Contributors.

A DICTIONARY OF APPLIED CHEMISTRY. 3 vols. 8vo. Vols. I. and II., 42s. each. Vol. III., 63s.

QUANTITATIVE CHEMICAL ANALYSIS. With 88 Woodcuts. Fcp. 8vo., 4s. 6d.

THORPE AND MUIR.—QUALITATIVE CHEMICAL ANALYSIS AND LABORATORY PRACTICE. By T. E. THORPE, C.B., Ph.D., D.Sc., F.R.S., and M. M. PATTISON MUIR, M.A. With Plate of Spectra and 57 Illustrations. Fcp. 8vo., 3s. 6d.

TILDEN.—Works by WILLIAM A. TILDEN, D.Sc. London, F.R.S., Professor of Chemistry in the Royal College of Science, South Kensington.

A SHORT HISTORY OF THE PROGRESS OF SCIENTIFIC CHEMISTRY IN OUR OWN TIMES. Crown 8vo., 5s. net.

INTRODUCTION TO THE STUDY OF CHEMICAL PHILOSOPHY. The Principles of Theoretical and Systematic Chemistry. With 5 Illustrations. Fcp. 8vo., 5s. With ANSWERS to Problems. Fcp. 8vo., 5s. 6d.

PRACTICAL CHEMISTRY. The principles of Qualitative Analysis. Fcp. 8vo., 1s. 6d.

WATTS DICTIONARY OF CHEMISTRY. Revised and entirely Rewritten by H. FORSTER MORLEY, M.A., D.Sc., Fellow of, and lately Assistant Professor of Chemistry in, University College, London; and M. M. PATTISON MUIR, M.A., F.R.S.E., Fellow, and Prælector in Chemistry, of Gonville and Caius College, Cambridge. Assisted by Eminent Contributors. 4 vols. 8vo., £5 net.

WHITELEY.—Works by R. LLOYD WHITELEY, F.I.C., Principal of the Municipal Science School, West Bromwich.

CHEMICAL CALCULATIONS. With Explanatory Notes, Problems and Answers, specially adapted for use in Colleges and Science Schools. With a Preface by Professor F. CLOWES, D.Sc. (Lond.), F.I.C. Crown 8vo., 2s.

ORGANIC CHEMISTRY: the Fatty Compounds. With 45 Illustrations. Crown 8vo., 3s. 6d.

PHYSICS, ETC.

BIDGOOD.—ELEMENTARY PHYSICS AND CHEMISTRY FOR THE USE OF SCHOOLS. (In Three Books.) By JOHN BIDGOOD, B.Sc., Headmaster of the Gateshead School of Science.

Book I. Elementary Physics. With 120 Illustrations. Crown 8vo., 1s. 6d.

Book II. Physics and Chemistry. With 122 Illustrations. Crown 8vo., 1s. 6d.

BOSE.—RESPONSE IN THE LIVING AND NON-LIVING.

By JAGADIS CHUNDER BOSE, M.A. (Cantab.), D.Sc. (Lond.), Professor, Presidency College, Calcutta. With 117 Illustrations. 8vo., 10s. 6d.

**** This volume describes experimental investigations on animal, vegetable and inorganic substances regarding their response to stimulus. These researches show that the effects of fatigue, stimulants, depressants and poisons are alike in the organic and inorganic, and demonstrate that the response phenomena in the 'living' have been foreshadowed in the 'non-living'.*

GANOT.—Works by PROFESSOR GANOT. Translated and Edited by E. ATKINSON, Ph.D., F.C.S., and A. W. REINOLD, M.A., F.R.S.

ELEMENTARY TREATISE ON PHYSICS, Experimental and Applied. With 9 Coloured Plates and Maps, and 1048 Woodcuts, and Appendix of Problems and Examples with Answers. Crown 8vo., 15s.

NATURAL PHILOSOPHY FOR GENERAL READERS AND YOUNG PEOPLE. With 7 Plates, 632 Woodcuts, and an Appendix of Questions. Crown 8vo. 7s. 6d.

GLAZEBROOK AND SHAW.—PRACTICAL PHYSICS. By R. T. GLAZEBROOK, M.A., F.R.S., and W. N. SHAW, M.A. With 134 Illustrations. Fcp. 8vo., 7s. 6d.

GUTHRIE.—MOLECULAR PHYSICS AND SOUND. By F. GUTHRIE, Ph.D. With 91 Diagrams. Fcp. 8vo., 1s. 6d.

HELMHOLTZ.—POPULAR LECTURES ON SCIENTIFIC SUBJECTS. By HERMANN VON HELMHOLTZ. Translated by E. ATKINSON, Ph.D., F.C.S., formerly Professor of Experimental Science, Staff College. With 68 Illustrations. 2 vols., crown 8vo., 3s. 6d. each.

HENDERSON.—ELEMENTARY PHYSICS. By JOHN HENDERSON, D.Sc. (Edin.), A.I.E.E., Physics Department, Borough Road Polytechnic. Crown 8vo., 2s. 6d.

MACLEAN.—EXERCISES IN NATURAL PHILOSOPHY. By MAGNUS MACLEAN, D.Sc., Professor of Electrical Engineering at the Glasgow and West of Scotland Technical College. Crown 8vo., 4s. 6d.

MEYER.—THE KINETIC THEORY OF GASES. Elementary Treatise, with Mathematical Appendices. By Dr. OSKAR EMIL MEYER, Professor of Physics at the University of Breslau. Second Revised Edition. Translated by ROBERT E. BAYNES, M.A., Student of Christ Church, Oxford, and Dr. Lee's Reader in Physics. 8vo., 15s. net.

VAN 'T'HOFF.—THE ARRANGEMENT OF ATOMS IN SPACE. By J. H. VAN 'T'HOFF. Second, Revised, and Enlarged Edition. With a Preface by JOHANNES WISLICENUS, Professor of Chemistry at the University of Leipzig; and an Appendix 'Stereo-chemistry among Inorganic Substances,' by ALFRED WERNER, Professor of Chemistry at the University of Zürich. Translated and Edited by ARNOLD EILOART. Crown 8vo., 6s. 6d.

PHYSICS, ETC.—Continued.

WATSON.—Works by W. WATSON, A.R.C.S., F.R.S., D.Sc., Assistant Professor of Physics at the Royal College of Science, London.

ELEMENTARY PRACTICAL PHYSICS: a Laboratory Manual for Use in Organised Science Schools. With 120 Illustrations and 193 Exercises. Crown 8vo., 2s. 6d.

A TEXT-BOOK OF PHYSICS. With 564 Diagrams and Illustrations. Large crown 8vo., 10s. 6d.

WORTHINGTON.—A FIRST COURSE OF PHYSICAL LABORATORY PRACTICE. Containing 264 Experiments. By A. M. WORTHINGTON, M.A., F.R.S. With Illustrations. Crown 8vo., 4s. 6d.

WRIGHT.—ELEMENTARY PHYSICS. By MARK R. WRIGHT, M.A., Professor of Normal Education, Durham College of Science. With 242 Illustrations. Crown 8vo., 2s. 6d.

MECHANICS, DYNAMICS, STATICS, HYDRO-STATICS, ETC.

BALL.—A CLASS-BOOK OF MECHANICS. By Sir R. S. BALL, LL.D. 89 Diagrams. Fcp. 8vo., 1s. 6d.

GOODEVE.—Works by T. M. GOODEVE, M.A., formerly Professor of Mechanics at the Normal School of Science, and the Royal School of Mines.

THE ELEMENTS OF MECHANISM. With 357 Illustrations. Crown 8vo., 6s.

PRINCIPLES OF MECHANICS. With 253 Illustrations and numerous Examples. Crown 8vo., 6s.

A MANUAL OF MECHANICS: an Elementary Text-Book for Students of Applied Mechanics. With 138 Illustrations and Diagrams and 188 Examples taken from the Science Department Examination Papers, with Answers. Fcp. 8vo., 2s. 6d.

GOODMAN.—MECHANICS APPLIED TO ENGINEERING. By JOHN GOODMAN, Wh.Sch., A.M.I.C.E., M.I.M.E., Professor of Engineering in the Yorkshire College, Leeds (Victoria University). With 620 Illustrations and numerous examples. Crown 8vo., 7s. 6d. net.

GRIEVE.—LESSONS IN ELEMENTARY MECHANICS. By W. H. GRIEVE, late Engineer, R.N., Science Demonstrator for the London School Board, etc.

Stage 1. With 165 Illustrations and a large number of Examples. Fcp. 8vo., 1s. 6d.

Stage 2. With 122 Illustrations. Fcp. 8vo., 1s. 6d.

Stage 3. With 103 Illustrations. Fcp. 8vo., 1s. 6d.

MECHANICS, DYNAMICS, STATICS, HYDROSTATICS, ETC.—

Continued.

MAGNUS.—Works by SIR PHILIP MAGNUS, B.Sc., B.A.

LESSONS IN ELEMENTARY MECHANICS. Introductory to the study of Physical Science. Designed for the Use of Schools, and of Candidates for the London Matriculation and other Examinations. With numerous Exercises, Examples, Examination Questions, and Solutions, etc., from 1870-1895. With Answers, and 131 Woodcuts. Fcp. 8vo., 3s. 6d.

Key for the use of Teachers only, price 5s. 3½d.

HYDROSTATICS AND PNEUMATICS. Fcp. 8vo., 1s. 6d.; or, with Answers, 2s. The Worked Solutions of the Problems, 2s.

PULLEN.—MECHANICS: Theoretical, Applied, and Experimental. By W. W. F. PULLEN, WH. SC. M.I.M.E., A.M.I.C.E. With 318 Diagrams and numerous Examples. Crown 8vo., 4s. 6d.

ROBINSON.—ELEMENTS OF DYNAMICS (Kinetics and Statics). With numerous Exercises. A Text-book for Junior Students. By the Rev. J. L. ROBINSON, M.A. Crown 8vo., 6s.

SMITH.—Works by J. HAMBLIN SMITH, M.A.

ELEMENTARY STATICS. Crown 8vo., 3s.

ELEMENTARY HYDROSTATICS. Crown 8vo., 3s.

KEY TO STATICS AND HYDROSTATICS. Crown 8vo., 6s.

TARLETON.—AN INTRODUCTION TO THE MATHEMATICAL THEORY OF ATTRACTION. By FRANCIS A. TARLETON, LL.D., Sc.D., Fellow of Trinity College, and Professor of Natural Philosophy in the University of Dublin. Crown 8vo., 10s. 6d.

TAYLOR.—Works by J. E. TAYLOR, M.A., B.Sc. (Lond.).

THEORETICAL MECHANICS, including Hydrostatics and Pneumatics. With 175 Diagrams and Illustrations, and 522 Examination Questions and Answers. Crown 8vo., 2s. 6d.

THEORETICAL MECHANICS—SOLIDS. With 163 Illustrations, 120 Worked Examples and over 500 Examples from Examination Papers, etc. Crown 8vo., 2s. 6d.

THEORETICAL MECHANICS.—FLUIDS. With 122 Illustrations, numerous Worked Examples, and about 500 Examples from Examination Papers, etc. Crown 8vo., 2s. 6d.

THORNTON.—THEORETICAL MECHANICS—SOLIDS. Including Kinematics, Statics and Kinetics. By ARTHUR THORNTON, M.A., F.R.A.S. With 200 Illustrations, 130 Worked Examples, and over 900 Examples from Examination Papers, etc. Crown 8vo., 4s. 6d.

MECHANICS, DYNAMICS, STATICS, HYDROSTATICS, ETC.—

Continued.

TWISDEN.—Works by the Rev. JOHN F. TWISDEN, M.A.

PRACTICAL MECHANICS; an Elementary Introduction to their Study. With 855 Exercises, and 184 Figures and Diagrams. Crown 8vo., 10s. 6d.

THEORETICAL MECHANICS. With 172 Examples, numerous Exercises, and 154 Diagrams. Crown 8vo., 8s. 6d.

WILLIAMSON.—INTRODUCTION TO THE MATHEMATICAL THEORY OF THE STRESS AND STRAIN OF ELASTIC SOLIDS. By BENJAMIN WILLIAMSON, D.Sc., F.R.S. Crown 8vo., 5s.

WILLIAMSON AND TARLETON.—AN ELEMENTARY TREATISE ON DYNAMICS. Containing Applications to Thermodynamics, with numerous Examples. By BENJAMIN WILLIAMSON, D.Sc., F.R.S., and FRANCIS A. TARLETON, LL.D. Crown 8vo., 10s. 6d.

WORTHINGTON.—DYNAMICS OF ROTATION: an Elementary Introduction to Rigid Dynamics. By A. M. WORTHINGTON, M.A., F.R.S. Crown 8vo., 4s. 6d.

MENSURATION, SURVEYING, ETC.

BRABANT.—THE ELEMENTS OF PLANE AND SOLID MENSURATION. With Copious Examples and Answers. By F. G. BRABANT, M.A. Crown 8vo., 3s. 6d.

GRIBBLE.—PRELIMINARY SURVEY AND ESTIMATES. By THEODORE GRAHAM GRIBBLE, Civil Engineer. Including Elementary Astronomy, Route Surveying, Tacheometry, Curve Ranging, Graphic Mensuration, Estimates, Hydrography and Instruments. With 133 Illustrations, Quantity Diagrams, and a Manual of the Slide-Rule. Fcp. 8vo., 7s. 6d.

LODGE.—MENSURATION FOR SENIOR STUDENTS. By ALFRED LODGE, M.A., late Fereday Fellow of St. John's College, Oxford; Professor of Pure Mathematics at the Royal Indian Engineering College, Cooper's Hill. With Answers. Crown 8vo., 4s. 6d.

LUPTON.—A PRACTICAL TREATISE ON MINE SURVEYING. By ARNOLD LUPTON, Mining Engineer, Certificated Colliery Manager, Surveyor, Member of the Institution of Civil Engineers, etc. With 216 Illustrations. Medium 8vo., 12s. net.

NESBIT.—Works by A. NESBIT.

PRACTICAL MENSURATION. Illustrated by 700 Practical Examples and 700 Woodcuts. 12mo., 3s. 6d. KEY, 5s.

PRACTICAL LAND-SURVEYING, for the Use of Schools and Private Students. Edited by W. BURNES, F.R.A.S. With 14 Plates, 221 Figures, and a Field-Book. 8vo., 12s.

SMITH.—CIRCULAR SLIDE RULE. By G. L. SMITH. Fcp. 8vo., 1s. net.

ALGEBRA, ETC.

* * * For other Books, see Longmans & Co.'s Catalogue of Educational and School Books.

ANNALS OF MATHEMATICS. (*PUBLISHED UNDER THE AUSPICES OF HARVARD UNIVERSITY.*) Issued Quarterly. 4to., 2s. net.

BURNSIDE AND PANTON.—Works by WILLIAM SNOW BURNSIDE, M.A., Fellow of Trinity College, Dublin; and ARTHUR WILLIAM PANTON, M.A., Fellow and Tutor of Trinity College, Dublin.

THE THEORY OF EQUATIONS. With an Introduction to the Theory of Binary Algebraic Forms. 2 vols. 8vo., 9s. 6d. each.

AN INTRODUCTION TO DETERMINANTS: being a Chapter from the Theory of Equations (being the First Chapter of the Second Volume of 'The Theory of Equations'). 8vo., sewed, 2s. 6d.

CRACKNELL.—PRACTICAL MATHEMATICS. By A. G. CRACKNELL, M.A., B.Sc., Sixth Wrangler, etc. With Answers to the Examples. Crown 8vo., 3s. 6d.

GRIFFIN.—Works by Rev. WILLIAM NATHANIEL GRIFFIN, B.D., sometime Fellow of St. John's College, Cambridge.

THE ELEMENTS OF ALGEBRA AND TRIGONOMETRY. Fcp. 8vo., 3s. 6d.

NOTES ON THE ELEMENTS OF ALGEBRA AND TRIGONOMETRY. With Solutions of the more Difficult Questions. Fcp. 8vo., 3s. 6d.

MELLOR.—HIGHER MATHEMATICS FOR STUDENTS OF CHEMISTRY AND PHYSICS. With special reference to Practical Work. By J. W. MELLOR, D.Sc., Research Fellow, The Owens College, Manchester. With 142 Diagrams. 8vo., 12s. 6d. net.

WELSFORD AND MAYO.—ELEMENTARY ALGEBRA. By J. W. WELSFORD, M.A., formerly Fellow of Gonville and Caius College, Cambridge, and C. H. P. MAYO, M.A., formerly Scholar of St. Peter's College, Cambridge; Assistant Masters at Harrow School. Crown 8vo., 3s. 6d., or with Answers, 4s. 6d.

CONIC SECTIONS, ETC.

CASEY.—A TREATISE ON THE ANALYTICAL GEOMETRY OF THE POINT, LINE, CIRCLE, AND CONIC SECTIONS. By JOHN CASEY, LL.D., F.R.S. Crown 8vo., 12s.

RICHARDSON.—GEOMETRICAL CONIC SECTIONS. By G. RICHARDSON, M.A. Crown 8vo., 4s. 6d.

SALMON.—A TREATISE ON CONIC SECTIONS, containing an Account of some of the most Important Modern Algebraic and Geometric Methods. By G. SALMON, D.D., F.R.S. 8vo., 12s.

SMITH.—GEOMETRICAL CONIC SECTIONS. By J. HAMBLIN SMITH, M.A. Crown 8vo., 3s. 6d.

THE CALCULUS, LOGARITHMS, ETC.

BARKER.—GRAPHICAL CALCULUS. By ARTHUR H. BARKER, B.A., B.Sc. With an Introduction by JOHN GOODMAN, A.M.I.C.E. With 61 Diagrams. Crown 8vo., 4s. 6d.

MURRAY.—AN INTRODUCTORY COURSE IN DIFFERENTIAL EQUATIONS. By DANIEL ALEXANDER MURRAY, Ph.D. Crown 8vo., 4s. 6d.

TATE.—PRINCIPLES OF THE DIFFERENTIAL AND INTEGRAL CALCULUS. Applied to the Solution of Useful Problems in Mathematics and Mechanics. By THOMAS TATE. 12mo, 4s. 6d.

TAYLOR.—Works by F. GLANVILLE TAYLOR.

AN INTRODUCTION TO THE DIFFERENTIAL AND INTEGRAL CALCULUS AND DIFFERENTIAL EQUATIONS. Crown 8vo., 9s.

AN INTRODUCTION TO THE PRACTICAL USE OF LOGARITHMS, WITH EXAMPLES IN MENSURATION. With Answers to Exercises. Crown 8vo., 1s. 6d.

WILLIAMSON.—Works by BENJAMIN WILLIAMSON, D.Sc.

AN ELEMENTARY TREATISE ON THE DIFFERENTIAL CALCULUS; containing the Theory of Plane Curves, with numerous Examples. Crown 8vo., 10s. 6d.

AN ELEMENTARY TREATISE ON THE INTEGRAL CALCULUS; containing Applications to Plane Curves and Surfaces, and also a Chapter on the Calculus of Variations, with numerous Examples. Crown 8vo., 10s. 6d.

GEOMETRY AND EUCLID.

* * *For other Works, see Longmans & Co.'s Catalogue of Educational and School Books.*

ALLMAN.—GREEK GEOMETRY FROM THALES TO EUCLID. By G. J. ALLMAN. 8vo., 10s. 6d.

CASEY.—Works by JOHN CASEY, LL.D., F.R.S.

THE ELEMENTS OF EUCLID, BOOKS I.-VI. and Propositions, I.-XXI. of Book XI., and an Appendix of the Cylinder, Sphere, Cone, etc. With Copious Annotations and numerous Exercises. Fcp. 8vo., 4s. 6d. KEY to Exercises. Fcp. 8vo., 6s.

A SEQUEL TO THE ELEMENTS OF EUCLID. Part I. Books I.-VI. With numerous Examples. Fcp. 8vo., 3s. 6d.

A TREATISE ON THE ANALYTICAL GEOMETRY OF THE POINT, LINE, CIRCLE AND CONIC SECTIONS. Containing an Account of its most recent Extension. Crown 8vo., 12s.

GEOMETRY AND EUCLID—Continued.

HAMILTON.—ELEMENTS OF QUATERNIONS. By the late Sir WILLIAM ROWAN HAMILTON, LL.D., M.R.I.A. Edited by CHARLES JASPER JOLY, M.A., Fellow of Trinity College, Dublin. 2 vols. 4to. 21s. net each.

HIME.—THE OUTLINES OF QUATERNIONS. By Lieut.-Colonel H. W. L. HIME, late Royal Artillery. Crown 8vo., 10s.

LOW.—TEXT-BOOK ON PRACTICAL, SOLID, AND DESCRIPTIVE GEOMETRY. By DAVID ALLAN LOW, Professor of Engineering, East London Technical College. Crown 8vo.

Part I. With 114 Figures, 2s.

Part II. With 64 Figures, 3s.

MORRIS.—Works by I. HAMMOND MORRIS.

PRACTICAL PLANE AND SOLID GEOMETRY, including Graphic Arithmetic fully Illustrated with Drawings prepared specially by the Author. Crown 8vo., 2s. 6d.

GEOMETRICAL DRAWING FOR ART STUDENTS. Embracing Plane Geometry and its Applications, the Use of Scales, and the Plans and Elevations of Solids as required in Section I. of Science Subjects. Crown 8vo., 2s.

SMITH.—ELEMENTS OF GEOMETRY. By J. HAMBLIN SMITH, M.A. Containing Books 1 to 6, and portions of Books 11 and 12, of Euclid, with Exercises and Notes. Crown 8vo., 3s. 6d. KEY, crown 8vo., 8s. 6d.

Books 1 and 2, limp cloth, 1s. 6d., may be had separately.

SPOONER.—THE ELEMENTS OF GEOMETRICAL DRAWING: an Elementary Text-book on Practical Plane Geometry, including an Introduction to Solid Geometry. Written to include the requirements of the Syllabus of the Board of Education in Geometrical Drawing and for the use of Students preparing for the Military Entrance Examinations. By HENRY J. SPOONER, C.E., M.Inst.M.E.; Director of the Polytechnic School of Engineering, etc. Crown 8vo., 3s. 6d.

WATSON.—ELEMENTS OF PLANE AND SOLID GEOMETRY. By H. W. WATSON, M.A. Fcp. 8vo., 3s. 6d.

WILSON.—GEOMETRICAL DRAWING. For the use of Candidates for Army Examinations, and as an Introduction to Mechanical Drawing. By W. N. WILSON, M.A. Parts I. and II. Crown 8vo., 4s. 6d. each.

WINTER.—ELEMENTARY GEOMETRICAL DRAWING. By S. H. WINTER.

Part I. Including Practical Plane Geometry, the Construction of Scales, the Use of the Sector, the Marquois Scales, and the Protractor. With 3 Plates and 1000 Exercises and Examination Papers. Post 8vo., 5s.

TRIGONOMETRY.

CASEY.—A TREATISE ON ELEMENTARY TRIGONOMETRY. By JOHN CASEY, LL.D., F.R.S., late Fellow of the Royal University of Ireland. With numerous Examples and Questions for Examination. 12mo, 3s.

CLARKE.—PLANE TRIGONOMETRY. Containing the more advanced Propositions, Solution of Problems and a complete Summary of Formulæ, Bookwork, etc., together with recent Examination Papers for the Army, Woolwich, etc. With Answers. By the Rev. A. DAWSON CLARKE, M.A., St. John's College, Cambridge. Crown 8vo., 5s.

GOODWIN.—Works by H. B. GOODWIN, M.A.

PLANE AND SPHERICAL TRIGONOMETRY. In Three Parts, comprising those portions of the subjects, theoretical and practical, which are required in the Final Examination for Rank of Lieutenant at Greenwich. 8vo., 8s. 6d.

ELEMENTARY PLANE TRIGONOMETRY. With numerous Examples and Examination Papers set at the Royal Naval College in recent years. With Answers. 8vo., 5s.

JONES.—THE BEGINNINGS OF TRIGONOMETRY. By A. CLEMENT JONES, M.A., Ph.D., late Open Scholar and Senior Hulme Exhibitioner of Brasenose College, Oxford; Senior Mathematical Master of Bradford Grammar School. Crown 8vo., 2s.

MURRAY.—PLANE TRIGONOMETRY FOR COLLEGES AND SECONDARY SCHOOLS. By DANIEL A. MURRAY, B.A., Ph.D., Instructor in Mathematics in Cornell University. Crown 8vo., 3s. 6d. With Logarithmic and Trigonometric Tables. Crown 8vo., 5s.

SMITH.—ELEMENTARY TRIGONOMETRY. By J. HAMBLIN SMITH, M.A. Crown 8vo., 4s. 6d. Key, 7s. 6d.

OPTICS, PHOTOGRAPHY, ETC.

ABNEY.—A TREATISE ON PHOTOGRAPHY. By Sir WILLIAM DE WIVELESIE ABNEY, K.C.B., F.R.S., Principal Assistant Secretary of the Secondary Department of the Board of Education. With 134 Illustrations. Fcp. 8vo., 5s.

DRUDE.—THE THEORY OF OPTICS. By PAUL DRUDE, Professor of Physics at the University of Giessen. Translated from the German by C. RIBORG MANN and ROBERT A. MILLIKAN, Assistant Professors of Physics at the University of Chicago. With 110 Diagrams. 8vo., 15s. net.

GLAZEBROOK.—PHYSICAL OPTICS. By R. T. GLAZEBROOK, M.A., F.R.S., Principal of University College, Liverpool. With 183 Woodcuts of Apparatus, etc. Fcp. 8vo., 6s.

VANDERPOEL.—COLOR PROBLEMS: a Practical Manual for the Lay Student of Color. By EMILY NOYES VANDERPOEL. With 117 Plates in Color. Square 8vo., 21s. net.

WRIGHT.—OPTICAL PROJECTION: a Treatise on the Use of the Lantern in Exhibition and Scientific Demonstration. By LEWIS WRIGHT, Author of 'Light: a Course of Experimental Optics'. With 232 Illustrations. Crown 8vo., 6s.

SOUND, LIGHT, HEAT, AND THERMODYNAMICS.

DEXTER.—ELEMENTARY PRACTICAL SOUND, LIGHT AND HEAT. By JOSEPH S. DEXTER, B.Sc. (Lond.), Physics Master, Technical Day School, The Polytechnic Institute, Regent Street. With 152 Illustrations. Crown 8vo., 2s. 6d.

EMTAGE.—LIGHT. By W. T. A. EMTAGE, M.A., Director of Education, Mauritius. With 232 Illustrations. Crown 8vo., 6s.

HELMHOLTZ.—ON THE SENSATIONS OF TONE AS A PHYSIOLOGICAL BASIS FOR THE THEORY OF MUSIC. By HERMANN VON HELMHOLTZ. Royal 8vo., 28s.

MAXWELL.—THEORY OF HEAT. By J. CLERK MAXWELL, M.A., F.R.S.S., L. and E. With Corrections and Additions by Lord RAY LEIGH. With 38 Illustrations. Fcp. 8vo., 4s. 6d.

SMITH.—THE STUDY OF HEAT. By J. HAMBLIN SMITH, M.A., of Gonville and Caius College, Cambridge. Crown 8vo., 3s.

TYNDALL.—Works by JOHN TYNDALL, D.C.L., F.R.S.
See p. 36.

WORMELL.—A CLASS-BOOK OF THERMODYNAMICS. By RICHARD WORMELL, B.Sc., M.A. Fcp. 8vo., 1s. 6d.

WRIGHT.—Works by MARK R. WRIGHT, M.A.

SOUND, LIGHT, AND HEAT. With 160 Diagrams and Illustrations. Crown 8vo., 2s. 6d.

ADVANCED HEAT. With 136 Diagrams and numerous Examples and Examination Papers. Crown 8vo., 4s. 6d.

STEAM, OIL, AND GAS ENGINES.

BALE.—A HAND-BOOK FOR STEAM USERS; being Rules for Engine Drivers and Boiler Attendants, with Notes on Steam Engine and Boiler Management and Steam Boiler Explosions. By M. POWIS BALE, M.I.M.E., A.M.I.C.E. Fcp. 8vo., 2s. 6d.

CLERK.—THE GAS AND OIL ENGINE. By DUGALD CLERK, Member of the Institution of Civil Engineers, Fellow of the Chemical Society, Member of the Royal Institution, Fellow of the Institute of Patent Agents. With 228 Illustrations. 8vo., 15s.

STEAM, OIL, AND GAS ENGINES—Continued.

HOLMES.—THE STEAM ENGINE. By GEORGE C. V. HOLMES, Chairman of the Board of Works, Ireland. With 212 Illustrations. Fcp. 8vo., 6s.

NEILSON.—THE STEAM TURBINE. By ROBERT M. NEILSON, Whitworth Exhibitioner, Associate Member of the Institute of Mechanical Engineers, Lecturer on Steam and the Steam Engine at the Heginbottom Technical School, Ashton-under-Lyne. With 145 Illustrations. 8vo., 7s. 6d. net.

NORRIS.—A PRACTICAL TREATISE ON THE 'OTTO' CYCLE GAS ENGINE. By WILLIAM NORRIS, M.I.Mech.E. With 207 Illustrations. 8vo., 10s. 6d.

RIPPER.—Works by WILLIAM RIPPER, Professor of Mechanical Engineering in the Sheffield Technical School.

STEAM. With 185 Illustrations. Crown 8vo., 2s. 6d.

STEAM ENGINE THEORY AND PRACTICE. With 438 Illustrations. 8vo., 9s.

SENNETT AND ORAM.—THE MARINE STEAM ENGINE: A Treatise for Engineering Students, Young Engineers and Officers of the Royal Navy and Mercantile Marine. By the late RICHARD SENNETT, Engineer-in-Chief of the Navy, etc.; and HENRY J. ORAM, Senior Engineer Inspector at the Admiralty, Inspector of Machinery in H.M. Fleet, etc. With 414 Diagrams. 8vo., 21s.

STROMEYER.—MARINE BOILER MANAGEMENT AND CONSTRUCTION. Being a Treatise on Boiler Troubles and Repairs, Corrosion, Fuels, and Heat, on the properties of Iron and Steel, on Boiler Mechanics, Workshop Practices, and Boiler Design. By C. E. STROMEYER, Chief Engineer of the Manchester Steam Users' Association, Member of Council of the Institution of Naval Architects, etc. With 452 Diagrams, etc. 8vo., 12s. net.

ARCHITECTURE, BUILDING CONSTRUCTION, ETC.

ADVANCED BUILDING CONSTRUCTION. By the Author of 'Rivingtons' Notes on Building Construction'. With 385 Illustrations. Crown 8vo., 4s. 6d.

BURRELL.—BUILDING CONSTRUCTION. By EDWARD J. BURRELL, Second Master of the People's Palace Technical School, London. With 303 Working Drawings. Crown 8vo., 2s. 6d.

GWILT.—AN ENCYCLOPÆDIA OF ARCHITECTURE. By JOSEPH GWILT, F.S.A. Revised (1888), with Alterations and Considerable Additions by WYATT PAPWORTH. With 1700 Engravings. 8vo., 21s. net.

PARKER AND UNWIN.—THE ART OF BUILDING A HOME: A Collection of Lectures and Illustrations. By BARRY PARKER and RAYMOND UNWIN. With 68 Full-page Plates. 8vo., 10s. 6d. net.

RICHARDS.—BRICKLAYING AND BRICKCUTTING. By H. W. RICHARDS, Examiner in Brickwork and Masonry to the City and Guilds of London Institute, Head of Building Trades Department, Northern Polytechnic Institute, London, N. With over 200 Illustrations. 8vo., 3s. 6d.

ARCHITECTURE, BUILDING CONSTRUCTION, ETC.—Continued.

SEDDON.—**BUILDER'S WORK AND THE BUILDING TRADES.** By Col. H. C. SEDDON, R.E., late Superintending Engineer, H.M.'s Dockyard, Portsmouth; Examiner in Building Construction, Science and Art Department, South Kensington. With numerous Illustrations. Medium 8vo., 16s.

VALDER.—**BOOK OF TABLES**, giving the Cubic Contents of from One to Thirty Pieces Deals, Battens and Scantlings of the Sizes usually imported or used in the Building Trades, together with an Appendix showing a large number of sizes, the Contents of which may be found by referring to the aforesaid Tables. By THOMAS VALDER. Oblong 4to., 6s. net.

RIVINGTONS' COURSE OF BUILDING CONSTRUCTION.

NOTES ON BUILDING CONSTRUCTION. Arranged to meet the requirements of the syllabus of the Board of Education. Medium 8vo.

Part I. Elementary Stage. With 552 Illustrations, 9s. net.

Part II. Advanced Stage. With 479 Illustrations, 9s. net.

Part III. Materials. Course for Honours. With 188 Illustrations, 18s. net.

Part IV. Calculations for Building Structures. Course for Honours. With 551 Illustrations, 13s. net.

ELECTRICITY AND MAGNETISM.

ARRHENIUS.—**A TEXT-BOOK OF ELECTROCHEMISTRY.** By SVANTE ARRHENIUS, Professor at the University of Stockholm. Translated from the German Edition by JOHN McCRAE, Ph.D. With 58 Illustrations. 8vo., 9s. 6d. net.

CARUS-WILSON.—**ELECTRO-DYNAMICS**: the Direct-Current Motor. By CHARLES ASHLEY CARUS-WILSON, M.A. Cantab. With 71 Diagrams, and a Series of Problems, with Answers. Crown 8vo., 7s. 6d.

CUMMING.—**ELECTRICITY TREATED EXPERIMENTALLY.** By LINNÆUS CUMMING, M.A. With 242 Illustrations. Cr. 8vo., 4s. 6d.

DAY.—**EXERCISES IN ELECTRICAL AND MAGNETIC MEASUREMENTS**, with Answers. By R. E. DAY. 12mo., 3s. 6d.

FITZGERALD.—**THE SCIENTIFIC WRITINGS OF THE LATE GEORGE FRANCIS FITZGERALD**, Sc.D., F.R.S., F.R.S.E., Fellow of Trinity College, Dublin. Collected and Edited, with an Historical Introduction, by JOSEPH LARMOR, Sec.R.S., Fellow of St. John's College, Cambridge. With Portrait. 8vo., 15s.

GORE.—**THE ART OF ELECTRO-METALLURGY**, including all known Processes of Electro-Deposition. By G. GORE, LL.D., F.R.S. With 56 Illustrations. Fcp. 8vo., 6s.

HENDERSON.—**Works by JOHN HENDERSON**, D.Sc., F.R.S.E. **PRACTICAL ELECTRICITY AND MAGNETISM.** With 159 Illustrations and Diagrams. Crown 8vo., 6s. 6d.

PRELIMINARY PRACTICAL MAGNETISM AND ELECTRICITY. Crown 8vo., 1s.

ELECTRICITY AND MAGNETISM—Continued.

JENKIN.—ELECTRICITY AND MAGNETISM. By FLEEMING JENKIN, F.R.S., M.I.C.E. With 177 Illustrations. Fcp. 8vo., 3s. 6d.

JOUBERT.—ELEMENTARY TREATISE ON ELECTRICITY AND MAGNETISM. Founded on JOUBERT'S 'Traité Élémentaire d'Électricité'. By G. C. FOSTER, F.R.S., and E. ATKINSON, Ph.D. With Illustrations. Crown 8vo., 1s. [New Edition in the Press.]

JOYCE.—EXAMPLES IN ELECTRICAL ENGINEERING. By SAMUEL JOYCE, A.I.E.E. Crown 8vo., 5s.

MACLEAN AND MARCHANT.—ELEMENTARY QUESTIONS IN ELECTRICITY AND MAGNETISM. With Answers. Compiled by MAGNUS MACLEAN, D.Sc., M.I.E.E., and E. W. MARCHANT, D.Sc., A.I.E.E. Crown 8vo., 1s.

MERRIFIELD.—MAGNETISM AND DEVIATION OF THE COMPASS. By JOHN MERRIFIELD, LL.D., F.R.A.S., 18mo., 2s. 6d.

PARR.—PRACTICAL ELECTRICAL TESTING IN PHYSICS AND ELECTRICAL ENGINEERING. By G. D. ASPINALL PARR, Assoc. M.I.E.E. With 231 Illustrations. 8vo., 8s. 6d.

POYSER.—Works by A. W. POYSER, M.A.

MAGNETISM AND ELECTRICITY. With 235 Illustrations. Crown 8vo., 2s. 6d.

ADVANCED ELECTRICITY AND MAGNETISM. With 317 Illustrations. Crown 8vo., 4s. 6d.

RHODES.—AN ELEMENTARY TREATISE ON ALTERNATING CURRENTS. By W. G. RHODES, M.Sc. (Vict.), Consulting Engineer. With 80 Diagrams. 8vo., 7s. 6d. net.

SLINGO AND BROOKER.—Works by W. SLINGO and A. BROOKER.

ELECTRICAL ENGINEERING FOR ELECTRIC LIGHT ARTISANS AND STUDENTS. With 371 Illustrations. Crown 8vo., 12s.

PROBLEMS AND SOLUTIONS IN ELEMENTARY ELECTRICITY AND MAGNETISM. With 98 Illustrations. Cr. 8vo., 2s.

TYNDALL.—Works by JOHN TYNDALL, D.C.L., F.R.S. Seep. 36.

TELEGRAPHY AND THE TELEPHONE.

HOPKINS.—TELEPHONE LINES AND THEIR PROPERTIES. By WILLIAM J. HOPKINS, Professor of Physics in the Drexel Institute, Philadelphia. Crown 8vo., 6s.

PREECE AND SIVEWRIGHT.—TELEGRAPHY. By Sir W. H. PREECE, K.C.B., F.R.S., V.P.Inst., C.E., etc., Consulting Engineer and Electrician, Post Office Telegraphs; and Sir J. SIVEWRIGHT, K.C.M.G., General Manager, South African Telegraphs. With 267 Illustrations. Fcp. 8vo., 6s.

ENGINEERING, STRENGTH OF MATERIALS, ETC.

ANDERSON.—THE STRENGTH OF MATERIALS AND STRUCTURES: the Strength of Materials as depending on their Quality and as ascertained by Testing Apparatus. By Sir J. ANDERSON, C.E., LL.D., F.R.S.E. With 66 Illustrations. Fcp. 8vo., 3s. 6d.

BARRY.—RAILWAY APPLIANCES: a Description of Details of Railway Construction subsequent to the completion of the Earthworks and Structures. By Sir JOHN WOLFE BARRY, K.C.B., F.R.S., M.I.C.E. With 218 Illustrations. Fcp. 8vo., 4s. 6d.

DIPLOCK.—A NEW SYSTEM OF HEAVY GOODS TRANSPORT ON COMMON ROADS. By BRAHAM JOSEPH DIPLOCK. With 27 Illustrations. 8vo.

GOODMAN.—MECHANICS APPLIED TO ENGINEERING. By JOHN GOODMAN, Wh.Sch., A.M.I.C.E., M.I.M.E., Professor of Engineering in the Yorkshire College, Leeds (Victoria University). With 620 Illustrations and numerous Examples. Crown 8vo., 7s. 6d. net.

LOW.—A POCKET-BOOK FOR MECHANICAL ENGINEERS. By DAVID ALLAN LOW (Whitworth Scholar), M.I.Mech.E., Professor of Engineering, East London Technical College (People's Palace), London. With over 1000 specially prepared Illustrations. Fcp. 8vo., gilt edges, rounded corners, 7s. 6d.

PARKINSON.—LIGHT RAILWAY CONSTRUCTION. By RICHARD MARION PARKINSON, Assoc.M.Inst.C.E. With 85 Diagrams. 8vo., 10s. 6d. net.

SMITH.—GRAPHICS, or the Art of Calculation by Drawing Lines, applied especially to Mechanical Engineering. By ROBERT H. SMITH, Professor of Engineering, Mason College, Birmingham. Part I. With separate Atlas of 29 Plates containing 97 Diagrams. 8vo., 15s.

STONEY.—THE THEORY OF STRESSES IN GIRDERS AND SIMILAR STRUCTURES; with Practical Observations on the Strength and other Properties of Materials. By BINDON B. STONEY, LL.D., F.R.S., M.I.C.E. With 5 Plates and 143 Illust. in the Text. Royal 8vo., 36s.

UNWIN.—Works by W. CAWTHORNE UNWIN, F.R.S., B.Sc.
THE TESTING OF MATERIALS OF CONSTRUCTION.
A Text-book for the Engineering Laboratory and a Collection of the Results of Experiment. With 5 Plates and 188 Illustrations and Diagrams. 8vo., 16s. net.

ON THE DEVELOPMENT AND TRANSMISSION OF POWER FROM CENTRAL STATIONS: being the Howard Lectures delivered at the Society of Arts in 1893. With 81 Diagrams. 8vo., 10s. net.

WARREN.—ENGINEERING CONSTRUCTION IN IRON, STEEL, AND TIMBER. By WILLIAM HENRY WARREN, Challis Professor of Civil and Mechanical Engineering, University of Sydney. With 13 Folding Plates and 375 Diagrams. Royal 8vo., 16s. net.

WHEELER.—THE SEA COAST: Destruction, Littoral Drift, Protection. By W. H. WHEELER, M.Inst. C.E. With 38 Illustrations and Diagram. Medium 8vo., 10s. 6d. net.

LONGMANS' CIVIL ENGINEERING SERIES.

CIVIL ENGINEERING AS APPLIED TO CONSTRUCTION.

By LEVESON FRANCIS VERNON-HARCOURT, M.A., M.Inst.C.E. With 368 Illustrations. Medium 8vo., 14s. net.

CONTENTS.—Materials, Preliminary Works, Foundations and Roads—Railway Bridge and Tunnel Engineering—River and Canal Engineering—Irrigation Works—Dock Works and Maritime Engineering—Sanitary Engineering.

NOTES ON DOCKS AND DOCK CONSTRUCTION. By C.

COLSON, C.B., M.Inst.C.E. With 365 Illustrations. Medium 8vo., 21s. net.

CALCULATIONS IN HYDRAULIC ENGINEERING: a

Practical Text-Book for the use of Students, Draughtsmen and Engineers. By T. CLAXTON FIDLER, M.Inst.C.E.

Part I. Fluid Pressure and the Calculation of its Effects in Engineering Structures. With numerous Illustrations and Examples. 8vo., 6s. 6d. net.

Part II. Calculations in Hydro-Kinetics. With numerous Illustrations and Examples. 8vo., 7s. 6d. net.

RAILWAY CONSTRUCTION. By W. H. MILLS, M.I.C.E.,

Engineer-in-Chief of the Great Northern Railway of Ireland. With 516 Illustrations and Diagrams. 8vo., 18s. net.

PRINCIPLES AND PRACTICE OF HARBOUR CONSTRUCTION. By WILLIAM SHIELD, F.R.S.E., M.Inst.C.E. With 97 Illustrations. Medium 8vo., 15s. net.

TIDAL RIVERS: their (1) Hydraulics, (2) Improvement, (3)

Navigation. By W. H. WHEELER, M.Inst.C.E. With 75 Illustrations. Medium 8vo., 16s. net.

MACHINE DRAWING AND DESIGN.

LOW.—Works by DAVID ALLAN LOW, Professor of Engineering, East London Technical College (People's Palace).

IMPROVED DRAWING SCALES. 6d. in case.

AN INTRODUCTION TO MACHINE DRAWING AND DESIGN. With 153 Illustrations and Diagrams. Crown 8vo., 2s. 6d.

LOW AND BEVIS.—A MANUAL OF MACHINE DRAWING AND DESIGN. By DAVID ALLAN LOW and ALFRED WILLIAM BEVIS, M.I.Mech.E. With 700 Illustrations. 8vo., 7s. 6d.

UNWIN.—THE ELEMENTS OF MACHINE DESIGN. By W. CAWTHORNE UNWIN, F.R.S.

Part I. General Principles, Fastenings, and Transmissive Machinery. With 345 Diagrams, etc. Fcp. 8vo., 7s. 6d.

Part II. Chiefly on Engine Details. With 259 Illustrations. Fcp. 8vo., 6s.

NAVAL ARCHITECTURE.

- ATTWOOD.**—TEXT-BOOK OF THEORETICAL NAVAL ARCHITECTURE: a Manual for Students of Science Classes and Draughtsmen Engaged in Shipbuilders' and Naval Architects' Drawing Offices. By EDWARD LEWIS ATTWOOD, Assistant Constructor, Royal Navy. With 114 Diagrams. Crown 8vo., 7s. 6d.
- WATSON.**—NAVAL ARCHITECTURE: A Manual of Laying-off Iron, Steel and Composite Vessels. By THOMAS H. WATSON, Lecturer on Naval Architecture at the Durham College of Science, Newcastle-upon-Tyne. With numerous Illustrations. Royal 8vo., 15s. net.

WORKSHOP APPLIANCES, ETC.

- NORTHCOTT.**—LATHES AND TURNING, Simple, Mechanical and Ornamental. By W. H. NORTHCOTT. With 338 Illustrations. 8vo., 18s.
- SHELLEY.**—WORKSHOP APPLIANCES, including Descriptions of some of the Gauging and Measuring Instruments, Hand-cutting Tools, Lathes, Drilling, Planeing, and other Machine Tools used by Engineers. By C. P. B. SHELLEY, M.I.C.E. With an additional Chapter on Milling by R. R. LISTER. With 323 Illustrations. Fcp. 8vo., 5s.

MINERALOGY, MINING, METALLURGY, ETC.

- BAUERMAN.**—Works by HILARY BAUERMAN, F.G.S.
- SYSTEMATIC MINERALOGY.** With 373 Illustrations. Fcp. 8vo., 6s.
- DESCRIPTIVE MINERALOGY.** With 236 Illustrations. Fcp. 8vo., 6s.
- BREARLEY AND IBBOTSON.**—THE ANALYSIS OF STEEL-WORKS MATERIALS. By HARRY BREARLEY and FRED IBBOTSON, B.Sc. (Iond.), Demonstrator of Micrographic Analysis, University College, Sheffield. With 85 Illustrations. 8vo., 14s. net.
- GORE.**—THE ART OF ELECTRO-METALLURGY. By G. GORE, LL.D., F.R.S. With 56 Illustrations. Fcp. 8vo., 6s.
- HUNTINGTON AND M'MILLAN.**—METALS: their Properties and Treatment. By A. K. HUNTINGTON, Professor of Metallurgy in King's College, London, and W. G. M'MILLAN, Lecturer on Metallurgy in Mason's College, Birmingham. With 122 Illustrations. Fcp. 8vo., 7s. 6d.
- LUPTON.**—Works by ARNOLD LUPTON, M.I.C.E., F.G.S., etc.
- MINING.** An Elementary Treatise on the Getting of Minerals. With 596 Diagrams and Illustrations. Crown 8vo., 9s. net.
- A PRACTICAL TREATISE ON MINE SURVEYING.** With 209 Illustrations. 8vo., 12s. net.
- RHEAD.**—METALLURGY. By E. L. RHEAD, Lecturer on Metallurgy at the Municipal Technical School, Manchester. With 94 Illustrations. Fcp. 8vo., 3s. 6d.
- RHEAD AND SEXTON.**—ASSAYING AND METALLURGICAL ANALYSIS for the use of Students, Chemists and Assayers. By E. L. RHEAD, Lecturer on Metallurgy, Municipal School of Technology, Manchester; and A. HUMBOLDT SEXTON, F.I.C., F.C.S., Professor of Metallurgy, Glasgow and West of Scotland Technical College. 8vo., 10s. 6d. net.
- RUTLEY.**—THE STUDY OF ROCKS: an Elementary Text-book of Petrology. By F. RUTLEY, F.G.S. With 6 Plates and 88 other Illustrations. Fcp. 8vo., 4s. 6d.

ASTRONOMY, NAVIGATION, ETC.

ABBOTT.—ELEMENTARY THEORY OF THE TIDES: the Fundamental Theorems Demonstrated without Mathematics and the Influence on the Length of the Day Discussed. By T. K. ABBOTT, B.D., Fellow and Tutor, Trinity College, Dublin. Crown 8vo., 2s.

BALL.—Works by Sir ROBERT S. BALL, LL.D., F.R.S.

ELEMENTS OF ASTRONOMY. With 130 Figures and Diagrams. Fcp. 8vo., 6s. 6d.

A CLASS-BOOK OF ASTRONOMY. With 41 Diagrams. Fcp. 8vo., 1s. 6d.

GILL.—TEXT-BOOK ON NAVIGATION AND NAUTICAL ASTRONOMY. By J. GILL, F.R.A.S., late Head Master of the Liverpool Corporation Nautical College. 8vo., 10s. 6d.

GOODWIN.—AZIMUTH TABLES FOR THE HIGHER DECLINATIONS. (Limits of Declination 24° to 30° , both inclusive.) Between the Parallels of Latitude 0° and 60° . With Examples of the Use of the Tables in English and French. By H. B. GOODWIN, Naval Instructor, Royal Navy. Royal 8vo., 7s. 6d.

HERSCHEL.—OUTLINES OF ASTRONOMY. By Sir JOHN F. W. HERSCHEL, Bart., K.H., etc. With 9 Plates and numerous Diagrams. 8vo., 12s.

LAUGHTON.—AN INTRODUCTION TO THE PRACTICAL AND THEORETICAL STUDY OF NAUTICAL SURVEYING. By JOHN KNOX LAUGHTON, M.A., F.R.A.S. With 35 Diagrams. Crown 8vo., 6s.

LOWELL.—MARS. By PERCIVAL LOWELL, Fellow American Academy, Member Royal Asiatic Society, Great Britain and Ireland, etc. With 24 Plates. 8vo., 12s. 6d.

MARTIN.—NAVIGATION AND NAUTICAL ASTRONOMY. Compiled by Staff Commander W. R. MARTIN, R.N. Royal 8vo., 18s.

MERRIFIELD.—A TREATISE ON NAVIGATION. For the Use of Students. By J. MERRIFIELD, LL.D., F.R.A.S., F.M.S. With Charts and Diagrams. Crown 8vo., 5s.

PARKER.—ELEMENTS OF ASTRONOMY. With Numerous Examples and Examination Papers. By GEORGE W. PARKER, M.A., of Trinity College, Dublin. With 84 Diagrams. 8vo., 5s. 6d. net.

WEBB.—CELESTIAL OBJECTS FOR COMMON TELESCOPES. By the Rev. T. W. WEBB, M.A., F.R.A.S. Fifth Edition, Revised and greatly Enlarged by the Rev. T. E. ESPIN, M.A., F.R.A.S. (Two Volumes.) Vol. I., with Portrait and a Reminiscence of the Author, 2 Plates, and numerous Illustrations. Crown 8vo., 6s. Vol. II., with numerous Illustrations. Crown 8vo., 6s. 6d.

WORKS BY RICHARD A. PROCTOR.

THE MOON: Her Motions, Aspect, Scenery, and Physical Condition. With many Plates and Charts, Wood Engravings, and 2 Lunar Photographs. Crown 8vo., 3s. 6d.

OTHER WORLDS THAN OURS: the Plurality of Worlds Studied Under the Light of Recent Scientific Researches. With 14 Illustrations; Map, Charts, etc. Crown 8vo., 3s. 6d.

OUR PLACE AMONG INFINITIES: a Series of Essays contrasting our Little Abode in Space and Time with the Infinities around us. Crown 8vo., 3s. 6d.

MYTHS AND MARVELS OF ASTRONOMY. Crown 8vo., 3s. 6d.

LIGHT SCIENCE FOR LEISURE HOURS: Familiar Essays on Scientific Subjects, Natural Phenomena, etc. Crown 8vo., 3s. 6d.

THE ORBS AROUND US; Essays on the Moon and Planets, Meteors and Comets, the Sun and Coloured Pairs of Suns. Crown 8vo., 3s. 6d.

THE EXPANSE OF HEAVEN: Essays on the Wonders of the Firmament. Crown 8vo., 3s. 6d.

OTHER SUNS THAN OURS: a Series of Essays on Suns—Old, Young, and Dead. With other Science Gleanings. Two Essays on Whist, and Correspondence with Sir John Herschel. With 9 Star-Maps and Diagrams. Crown 8vo., 3s. 6d.

HALF-HOURS WITH THE TELESCOPE: a Popular Guide to the Use of the Telescope as a means of Amusement and Instruction. With 7 Plates. Fcp. 8vo., 2s. 6d.

NEW STAR ATLAS FOR THE LIBRARY, the School, and the Observatory, in Twelve Circular Maps (with Two Index-Plates). With an Introduction on the Study of the Stars. Illustrated by 9 Diagrams. Cr. 8vo., 5s.

THE SOUTHERN SKIES: a Plain and Easy Guide to the Constellations of the Southern Hemisphere. Showing in 12 Maps the position of the principal Star-Groups night after night throughout the year. With an Introduction and a separate Explanation of each Map. True for every Year. 4to., 5s.

HALF-HOURS WITH THE STARS: a Plain and Easy Guide to the Knowledge of the Constellations. Showing in 12 Maps the position of the principal Star-Groups night after night throughout the year. With Introduction and a separate Explanation of each Map. True for every Year. 4to., 3s. net.

LARGER STAR ATLAS FOR OBSERVERS AND STUDENTS.
In Twelve Circular Maps, showing 6000 Stars, 1500 Double Stars, Nebulæ, etc. With 2 Index-Plates. Folio, 15s.

WORKS BY RICHARD A. PROCTOR—Continued.

THE STARS IN THEIR SEASONS: an Easy Guide to a Knowledge of the Star-Groups. In 12 Large Maps. Imperial 8vo., 5s.

ROUGH WAYS MADE SMOOTH. Familiar Essays on Scientific Subjects. Crown 8vo., 3s. 6d.

PLEASANT WAYS IN SCIENCE. Crown 8vo., 3s. 6d.

NATURE STUDIES. By R. A. PROCTOR, GRANT ALLEN, A. WILSON, T. FOSTER, and E. CLODD. Crown 8vo., 3s. 6d.

LEISURE READINGS. By R. A. PROCTOR, E. CLODD, A. WILSON, T. FOSTER, and A. C. RANYARD. Crown 8vo., 3s. 6d.

PHYSIOGRAPHY AND GEOLOGY.

BIRD.—Works by CHARLES BIRD, B.A.

ELEMENTARY GEOLOGY. With Geological Map of the British Isles, and 247 Illustrations. Crown 8vo., 2s. 6d.

ADVANCED GEOLOGY. A Manual for Students in Advanced Classes and for General Readers. With over 300 Illustrations, a Geological Map of the British Isles (coloured), and a set of Questions for Examination. Crown 8vo., 7s. 6d.

GREEN.—**PHYSICAL GEOLOGY FOR STUDENTS AND GENERAL READERS.** By A. H. GREEN, M.A., F.G.S. With 236 Illustrations. 8vo., 21s.

MORGAN.—Works by ALEX. MORGAN, M.A., D.Sc., F.R.S.E.

ELEMENTARY PHYSIOGRAPHY. Treated Experimentally. With 4 Maps and 243 Diagrams. Crown 8vo., 2s. 6d.

ADVANCED PHYSIOGRAPHY. With 215 Illustrations. Crown 8vo., 4s. 6d.

THORNTON.—Works by J. THORNTON, M.A.

ELEMENTARY PRACTICAL PHYSIOGRAPHY.

Part I. With 215 Illustrations. Crown 8vo., 2s. 6d.

Part II. With 98 Illustrations. Crown 8vo., 2s. 6d.

ELEMENTARY PHYSIOGRAPHY: an Introduction to the Study of Nature. With 13 Maps and 295 Illustrations. With Appendix on Astronomical Instruments and Measurements. Crown 8vo., 2s. 6d.

ADVANCED PHYSIOGRAPHY. With 11 Maps and 255 Illustrations. Crown 8vo., 4s. 6d.

NATURAL HISTORY AND GENERAL SCIENCE.

BEDDARD.—THE STRUCTURE AND CLASSIFICATION OF BIRDS. By FRANK E. BEDDARD, M.A., F.R.S., Prosector and Vice-Secretary of the Zoological Society of London. With 252 Illus. 8vo., 21s. net.

FURNEAUX.—Works by WILLIAM FURNEAUX, F.R.G.S.

THE OUTDOOR WORLD; or, The Young Collector's Handbook. With 18 Plates, 16 of which are coloured, and 549 Illustrations in the Text. Crown 8vo., 6s. net.

LIFE IN PONDS AND STREAMS. With 8 Coloured Plates and 331 Illustrations in the Text. Crown 8vo., 6s. net.

BUTTERFLIES AND MOTHS (British). With 12 Coloured Plates and 241 Illustrations in the Text. Crown 8vo., 6s. net.

HUDSON.—BRITISH BIRDS. By W. H. HUDSON, C.M.Z.S. With 8 Coloured Plates from Original Drawings by A. THORBURN, and 8 Plates and 100 Figures by C. E. LODGE, and 3 Illustrations from Photographs. Crown 8vo., 6s. net.

MILLAIS.—THE NATURAL HISTORY OF THE BRITISH SURFACE-FEEDING DUCKS. By JOHN GUILLE MILLAIS, F.Z.S., etc. With 6 Photogravures and 66 Plates (41 in colours) from Drawings by the Author ARCHIBALD THORBURN, and from Photographs. Royal 4to., £6 6s. net.

NANSEN.—THE NORWEGIAN NORTH POLAR EXPEDITION, 1893-1896: Scientific Results. Edited by FRIDTJOF NANSEN. Volume I. With 44 Plates and numerous Illustrations in the Text. Demy 4to., 40s. net.

CONTENTS: The Fram—The Jurassic Fauna of Cape Flora. With a Geological Sketch of Cape Flora and its Neighbourhood—Fossil Plants from Franz Josef Land—An Account of the Birds—Crustacea.

Volume II. With 2 Charts and 17 Plates. Demy 4to., 30s. net.

CONTENTS: Astronomical Observations—Terrestrial Magnetism—Results of the Pendulum—Observations and some Remarks on the Constitution of the Earth's Crust.

Volume III. With 33 Plates. Demy 4to., 32s. net.

CONTENTS: The Oceanography of the North Polar Basin—On Hydrometers and the Surface Tension of Liquids.

STANLEY.—A FAMILIAR HISTORY OF BIRDS. By E. STANLEY, D.D., formerly Bishop of Norwich. With 160 Illustrations. Crown 8vo., 3s. 6d

MANUFACTURES, TECHNOLOGY, ETC.

BELL.—JACQUARD WEAVING AND DESIGNING. By F. T. BELL. With 199 Diagrams. 8vo., 12s. net.

CROSS AND BEVAN.—Works by C. F. CROSS and E. J. BEVAN.

CELLULOSE: an Outline of the Chemistry of the Structural Elements of Plants. With reference to their Natural History and Industrial Uses. (C. F. CROSS, E. J. BEVAN and C. BEADLE.) With 14 Plates. Crown 8vo., 12s. net.

RESEARCHES ON CELLULOSE, 1895-1900. Crown 8vo., 6s. net.

MANUFACTURES, TECHNOLOGY, ETC.—Continued.

MORRIS AND WILKINSON.—THE ELEMENTS OF COTTON SPINNING. By JOHN MORRIS and F. WILKINSON. With a Preface by Sir B. A. DOBSON, C.E., M.I.M.E. With 169 Diagrams and Illustrations. Crown 8vo., 7s. 6d. net.

RICHARDS.—BRICKLAYING AND BRICK-CUTTING. By H. W. RICHARDS, Examiner in Brickwork and Masonry to the City and Guilds of London Institute, Head of Building Trades Department, Northern Polytechnic Institute, London, N. With over 200 Illustrations. Med. 8vo., 3s. 6d.

TAYLOR.—COTTON WEAVING AND DESIGNING. By JOHN T. TAYLOR. With 373 Diagrams. Crown 8vo., 7s. 6d. net.

WATTS.—AN INTRODUCTORY MANUAL FOR SUGAR GROWERS. By FRANCIS WATTS, F.C.S., F.I.C. With 20 Illustrations. Crown 8vo., 6s.

HEALTH AND HYGIENE.

ASHBY.—HEALTH IN THE NURSERY. By HENRY ASHBY, M.D., F.R.C.P. With 25 Illustrations. Crown 8vo., 3s. net.

BUCKTON.—HEALTH IN THE HOUSE. By Mrs. C. M. BUCKTON. With 41 Woodcuts and Diagrams. Crown 8vo., 2s.

CORFIELD.—THE LAWS OF HEALTH. By W. H. CORFIELD, M.A., M.D. Fcp. 8vo., 1s. 6d.

FURNEAUX.—ELEMENTARY PRACTICAL HYGIENE.—Section I. By WILLIAM S. FURNEAUX. With 146 Illustrations. Cr. 8vo., 2s. 6d.

NOTTER AND FIRTH.—Works by J. L. NOTTER, M.A., M.D., and R. H. FIRTH, F.R.C.S.

HYGIENE. With 95 Illustrations. Crown 8vo., 3s. 6d.

PRACTICAL DOMESTIC HYGIENE. With 83 Illustrations. Crown 8vo., 2s. 6d.

POORE.—Works by GEORGE VIVIAN POORE, M.D.

ESSAYS ON RURAL HYGIENE. Crown 8vo., 6s. 6d.

THE DWELLING-HOUSE. With 36 Illustrations. Crown 8vo., 3s. 6d.

THE EARTH IN RELATION TO THE PRESERVATION AND DESTRUCTION OF CONTAGIA: being the Milroy Lectures delivered at the Royal College of Physicians in 1899, together with other Papers on Sanitation. With 13 Illustrations. Crown 8vo., 5s.

WILSON.—A MANUAL OF HEALTH-SCIENCE. By ANDREW WILSON, F.R.S.E., F.L.S., etc. With 74 Illustrations. Crown 8vo., 2s. 6d.

MEDICINE AND SURGERY.

ASHBY AND WRIGHT.—THE DISEASES OF CHILDREN, MEDICAL AND SURGICAL. By HENRY ASHBY, M.D., Lond., F.R.C.P., Physician to the General Hospital for Sick Children, Manchester; and G. A. WRIGHT, B.A., M.B. Oxon., F.R.C.S., Eng., Assistant-Surgeon to the Manchester Royal Infirmary, and Surgeon to the Children's Hospital. Enlarged and Improved Edition. With 192 Illustrations. 8vo., 25s.

BENNETT.—Works by SIR WILLIAM BENNETT, K.C.V.O., F.R.C.S., Surgeon to St. George's Hospital; Member of the Board of Examiners, Royal College of Surgeons of England.

CLINICAL LECTURES ON VARICOSE VEINS OF THE LOWER EXTREMITIES. With 3 Plates. 8vo., 6s.

ON VARICOCELE; A PRACTICAL TREATISE. With 4 Tables and a Diagram. 8vo., 5s.

CLINICAL LECTURES ON ABDOMINAL HERNIA: chiefly in relation to Treatment, including the Radical Cure. With 12 Diagrams in the Text. 8vo., 8s. 6d.

ON VARIX, ITS CAUSES AND TREATMENT, WITH ESPECIAL REFERENCE TO THROMBOSIS. 8vo., 3s. 6d.

THE PRESENT POSITION OF THE TREATMENT OF SIMPLE FRACTURES OF THE LIMBS. 8vo., 2s. 6d.

LECTURES ON THE USE OF MASSAGE AND EARLY PASSIVE MOVEMENTS IN RECENT FRACTURES AND OTHER COMMON SURGICAL INJURIES: The Treatment of Internal Derangements of the Knee Joint and Management of Stiff Joints. With 17 Illustrations. 8vo., 6s.

BENTLEY.—A TEXT-BOOK OF ORGANIC MATERIA MEDICA. Comprising a Description of the Vegetable and Animal Drugs of the British Pharmacopœia, with some others in common use. Arranged Systematically, and Especially Designed for Students. By ROBERT BENTLEY, M.R.C.S. Eng., F.L.S. With 62 Illustrations on Wood. Crown 8vo., 7s. 6d.

CABOT.—A GUIDE TO THE CLINICAL EXAMINATION OF THE BLOOD FOR DIAGNOSTIC PURPOSES. By RICHARD C. CABOT, M.D., Physician to Out-patients, Massachusetts General Hospital. With 3 Coloured Plates and 28 Illustrations in the Text. 8vo., 16s.

CARR, PICK, DORAN, AND DUNCAN.—THE PRACTITIONER'S GUIDE. By J. WALTER CARR, M.D. (Lond.), F.R.C.P.; T. PICKERING PICK, F.R.C.S.; ALBAN H. G. DORAN, F.R.C.S.; ANDREW DUNCAN, M.D., B.Sc. (Lond.), F.R.C.S., M.R.C.P. 8vo., 21s. net.

CELLI.—MALARIA, ACCORDING TO THE NEW RESEARCHES. By Prof. ANGELO CELLI, Director of the Institute of Hygiene, University of Rome. Translated from the Second Italian Edition by JOHN JOSEPH EYRE, M.R.C.P., L.R.C.S. Ireland, D.P.H. Cambridge. With an Introduction by Dr. PATRICK MANSON, Medical Adviser to the Colonial Office. 8vo., 10s. 6d.

MEDICINE AND SURGERY—Continued.

CHEYNE AND BURGHARD.—A MANUAL OF SURGICAL

TREATMENT. By W. WATSON CHEYNE, C.B. M.B., F.R.C.S., F.R.S., Professor of Surgery in King's College, London, Surgeon to King's College Hospital, etc.; and F. F. BURGHARD, M.D. and M.S., F.R.C.S., Teacher of Practical Surgery in King's College, London, Surgeon to King's College Hospital (Lond.), etc.

Part I. The Treatment of General Surgical Diseases, including Inflammation, Suppuration, Ulceration, Gangrene, Wounds and their Complications, Infective Diseases and Tumours; the Administration of Anæsthetics. With 66 Illustrations. Royal 8vo., 10s. 6d.

Part II. The Treatment of the Surgical Affections of the Tissues, including the Skin and Subcutaneous Tissues, the Nails, the Lymphatic Vessels and Glands, the Fasciæ, Bursæ, Muscles, Tendons and Tendon-sheaths, Nerves, Arteries and Veins. Deformities. With 141 Illustrations. Royal 8vo., 14s.

Part III. The Treatment of the Surgical Affections of the Bones. Amputations. With 100 Illustrations. Royal 8vo., 12s.

Part IV. The Treatment of the Surgical Affections of the Joints (including Excisions) and the Spine. With 138 Illustrations. Royal 8vo., 14s.

Part V. The Treatment of the Surgical Affections of the Head, Face, Jaws, Lips, Larynx and Trachea; and the Intrinsic Diseases of the Nose, Ear and Larynx, by H. LAMBERT LACK, M.D. (Lond.), F.R.C.S., Surgeon to the Hospital for Diseases of the Throat, Golden Square, and to the Throat and Ear Department, The Children's Hospital, Paddington Green. With 145 Illustrations. Royal 8vo., 18s.

Part VI. Section I. The Treatment of the Surgical Affections of the Tongue and Floor of the Mouth, the Pharynx, Neck, Esophagus, Stomach and Intestines. With 124 Illustrations. Royal 8vo., 18s.

Section II. The Treatment of the Surgical Affections of the Rectum, Liver, Spleen, Pancreas, Throat, Breast and Genito-urinary Organs. With Illustrations. Royal 8vo. [*In preparation.*]

CLARKE.—POST-MORTEM EXAMINATIONS IN MEDICO-LEGAL AND ORDINARY CASES. With Special Chapters on the Legal Aspects of Post-mortems, and on Certificates of Death. By J. JACKSON CLARKE, M.B. Lond., F.R.C.S., Assistant Surgeon at the North-west London and City Orthopædic Hospitals, etc. Fcp. 8vo., 2s. 6d.

COATS.—A MANUAL OF PATHOLOGY. By JOSEPH COATS, M.D., late Professor of Pathology in the University of Glasgow. Fourth Edition. Revised throughout and Edited by LEWIS R. SUTHERLAND, M.D., Professor of Pathology, University of St. Andrews. With 490 Illustrations. 8vo., 31s. 6d.

COOKE.—Works by THOMAS COOKE, F.R.C.S. Eng., B.A., B.Sc., M.D., Paris.

TABLETS OF ANATOMY. Being a Synopsis of Demonstrations given in the Westminster Hospital Medical School. Eleventh Edition in Three Parts, thoroughly brought up to date, and with over 700 Illustrations from all the best Sources, British and Foreign. Post 4to.

Part I. The Bones. 7s. 6d. net.

Part II. Limbs, Abdomen, Pelvis. 10s. 6d. net.

Part III. Head and Neck, Thorax, Brain. 10s. 6d. net.

MEDICINE AND SURGERY—Continued.

COOKE.—Works by THOMAS COOKE (*continued*).

APHORISMS IN APPLIED ANATOMY AND OPERATIVE SURGERY. Including 100 Typical *vivâ voce* Questions on Surface Marking, etc. Crown 8vo., 3s. 6d.

DAKIN.—A HANDBOOK OF MIDWIFERY. By WILLIAM RADFORD DAKIN, M.D., F.R.C.P., Obstetric Physician and Lecturer on Midwifery at St. George's Hospital, etc. With 394 Illustrations. Large crown 8vo., 18s.

DICKINSON.—Works by W. HOWSHIP DICKINSON, M.D. Cantab., F.R.C.P.

ON RENAL AND URINARY AFFECTIONS. With 12 Plates and 122 Woodcuts. Three Parts. 8vo., £3 4s. 6d.

THE TONGUE AS AN INDICATION OF DISEASE: being the Lumleian Lectures delivered March, 1888. 8vo., 7s. 6d.

OCCASIONAL PAPERS ON MEDICAL SUBJECTS, 1855-1896. 8vo., 12s.

MEDICINE OLD AND NEW. An Address Delivered on the Occasion of the Opening of the Winter Session, 1899-1900, at St. George's Hospital Medical School, on 2nd October, 1899. Crown 8vo., 2s. 6d.

DUCKWORTH.—Works by SIR DYCE DUCKWORTH, M.D., LL.D., Fellow and Treasurer of the Royal College of Physicians, etc.

THE SEQUELS OF DISEASE: being the Lumleian Lectures, 1896. 8vo., 10s. 6d.

THE INFLUENCE OF CHARACTER AND RIGHT JUDGMENT IN MEDICINE: the Harveian Oration, 1898. Post 4to. 2s. 6d.

ERICHSEN.—THE SCIENCE AND ART OF SURGERY; a Treatise on Surgical Injuries, Diseases, and Operations. By Sir JOHN ERIC ERICHSEN, Bart., F.R.S., LL.D. Edin., Hon. M.Ch. and F.R.C.S. Ireland. Illustrated by nearly 1000 Engravings on Wood. 2 vols. Royal 8vo., 48s.

FOWLER AND GODLEE.—THE DISEASES OF THE LUNGS. By JAMES KINGSTON FOWLER, M.A., M.D., F.R.C.P., Physician to the Middlesex Hospital and to the Hospital for Consumption and Diseases of the Chest, Brompton, etc.; and RICKMAN JOHN GODLEE, Honorary Surgeon in Ordinary to His Majesty, M.S., F.R.C.S., Fellow and Professor of Clinical Surgery, University College, London, etc. With 160 Illustrations. 8vo., 25s.

MEDICINE AND SURGERY—Continued.

GARROD.—Works by SIR ALFRED BARING GARROD, M.D., F.R.S., etc.

A TREATISE ON GOUT AND RHEUMATIC GOUT (RHEUMATOID ARTHRITIS). With 6 Plates, comprising 21 Figures (14 Coloured), and 27 Illustrations engraved on Wood. 8vo., 21s.

THE ESSENTIALS OF MATERIA MEDICA AND THERAPEUTICS. Crown 8vo., 12s. 6d.

GOADBY.—THE MYCOLOGY OF THE MOUTH : a Text-Book of Oral Bacteria. By KENNETH W. GOADBY, L.D.S. (Eng.), D.P.H. (Camb.), L.R.C.P., M.R.C.S., Bacteriologist and Lecturer on Bacteriology, National Dental Hospital, etc. With numerous Illustrations. 8vo.

GOODSALL AND MILES.—DISEASES OF THE ANUS AND RECTUM. By D. H. GOODSALL, F.R.C.S., Senior Surgeon, Metropolitan Hospital; Senior Surgeon, St. Mark's Hospital; and W. ERNEST MILES, F.R.C.S., Assistant Surgeon to the Cancer Hospital, Surgeon (out-patients), to the Gordon Hospital, etc. (In Two Parts.) Part I. With 91 Illustrations. 8vo., 7s. 6d. net.

GRAY.—ANATOMY, DESCRIPTIVE AND SURGICAL. By HENRY GRAY, F.R.S., late Lecturer on Anatomy at St. George's Hospital Medical School. The Fifteenth Edition Enlarged, edited by T. PICKERING PICK, F.R.C.S., Consulting Surgeon to St. George's Hospital, etc., and by ROBERT HOWDEN, M.A., M.B., C.M., Professor of Anatomy in the University of Durham, etc. With 772 Illustrations, a large proportion of which are Coloured, the Arteries being coloured red, the Veins blue, and the Nerves yellow. The attachments of the muscles to the bones, in the section on Osteology, are also shown in coloured outline. Royal 8vo., 32s. net.

HALLIBURTON.—Works by W. D. HALLIBURTON, M.D., F.R.S., Professor of Physiology in King's College, London.

A TEXT-BOOK OF CHEMICAL PHYSIOLOGY AND PATHOLOGY. With 104 Illustrations. 8vo., 28s.

ESSENTIALS OF CHEMICAL PHYSIOLOGY. With 77 Illustrations. 8vo., 5s.

LANG.—THE METHODICAL EXAMINATION OF THE EYE. Being Part I. of a Guide to the Practice of Ophthalmology for Students and Practitioners. By WILLIAM LANG, F.R.C.S. Eng., Surgeon to the Royal London Ophthalmic Hospital, Moorfields, etc. With 15 Illustrations. Crown 8vo., 3s. 6d.

LUFF.—TEXT-BOOK OF FORENSIC MEDICINE AND TOXICOLOGY. By ARTHUR P. LUFF, M.D., B.Sc. (Lond.), Physician-in-Charge of Out-Patients and Lecturer on Medical Jurisprudence and Toxicology in St. Mary's Hospital. With 13 full-page Plates (1 in colours) and 33 Illustrations in the Text. 2 vols. Crown 8vo., 24s.

MEDICINE AND SURGERY—Continued.

LIVERPOOL UNIVERSITY PRESS PUBLICATIONS, THE.

The Thomson Yates Laboratories Reports. Physiology; Pathology; Bacteriology; Tropical Medicine; Hygiene. Edited by RUPERT BOYCE and C. S. SHERRINGTON. With Plates and Illustrations in the text. Demy 4to. Vol I., 1898-9, 10s. 6d.; Vol. II., 1898-9, 25s.; Vol. III., Part I., 1900, 7s. 6d.; Vol. III., Part II., 1901, 12s. 6d.; Vol. IV., Part I., 1901, 20s.; Vol. IV., Part II., 1902, 21s.

THE LIVERPOOL SCHOOL OF TROPICAL MEDICINE MEMOIRS.

With Plates and Illustrations in the text. Demy 4to.

- I. Malarial Fever: Its Cause, Prevention and Treatment. Containing full details for the use of Travellers, Sportsmen, Soldiers, and Residents in Malarious Places. By RONALD ROSS, C.B., F.R.S., F.R.C.S. Ninth Edition, Revised and Enlarged. With Frontispiece. 8vo., 2s. 6d.
- II. Report of the Malaria Expedition to West Africa, August, 1899. By RONALD ROSS, C.B., F.R.S., F.R.C.S., H. E. ANNETT, M.D., D.P.H. and E. E. AUSTEN. With Supplementary Reports by Major G. M. GILES, M.B. and R. FIELDING-OULD, M.B. 21s.
- III. Report of the Malaria Expedition to Nigeria. Part I. Malarial Fever, etc. By H. E. ANNETT, M.D., J. EVERETT DUTTON, M.B. and J. H. ELLIOTT, M.D. 10s. 6d.
- IV. Report of the Malaria Expedition to Nigeria. Part II. Filariasis. By H. E. ANNETT, M.D., J. EVERETT DUTTON, M.B. and J. H. ELLIOTT, M.D. 15s.
- V. Part I. First Progress Report of the Campaign against Mosquitoes in Sierra Leone (1901). By RONALD ROSS, C.B., F.R.C.S., F.R.S. 8vo., 1s.
- VII. Report of the Yellow Fever Expedition to Pará (1900). By H. E. DURHAM, M.B., F.R.C.S., and the late WALTER MYERS, M.B. 4to., 7s. 6d.

MISCELLANEOUS.

Notes on Sanitary Conditions obtaining in Pará. By the MEMBERS OF THE YELLOW FEVER EXPEDITION. 8vo., 1s.

PAGET.—Edited by STEPHEN PAGET.

SELECTED ESSAYS AND ADDRESSES. By Sir JAMES PAGET. 8vo., 12s. 6d. net.

MEMOIRS AND LETTERS OF SIR JAMES PAGET, BART., F.R.S., D.C.L., late Sergeant-Surgeon to Her Majesty Queen Victoria. With 6 Portraits (3 Photogravures) and 4 other Illustrations. 8vo., 12s. 6d. net.

PICK.—SURGERY: a Treatise for Students and Practitioners. By T. PICKERING PICK, Consulting Surgeon to St. George's Hospital; Senior Surgeon to the Victoria Hospital for Children; H.M. Inspector of Anatomy in England and Wales. With 441 Illustrations. Medium 8vo., 25s.

POOLE.—COOKERY FOR THE DIABETIC. By W. H. and Mrs. POOLE. With Preface by Dr. PAVY. Fcap. 8vo., 2s. 6d.

PROBYN-WILLIAMS.—A PRACTICAL GUIDE TO THE ADMINISTRATION OF ANÆSTHETICS. By R. J. PROBYN-WILLIAMS, M.D., Anæsthetist and Instructor in Anæsthetics at the London Hospital; Lecturer in Anæsthetics at the London Hospital Medical College, etc. With 34 Illustrations. Crown 8vo., 4s. 6d. net.

MEDICINE AND SURGERY—Continued.

QUAIN.—**QUAIN'S (SIR RICHARD) DICTIONARY OF MEDICINE.** By Various Writers. Third Edition. Edited by H. MONTAGUE MURRAY, M.D., F.R.C.P., Joint Lecturer on Medicine, Charing Cross Medical School, and Physician to Out-Patients, Charing Cross Hospital; assisted by JOHN HAROLD, M.B., B.Ch., B.A.O., Physician to St. John's and St. Elizabeth's Hospital; and W. CECIL BOSANQUET, M.A., M.D., M.R.C.P., Physician to Out-Patients, Victoria Hospital for Children, Chelsea. With 21 Plates (14 in Colour) and numerous Illustrations in the Text. 8vo., 21s. net, buckram; or 30s. net, half-morocco.

QUAIN.—**QUAIN'S (JONES) ELEMENTS OF ANATOMY.** The Tenth Edition. Edited by EDWARD ALBERT SCHÄFER, F.R.S., Professor of Physiology in the University of Edinburgh; and GEORGE DANCER THANE, Professor of Anatomy in University College, London.

VOL. I., PART I. EMBRYOLOGY. By E. A. SCHÄFER, F.R.S. With 200 Illustrations. Royal 8vo., 9s.

VOL. I., PART II. GENERAL ANATOMY OR HISTOLOGY. By E. A. SCHÄFER, F.R.S. With 291 Illustrations. Royal 8vo., 12s. 6d.

VOL. II., PART I. OSTEOLOGY—ARTHROLOGY. By G. D. THANE. With 224 Illus. Royal 8vo., 11s.

VOL. II., PART II. MYOLOGY—ANGEIOLOGY. By G. D. THANE. With 199 Illustrations. Royal 8vo., 16s.

VOL. III., PART I. THE SPINAL CORD AND BRAIN. By E. A. SCHÄFER, F.R.S. With 139 Illustrations. Royal 8vo., 12s. 6d.

VOL. III., PART II. THE NERVES. By G. D. THANE. With 102 Illustrations. Royal 8vo., 9s.

VOL. III., PART III. THE ORGANS OF THE SENSES. By E. A. SCHÄFER, F.R.S. With 178 Illustrations. Royal 8vo., 9s.

VOL. III., PART IV. SPLANCHNOLOGY. By E. A. SCHÄFER, F.R.S., and JOHNSON SYMINGTON, M.D. With 337 Illustrations. Royal 8vo., 16s.

APPENDIX. SUPERFICIAL AND SURGICAL ANATOMY. By Professor G. D. THANE and Professor R. J. GODLEE, M.S. With 29 Illustrations. Royal 8vo., 6s. 6d.

SCHÄFER.—Works by E. A. SCHÄFER, F.R.S., Professor of Physiology in the University of Edinburgh.

THE ESSENTIALS OF HISTOLOGY. Descriptive and Practical. For the Use of Students. With 463 Illustrations. 8vo., 9s. net.

DIRECTIONS FOR CLASS WORK IN PRACTICAL PHYSIOLOGY: Elementary Physiology of Muscle and Nerve and of the Vascular and Nervous Systems. With 48 Diagrams and 24 pages of plain paper at end for Notes. 8vo., 3s. net.

SMALE AND COLYER.—**DISEASES AND INJURIES OF THE TEETH**, including Pathology and Treatment. By MORTON SMALE, M.R.C.S., L.S.A., L.D.S., Dental Surgeon to St. Mary's Hospital, Dean of the School, Dental Hospital of London, etc.; and J. F. COLYER, L.R.C.P., M.R.C.S., L.D.S., Dental Surgeon to Charing Cross Hospital and to the Dental Hospital of London. Second Edition Revised and Enlarged by J. F. COLYER. With 640 Illustrations. Large crown 8vo., 21s. net.

MEDICINE AND SURGERY—Continued.

SMITH (H. F.).—THE HANDBOOK FOR MIDWIVES. By HENRY FLY SMITH, B.A., M.B. Oxon., M.R.C.S. 41 Woodcuts. Cr. 8vo., 5s.

STEVENSON.—WOUNDS IN WAR: the Mechanism of their Production and their Treatment. By Surgeon-Colonel W. F. STEVENSON (Army Medical Staff), A.B., M.B., M.Ch. Dublin University, Professor of Military Surgery, Army Medical School, Netley. With 86 Illustrations. 8vo., 18s.

TAPPEINER. — INTRODUCTION TO CHEMICAL METHODS OF CLINICAL DIAGNOSIS. By Dr. H. TAPPEINER, Professor of Pharmacology and Principal of the Pharmacological Institute of the University of Munich. Translated by EDMOND J. MCWEENEY, M.A., M.D. (Royal Univ. of Ireland), L.R.C.P.I., etc. Crown 8vo., 3s. 6d.

WALLER.—Works by AUGUSTUS D. WALLER, M.D., Lecturer on Physiology at St. Mary's Hospital Medical School, London; late External Examiner at the Victorian University.

AN INTRODUCTION TO HUMAN PHYSIOLOGY. Third Edition, Revised. With 314 Illustrations. 8vo., 18s.

LECTURES ON PHYSIOLOGY. First Series. On Animal Electricity. 8vo., 5s. net.

VETERINARY MEDICINE, ETC.

FITZWYGRAM.—HORSES AND STABLES. By Lieut.-General Sir F. FITZWYGRAM, Bart. With 56 pages of Illustrations. 8vo., 3s. net.

STEEL.—Works by JOHN HENRY STEEL, F.R.C.V.S., F.Z.S., A.V.D., late Professor of Veterinary Science and Principal of Bombay Veterinary College.

A TREATISE ON THE DISEASES OF THE DOG; being a Manual of Canine Pathology. Especially adapted for the use of Veterinary Practitioners and Students. With 88 Illustrations. 8vo., 10s 6d.

A TREATISE ON THE DISEASES OF THE OX; being a Manual of Bovine Pathology. Especially adapted for the use of Veterinary Practitioners and Students. With 2 Plates and 117 Woodcuts. 8vo. 15s.

A TREATISE ON THE DISEASES OF THE SHEEP; being a Manual of Ovine Pathology for the use of Veterinary Practitioners and Students. With Coloured Plate and 99 Woodcuts. 8vo., 12s.

YOUATT.—Works by WILLIAM YOUATT.

THE HORSE. With 52 Wood Engravings. 8vo., 7s. 6d.

THE DOG. With 33 Wood Engravings. 8vo., 6s.

PHYSIOLOGY, BIOLOGY, ZOOLOGY, ETC.

(And see *MEDICINE AND SURGERY*, page 25.)

ASHBY.—NOTES ON PHYSIOLOGY FOR THE USE OF STUDENTS PREPARING FOR EXAMINATION. By HENRY ASHBY, M.D. Lond., F.R.C.P., Physician to the General Hospital for Sick Children, Manchester; formerly Demonstrator of Physiology, Liverpool School of Medicine. With 148 Illustrations. 18mo., 5s.

BARNETT.—THE MAKING OF THE BODY: a Children's Book on Anatomy and Physiology. By Mrs. S. A. BARNETT. With 113 Illustrations. Crown 8vo., 1s. 9d.

BEDDARD.—Works by FRANK E. BEDDARD, M.A. Oxon.
ELEMENTARY PRACTICAL ZOOLOGY. With 93 Illustrations. Crown 8vo., 2s. 6d.

THE STRUCTURE AND CLASSIFICATION OF BIRDS.
With 252 Illustrations. 8vo., 21s. net.

BIDGOOD.—A COURSE OF PRACTICAL ELEMENTARY BIOLOGY. By JOHN BIDGOOD, B.Sc., F.L.S. With 226 Illustrations. Crown 8vo., 4s. 6d.

BOSE.—RESPONSE IN THE LIVING AND NON-LIVING.
By JAGADIS CHUNDER BOSE, M.A. (Cantab.), D.Sc. (Lond.), Professor, Presidency College, Calcutta. With 117 Illustrations. 8vo., 10s. 6d.

* * This volume describes experimental investigations on animal, vegetable and inorganic substances regarding their response to stimulus. These researches show that the effects of fatigue, stimulants, depressants and poisons are alike in the organic and inorganic, and demonstrate that the response phenomena in the 'living' have been foreshadowed in the 'non-living'.

CHAPMAN.—THE FORAMINIFERA: An Introduction to the Study of the Protozoa. By FREDERICK CHAPMAN, A.L.S., F.R.M.S. With 14 Plates and 42 Illustrations in the Text. 8vo., 9s. net.

FURNEAUX.—HUMAN PHYSIOLOGY. By W. FURNEAUX, F.R.G.S. With 218 Illustrations. Crown 8vo., 2s. 6d.

HUDSON AND GOSSE.—THE ROTIFERA, or 'WHEEL-ANIMACULES'. By C. T. HUDSON, LL.D., and P. H. GOSSE, F.R.S. With 30 Coloured and 4 Uncoloured Plates. In 6 Parts. 4to., 10s. 6d. each. Supplement 12s. 6d. Complete in 2 vols., with Supplement, 4to., £4 4s.

MACALISTER.—Works by ALEXANDER MACALISTER, M.D.

AN INTRODUCTION TO THE SYSTEMATIC ZOOLOGY AND MORPHOLOGY OF VERTEBRATE ANIMALS. With 41 Diagrams. 8vo., 10s. 6d.

ZOOLOGY OF THE INVERTEBRATE ANIMALS. With 59 Diagrams. Fcp. 8vo., 1s. 6d.

ZOOLOGY OF THE VERTEBRATE ANIMALS. With 77 Diagrams. Fcp. 8vo., 1s. 6d.

PHYSIOLOGY, BIOLOGY, ZOOLOGY, ETC.—Continued.

MACDOUGAL.—Works by DANIEL TREMBLY MACDOUGALL, Ph.D., Director of the Laboratories of the New York Botanical Garden.

PRACTICAL TEXT-BOOK OF PLANT PHYSIOLOGY.

With 159 Illustrations. 8vo., 7s. 6d. net.

ELEMENTARY PLANT PHYSIOLOGY. With 108 Illustrations. Crown 8vo., 3s.

MOORE.—**ELEMENTARY PHYSIOLOGY.** By BENJAMIN MOORE, M.A., Lecturer on Physiology at the Charing Cross Hospital Medical School. With 125 Illustrations. Crown 8vo., 3s. 6d.

MORGAN.—**ANIMAL BIOLOGY:** an Elementary Text-Book. By C. LLOYD MORGAN, F.R.S., Principal of University College, Bristol. With 103 Illustrations. Crown 8vo., 8s. 6d.

SCHÄFER.—**DIRECTIONS FOR CLASS WORK IN PRACTICAL PHYSIOLOGY:** Elementary Physiology of Muscle and Nerve and of the Vascular and Nervous Systems. By E. A. SCHÄFER, LL.D., F.R.S., Professor of Physiology in the University of Edinburgh. With 48 Diagrams. 8vo., 3s. net.

THORNTON.—Works by JOHN THORNTON, M.A.

HUMAN PHYSIOLOGY. With 267 Illustrations, some Coloured. Crown 8vo., 6s.

ELEMENTARY BIOLOGY, Descriptive and Experimental. With numerous Illustrations. Crown 8vo., 3s. 6d.

BACTERIOLOGY.

CURTIS.—**THE ESSENTIALS OF PRACTICAL BACTERIOLOGY:** An Elementary Laboratory Book for Students and Practitioners. By H. J. CURTIS, B.S. and M.D. (Lond.), F.R.C.S. With 133 Illustrations. 8vo., 9s.

DHINGRA.—**AN INTRODUCTION TO BACTERIOLOGY.** (Specially designed for Indian Medical Students.) By M. L. DHINGRA, M.D.D.P.H.

FRANKLAND.—**MICRO-ORGANISMS IN WATER.** Together with an Account of the Bacteriological Methods involved in their Investigation. Specially designed for the use of those connected with the Sanitary Aspects of Water-Supply. By PERCY FRANKLAND, Ph.D., B.Sc. (Lond.), F.R.S., and Mrs. PERCY FRANKLAND. With 2 Plates and Numerous Diagrams. 8vo., 16s. net.

FRANKLAND.—**BACTERIA IN DAILY LIFE.** By Mrs. PERCY FRANKLAND, F.R.M.S. Crown 8vo.

GOADBY.—**THE MYCOLOGY OF THE MOUTH:** A Text-Book of Oral Bacteria. By KENNETH W. GOADBY, L.D.S. Eng., etc.; Bacteriologist and Lecturer on Bacteriology, National Dental Hospital, etc. With numerous Illustrations. 8vo.

BACTERIOLOGY—Continued.

KLÖCKER. — FERMENTATION ORGANISMS IN THE THEORY AND PRACTICE OF THE ALCOHOLIC FERMENTATION INDUSTRY, with special reference to the Construction and Work of Laboratories for the Physiology and Technology of Fermentation. By ALB. KLÖCKER. Translated by G. E. ALLAN, B.Sc., and J. H. MILLAR, F.I.C. With 146 Illustrations in the text. 8vo.

BOTANY.

AITKEN. — ELEMENTARY TEXT-BOOK OF BOTANY. By EDITH AITKEN, late Scholar of Girton College. With 400 Diagrams. Crown 8vo., 4s. 6d.

BENNETT AND MURRAY.—HANDBOOK OF CRYPTO-GAMIC BOTANY. By ALFRED W. BENNETT, M.A., B.Sc., F.L.S., Lecturer on Botany at St. Thomas's Hospital; and GEORGE MURRAY, F.L.S., Keeper of Botany, British Museum. With 378 Illustrations. 8vo., 16s.

CROSS AND BEVAN.—Works by C. F. CROSS, E. J. BEVAN and C. BEADLE.

CELLULOSE: an Outline of the Chemistry of the Structural Elements of Plants. With Reference to their Natural History and Industrial Uses. With 14 Plates. Crown 8vo., 12s. net.

RESEARCHES ON CELLULOSE, 1895-1900. Cr. 8vo., 6s. net.

EDMONDS.—Works by HENRY EDMONDS, B.Sc., London.

ELEMENTARY BOTANY. With 342 Illustrations. Cr. 8vo., 2s. 6d.

BOTANY FOR BEGINNERS. With 85 Illustrations. Fcp. 8vo., 1s. 6d.

FARMER.—A PRACTICAL INTRODUCTION TO THE STUDY OF BOTANY: Flowering Plants. By J. BRETLAND FARMER, F.R.S., M.A., Professor of Botany in the Royal College of Science, London. With 121 Illustrations. Crown 8vo., 2s. 6d.

KITCHENER.—A YEAR'S BOTANY. Adapted to Home and School Use. By FRANCES A. KITCHENER. With 195 Illustrations. Cr. 8vo., 5s.

LINDLEY AND MOORE.—THE TREASURY OF BOTANY. Edited by J. LINDLEY, M.D., F.R.S., and T. MOORE, F.L.S. With 20 Steel Plates and numerous Woodcuts. Two parts. Fcp. 8vo., 12s.

McNAB.—CLASS-BOOK OF BOTANY. By W. R. McNAB. MORPHOLOGY AND PHYSIOLOGY. With 42 Diagrams. CLASSIFICATION OF PLANTS. With 118 Diagrams. Fcp. 8vo., 1s. 6d.

SORAUER.—A POPULAR TREATISE ON THE PHYSIOLOGY OF PLANTS. By Dr. PAUL SORAUER. Translated by F. E. WEISS, B.Sc., F.L.S. With 33 Illustrations. 8vo., 9s. net.

BOTANY—Continued.

THOMÉ AND BENNETT.—STRUCTURAL AND PHYSIOLOGICAL BOTANY. By OTTO WILHELM THOMÉ and by ALFRED W. BENNETT, B.Sc., F.L.S. With Coloured Map and 600 Woodcuts. Fcp. 8vo., 6s.

TUBEUF.—DISEASES OF PLANTS INDUCED BY CRYPTOGAMIC PARASITES. Introduction to the Study of Pathogenic Fungi, Slime Fungi, Bacteria and Algæ. By Dr. KARL FREIHERR VON TUBEUF, Privatdocent in the University of Munich. English Edition by WILLIAM G. SMITH, B.Sc., Ph.D., Lecturer on Plant Physiology, University of Edinburgh. With 330 Illustrations. Royal 8vo., 18s. net.

WATTS.—A SCHOOL FLORA. For the use of Elementary Botanical Classes. By W. MARSHALL WATTS, D.Sc. Lond. Cr. 8vo., 2s. 6d.

AGRICULTURE AND GARDENING.

ADDYMAN.—AGRICULTURAL ANALYSIS. A Manual of Quantitative Analysis for Students of Agriculture. By FRANK T. ADDYMAN, B.Sc. (Lond.), F.I.C. With 49 Illustrations. Crown 8vo., 5s. net.

COLEMAN AND ADDYMAN.—PRACTICAL AGRICULTURAL CHEMISTRY. By J. BERNARD COLEMAN, A.R.C.Sc., F.I.C., and FRANK T. ADDYMAN, B.Sc. (Lond.), F.I.C. With 24 Illustrations. Crown 8vo., 1s. 6d. net.

HAGGARD.—Works by H. RIDER HAGGARD.

A FARMER'S YEAR: being his Commonplace Book for 1898. With 36 Illustrations by G. LEON LITTLE and three others. Crown 8vo., 7s. 6d. net.

RURAL ENGLAND. With Agricultural Maps and numerous Illustrations from Photographs. 2 vols. 8vo.

WEATHERS.—A PRACTICAL GUIDE TO GARDEN PLANTS. Containing Descriptions of the Hardest and most Beautiful Annuals and Biennials, Hardy Herbaceous and Bulbous Perennials, Hardy Water and Bog Plants, Flowering and Ornamental Trees and Shrubs, Conifers, Hardy Ferns, Hardy Bamboos and other Ornamental Grasses; and also the best kinds of Fruit and Vegetables that may be grown in the Open Air in the British Islands, with Full and Practical Instructions as to Culture and Propagation. By JOHN WEATHERS, F.R.H.S., late Assistant Secretary to the Royal Horticultural Society, formerly of the Royal Gardens, Kew, etc. With 163 Diagrams. 8vo., 21s. net.

WEBB.—Works by HENRY J. WEBB, Ph.D., B.Sc. (Lond.).

ELEMENTARY AGRICULTURE. A Text-Book specially adapted to the requirements of the Board of Education, the Junior Examination of the Royal Agricultural Society, and other Elementary Examinations. With 34 Illustrations. Crown 8vo., 2s. 6d.

AGRICULTURE. A Manual for Advanced Science Students. With 100 Illustrations. Crown 8vo., 7s. 6d. net.

WORKS BY JOHN TYNDALL, D.C.L., LL.D., F.R.S.

LECTURES ON SOUND. With Frontispiece of Fog-Syren, and 203 other Woodcuts and Diagrams in the Text. Crown 8vo., 10s. 6d.

HEAT, A MODE OF MOTION. With 125 Woodcuts and Diagrams. Crown 8vo., 12s.

LECTURES ON LIGHT DELIVERED IN THE UNITED STATES IN 1872 AND 1873. With Portrait, Lithographic Plate, and 59 Diagrams. Crown 8vo., 5s.

FRAGMENTS OF SCIENCE: a Series of Detached Essays, Addresses, and Reviews. 2 vols. Crown 8vo., 16s.

Vol. I.—The Constitution of Nature—Radiation—On Radiant Heat in Relation to the Colour and Chemical Constitution of Bodies—New Chemical Reactions produced by Light—On Dust and Disease—Voyage to Algeria to observe the Eclipse—Niagara—The Parallel Roads of Glen Roy—Alpine Sculpture—Recent Experiments on Fog-Signals—On the Study of Physics—On Crystalline and Slaty Cleavage—On Paramagnetic and Diamagnetic Forces—Physical Basis of Solar Chemistry—Elementary Magnetism—On Force—Contributions to Molecular Physics—Life and Letters of FARADAY—The Copley Medallist of 187c—The Copley Medallist of 1871—Death by Lightning—Science and the Spirits.

Vol. II.—Reflections on Prayer and Natural Law—Miracles and Special Providences—On Prayer as a Form of Physical Energy—Vitality—Matter and Force—Scientific Materialism—An Address to Students—Scientific Use of the Imagination—The Belfast Address—Apology for the Belfast Address—The Rev. JAMES MARTINEAU and the Belfast Address—Fermentation, and its Bearings on Surgery and Medicine—Spontaneous Generation—Science and Man—Professor VIRCHOW and Evolution—The Electric Light.

NEW FRAGMENTS. Crown 8vo., 10s. 6d.

CONTENTS.—The Sabbath—Goethe's 'Farbenlehre'—Atoms, Molecules, and Ether Waves—Count Rumford—Louis Pasteur, his Life and Labours—The Rainbow and its Congeners—Address delivered at the Birkbeck Institution on October 22, 1884—Thomas Young—Life in the Alps—About Common Water—Personal Recollections of Thomas Carlyle—On Unveiling the Statue of Thomas Carlyle—On the Origin, Propagation, and Prevention of Phthisis—Old Alpine Jottings—A Morning on Alp Lusen.

ESSAYS ON THE FLOATING MATTER OF THE AIR IN RELATION TO PUTREFACTION AND INFECTION. With 24 Woodcuts. Crown 8vo., 7s. 6d.

RESEARCHES ON DIAMAGNETISM AND MAGNETIC ACTION; including the Question of Diamagnetic Polarity. Crown 8vo., 12s.

NOTES OF A COURSE OF NINE LECTURES ON LIGHT, delivered at the Royal Institution of Great Britain, 1869. Crown 8vo., 1s. 6d.

NOTES OF A COURSE OF SEVEN LECTURES ON ELECTRICAL PHENOMENA AND THEORIES, delivered at the Royal Institution of Great Britain, 1870. Crown 8vo., 1s. 6d.

LESSONS IN ELECTRICITY AT THE ROYAL INSTITUTION 1875-1876. With 58 Woodcuts and Diagrams. Crown 8vo., 2s. 6d.

THE GLACIERS OF THE ALPS: being a Narrative of Excursions and Ascents. An Account of the Origin and Phenomena of Glaciers, and an Exposition of the Physical Principles to which they are related. With 7 Illustrations. Crown 8vo., 6s. 6d. net.

HOURS OF EXERCISE IN THE ALPS. With 7 Illustrations. Crown 8vo., 6s. 6d. net.

FARADAY AS A DISCOVERER. Crown 8vo., 3s. 6d.

TEXT-BOOKS OF SCIENCE.

- PHOTOGRAPHY.** By Sir WILLIAM DE WIVELESIE ABNEY, K.C.B., F.R.S. With 134 Illustrations. Fcp. 8vo., 5s.
- THE STRENGTH OF MATERIAL AND STRUCTURES.** By Sir J. ANDERSON, C.E. With 66 Illustrations. Fcp. 8vo., 3s. 6d.
- RAILWAY APPLIANCES.** By Sir JOHN WOLFE BARRY, K.C.B., F.R.S., M.I.C.E. With 218 Illustrations. Fcp. 8vo., 4s. 6d.
- INTRODUCTION TO THE STUDY OF INORGANIC CHEMISTRY.** By WILLIAM ALLEN MILLER, M.D., LL.D., F.R.S. With 72 Illustrations. 3s. 6d.
- QUANTITATIVE CHEMICAL ANALYSIS.** By T. E. THORPE, C.B., F.R.S., Ph.D. With 88 Illustrations. Fcp. 8vo., 4s. 6d.
- QUALITATIVE ANALYSIS AND LABORATORY PRACTICE.** By T. E. THORPE, C.B., Ph.D., F.R.S., and M. M. PATTISON MUIR, M.A. and F.R.S.E. With Plate of Spectra and 57 Illustrations. Fcp. 8vo., 3s. 6d.
- INTRODUCTION TO THE STUDY OF CHEMICAL PHILOSOPHY.** By WILLIAM A. TILDEN, D.Sc., London, F.R.S. With Illustrations. Fcp. 8vo., 5s. With Answers to Problems. Fcp. 8vo., 5s. 6d.
- ELEMENTS OF ASTRONOMY.** By Sir R. S. BALL, LL.D., F.R.S. With 130 Illustrations. Fcp. 8vo., 6s. 6d.
- SYSTEMATIC MINERALOGY.** By HILARY BAUERMAN, F.G.S. With 373 Illustrations. Fcp. 8vo., 6s.
- DESCRIPTIVE MINERALOGY.** By HILARY BAUERMAN, F.G.S., etc. With 236 Illustrations. Fcp. 8vo., 6s.
- METALS: THEIR PROPERTIES AND TREATMENT.** By A. K. HUNTINGTON and W. G. McMILLAN. With 122 Illustrations. Fcp. 8vo., 7s. 6d.
- THEORY OF HEAT.** By J. CLERK MAXWELL, M.A., LL.D., Edin., F.R.S.S., L. & E. With 38 Illustrations. Fcp. 8vo., 4s. 6d.
- PRACTICAL PHYSICS.** By R. T. GLAZEBROOK, M.A., F.R.S., and W. N. SHAW, M.A. With 134 Illustrations. Fcp. 8vo., 7s. 6d.
- PRELIMINARY SURVEY AND ESTIMATES.** By THEODORE GRAHAM GRIBBLE, Civil Engineer. Including Elementary Astronomy, Route Surveying, Tacheometry, Curve-ranging, Graphic Mensuration, Estimates, Hydrography and Instruments. With 133 Illustrations. Fcp. 8vo., 7s. 6d.
- ALGEBRA AND TRIGONOMETRY.** By WILLIAM NATHANIEL GRIFFIN, B.D. 3s. 6d. Notes on, with Solutions of the more difficult Questions. Fcp. 8vo., 3s. 6d.
- THE STEAM ENGINE.** By GEORGE C. V. HOLMES, Secretary of the Institution of Naval Architects. With 212 Illustrations. Fcp. 8vo., 6s.
- ELECTRICITY AND MAGNETISM.** By FLEEMING JENKIN, F.R.S.S., L. & E. With 177 Illustrations. Fcp. 8vo., 3s. 6d.
- THE ART OF ELECTRO-METALLURGY.** By G. GORE, LL.D., F.R.S. With 56 Illus. Fcp. 8vo., 6s.
- TELEGRAPHY.** By Sir W. H. PREECE, K.C.B., F.R.S., M.I.C.E., and Sir J. SIVEWRIGHT, M.A., K.C.M.G. With 267 Illustrations. Fcp. 8vo., 6s.
- PHYSICAL OPTICS.** By R. T. GLAZEBROOK, M.A., F.R.S. With 183 Illustrations. Fcp. 8vo., 6s.
- TECHNICAL ARITHMETIC AND MENSURATION.** By CHARLES W. MERRIEFIELD, F.R.S. 3s. 6d. Key, by the Rev. JOHN HUNTER, M.A. Fcp. 8vo., 3s. 6d.
- THE STUDY OF ROCKS.** By FRANK RUTLEY, F.G.S. With 6 Plates and 88 Illustrations. Fcp. 8vo., 4s. 6d.
- WORKSHOP APPLIANCES,** including Descriptions of some of the Machine Tools used by Engineers. By C. P. B. SHELLEY, M.I.C.E. With 323 Illustrations. Fcp. 8vo., 5s.
- ELEMENTS OF MACHINE DESIGN.** By W. CAWTHORNE UNWIN, F.R.S., B.Sc., M.I.C.E.
- PART I.** General Principles, Fastenings and Transmissive Machinery. With 345 Illustrations. Fcp. 8vo., 7s. 6d.
- PART II.** Chiefly on Engine Details. With 259 Illustrations. Fcp. 8vo., 6s.
- STRUCTURAL AND PHYSIOLOGICAL BOTANY.** By OTTO WILHELM THOMÉ, and A. W. BENNETT, M.A., B.Sc., F.L.S. With 600 Illustrations. Fcp. 8vo., 6s.
- PLANE AND SOLID GEOMETRY.** By H. W. WATSON, M.A. Fcp. 8vo., 3s. 6d.

ADVANCED SCIENCE MANUALS.

BUILDING CONSTRUCTION. By the Author of 'Rivington's Notes on Building Construction'. With 385 Illustrations and an Appendix of Examination Questions. Crown 8vo., 4s. 6d.

THEORETICAL MECHANICS. Solids, including Kinematics, Statics, and Kinetics. By A. THORNTON, M.A., F.R.A.S. With 220 Illustrations, 130 Worked Examples, and over 900 Examples from Examination Papers, etc. Crown 8vo., 4s. 6d.

HEAT. By MARK R. WRIGHT, Hon. Inter. B.Sc. (Lond.). With 136 Illustrations and numerous Examples and Examination Papers. Crown 8vo., 4s. 6d.

LIGHT. By W. J. A. EMTAGE, M.A. With 232 Illustrations. Cr. 8vo., 6s.

MAGNETISM AND ELECTRICITY. By ARTHUR WILLIAM POYSER, M.A. With 317 Illustrations. Crown 8vo., 4s. 6d.

INORGANIC CHEMISTRY, THEORETICAL AND PRACTICAL. By WILLIAM JAGO, F.C.S., F.I.C. With Plate of Spectra and 78 Woodcuts. Crown 8vo., 4s. 6d.

GEOLOGY: a Manual for Students in Advanced Classes and for General Readers. By CHARLES BIRD, B.A. (Lond.), F.G.S. With over 300 Illustrations, a Geological Map of the British Isles (coloured), and a set of Questions for Examination. Crown 8vo., 7s. 6d.

HUMAN PHYSIOLOGY: a Manual for Students in advanced Classes of the Science and Art Department. By JOHN THORNTON, M.A. With 268 Illustrations, some of which are Coloured, and a set of Questions for Examination. Crown 8vo., 6s.

PHYSIOGRAPHY. By JOHN THORNTON, M.A. With 11 Maps, 255 Illustrations, and Coloured Map of Ocean Deposits. Crown 8vo., 4s. 6d.

AGRICULTURE. By HENRY J. WEBB, Ph.D., B.Sc. With 100 Illustrations. Crown 8vo., 7s. 6d. net.

HYGIENE. By J. LANE NOTTER, M.A., M.D., Professor of Hygiene in the Army Medical School, Netley, Colonel, Royal Army Medical Corps; and R. H. FIRTH, F.R.C.S., late Assistant Professor of Hygiene in the Army Medical School, Netley, Major, Royal Army Medical Corps. With 95 Illustrations. Crown 8vo., 3s. 6d.

ELEMENTARY SCIENCE MANUALS.

* * *Written specially to meet the requirements of the ELEMENTARY STAGE OF SCIENCE SUBJECTS as laid down in the Syllabus of the Directory of the BOARD OF EDUCATION.*

PRACTICAL, PLANE, AND SOLID GEOMETRY, including Graphic Arithmetic. By I. H. MORRIS. Fully Illustrated with Drawings. Crown 8vo., 2s. 6d.

GEOMETRICAL DRAWING FOR ART STUDENTS. Embracing Plane Geometry and its Applications, the Use of Scales, and the Plans and Elevations of Solids. By I. H. MORRIS. Crown 8vo., 2s.

TEXT-BOOK ON PRACTICAL, SOLID, OR DESCRIPTIVE GEOMETRY. By DAVID ALLAN LOW (Whitworth Scholar). Part I. Crown 8vo., 2s. Part II. Crown 8vo., 3s.

AN INTRODUCTION TO MACHINE DRAWING AND DESIGN. By DAVID ALLAN LOW. With 153 Illustrations. Crown 8vo., 2s. 6d.

ELEMENTARY SCIENCE MANUALS—Continued.

- BUILDING CONSTRUCTION AND DRAWING.** By EDWARD J. BURRELL. With 308 Illustrations and Working Drawings. Crown 8vo., 2s. 6d.
- AN ELEMENTARY COURSE OF MATHEMATICS.** Containing Arithmetic; Euclid (Book I., with Deductions and Exercises); and Algebra. Crown 8vo., 2s. 6d.
- THEORETICAL MECHANICS.** Including Hydrostatics and Pneumatics. By J. E. TAYLOR, M.A., B.Sc. With numerous Examples and Answers, and 175 Diagrams and Illustrations. Crown 8vo., 2s. 6d.
- THEORETICAL MECHANICS—SOLIDS.** By J. E. TAYLOR, M.A., B.Sc. (Lond.). With 163 Illustrations, 120 Worked Examples, and over 500 Examples from Examination Papers, etc. Crown 8vo., 2s. 6d.
- THEORETICAL MECHANICS—FLUIDS.** By J. E. TAYLOR, M.A., B.Sc. (Lond.). With 122 Illustrations, numerous Worked Examples, and about 500 Examples from Examination Papers, etc. Crown 8vo., 2s. 6d.
- A MANUAL OF MECHANICS.** With 138 Illustrations and Diagrams, and 188 Examples taken from Examination Papers, with Answers. By T. M. GOODEVE, M.A. Crown 8vo., 2s. 6d.
- SOUND, LIGHT, AND HEAT.** By MARK R. WRIGHT, M.A. With 160 Diagrams and Illustrations. Crown 8vo., 2s. 6d.
- METALLURGY:** an Elementary Text-Book. By E. L. RHEAD. With 94 Illustrations. Crown 8vo., 3s. 6d.
- PHYSICS.** Alternative Course. By MARK R. WRIGHT, M.A. With 242 Illustrations. Crown 8vo., 2s. 6d.
- MAGNETISM AND ELECTRICITY.** By A. W. POYSER, M.A. With 235 Illustrations. Crown 8vo., 2s. 6d.
- PROBLEMS AND SOLUTIONS IN ELEMENTARY ELECTRICITY AND MAGNETISM.** By W. SLINGO and A. BROOKER. Embracing a Complete Set of Answers to the South Kensington Papers for the years 1885-1899, and a Series of Original Questions. With 67 Original Illustrations. Crown 8vo., 2s.
- ORGANIC CHEMISTRY:** the Fatty Compounds. By R. LLOYD WHITELEY, F.I.C., F.C.S. With 45 Illustrations. Crown 8vo., 3s. 6d.
- INORGANIC CHEMISTRY, THEORETICAL AND PRACTICAL.** By WILLIAM JAGO, F.C.S., F.I.C. With 63 Illustrations and numerous Questions and Exercises. Fep. 8vo. 2s. 6d.
- AN INTRODUCTION TO PRACTICAL INORGANIC CHEMISTRY.** By WILLIAM JAGO, F.C.S., F.I.C. Crown 8vo., 1s. 6d.
- PRACTICAL CHEMISTRY:** the Principles of Qualitative Analysis. By WILLIAM A. TILDEN, D.Sc. Fep. 8vo., 1s. 6d.
- ELEMENTARY INORGANIC CHEMISTRY.** By WILLIAM FURNEAUX, F.R.G.S. Crown 8vo., 2s. 6d.
- ELEMENTARY GEOLOGY.** By CHARLES BIRD, B.A., F.G.S. With Coloured Geological Map of the British Islands, and 247 Illustrations. Crown 8vo., 2s. 6d.
- HUMAN PHYSIOLOGY.** By WILLIAM FURNEAUX, F.R.G.S. With 218 Illustrations. Crown 8vo., 2s. 6d.
- A COURSE OF PRACTICAL ELEMENTARY BIOLOGY.** By J. BIDGOOD, B.Sc. With 226 Illustrations. Crown 8vo., 4s. 6d.
- ELEMENTARY BOTANY, THEORETICAL AND PRACTICAL.** By HENRY EDMONDS, B.Sc. With 342 Illustrations. Crown 8vo., 2s. 6d.
- STEAM.** By WILLIAM RIPPER, Member of the Institution of Mechanical Engineers. With 185 Illustrations. Crown 8vo., 2s. 6d.
- ELEMENTARY PHYSIOGRAPHY.** By J. THORNTON, M.A. With 13 Maps and 295 Illustrations. With Appendix on Astronomical Instruments and Measurements. Crown 8vo., 2s. 6d.
- AGRICULTURE.** By HENRY J. WEBB, Ph.D. With 34 Illustrations. Crown 8vo., 2s. 6d.

THE LONDON SCIENCE CLASS-BOOKS.

Edited by G. CAREY FOSTER, F.R.S., and by Sir PHILIP MAGNUS, B.Sc., B.A., of the City and Guilds of London Institute.

- ASTRONOMY.** By Sir ROBERT STAWELL BALL, LL.D., F.R.S. With 41 Diagrams. Fcp. 8vo., 1s. 6d.
- MECHANICS.** By Sir ROBERT STAWELL BALL, LL.D., F.R.S. With 89 Diagrams. Fcp. 8vo., 1s. 6d.
- THE LAWS OF HEALTH.** By W. H. CORFIELD, M.A., M.D., F.R.C.P. With 22 Illustrations. Fcp. 8vo., 1s. 6d.
- MOLECULAR PHYSICS AND SOUND.** By FREDERICK GUTHRIE, F.R.S. With 91 Diagrams. Fcp. 8vo., 1s. 6d.
- GEOMETRY, CONGRUENT FIGURES.** By O. HENRICI, Ph.D., F.R.S. With 141 Diagrams. Fcp. 8vo., 1s. 6d.
- ZOOLOGY OF THE INVERTEBRATE ANIMALS.** By ALEXANDER MACALISTER, M.D. With 59 Diagrams. Fcp. 8vo., 1s. 6d.
- ZOOLOGY OF THE VERTEBRATE ANIMALS.** By ALEXANDER MACALISTER, M.D. With 77 Diagrams. Fcp. 8vo., 1s. 6d.
- HYDROSTATICS AND PNEUMATICS.** By Sir PHILIP MAGNUS, B.Sc., B.A. With 79 Diagrams. Fcp. 8vo., 1s. 6d. (To be had also with *Answers*, 2s.) The Worked Solutions of the Problems. 2s.
- BOTANY.** Outlines of the Classification of Plants. By W. R. McNAB, M.D. With 118 Diagrams. Fcp. 8vo., 1s. 6d.
- BOTANY.** Outlines of Morphology and Physiology. By W. R. McNAB, M.D. With 42 Diagrams. Fcp. 8vo., 1s. 6d.
- THERMODYNAMICS.** By RICHARD WORMELL, M.A., D.Sc. With 41 Diagrams. Fcp. 8vo., 1s. 6d.

PRACTICAL ELEMENTARY SCIENCE SERIES.

- ELEMENTARY PRACTICAL PHYSIOGRAPHY.** (Section I.) By JOHN THORNTON, M.A. With 215 Illustrations and a Coloured Spectrum. Crown 8vo., 2s. 6d.
- ELEMENTARY PRACTICAL PHYSIOGRAPHY.** (Section II.) A Course of Lessons and Experiments in Elementary Science for the Queen's Scholarship Examination. By JOHN THORNTON, M.A. With 98 Illustrations and a Series of Questions. Crown 8vo., 2s. 6d.
- PRACTICAL DOMESTIC HYGIENE.** By J. LANE NOTTER, M.A., M.D., and R. H. FIRTH, F.R.C.S. With 83 Illustrations. Crown 8vo., 2s. 6d.
- A PRACTICAL INTRODUCTION TO THE STUDY OF BOTANY:** Flowering Plants. By J. BRETLAND FARMER, F.R.S., M.A. With 121 Illustrations. Crown 8vo., 2s. 6d.
- ELEMENTARY PRACTICAL HYGIENE.** Section I. By WILLIAM S. FURNEAUX. With 146 Illustrations. Crown 8vo., 2s. 6d.
- ELEMENTARY PRACTICAL SOUND, LIGHT, AND HEAT.** By JOSEPH S. DEXTER. With 152 Illustrations. Crown 8vo., 2s. 6d.
- PRACTICAL MATHEMATICS.** By A. G. CRACKNELL, M.A., B.Sc. Crown 8vo., 3s. 6d.
- ELEMENTARY PRACTICAL CHEMISTRY.** By G. S. NEWTH, F.I.C., F.C.S. With 108 Illustrations and 254 Experiments. Crown 8vo., 2s. 6d.
- ELEMENTARY PRACTICAL PHYSICS.** By W. WATSON, B.Sc. With 120 Illustrations and 193 Exercises. Crown 8vo., 2s. 6d.
- ELEMENTARY PRACTICAL ZOOLOGY.** By FRANK E. BEDDARD, M.A. Oxon., F.R.S. With 93 Illustrations. Crown 8vo., 2s. 6d.
- ELEMENTARY BIOLOGY.** By JOHN THORNTON, M.A. With 108 Illustrations. Crown 8vo., 3s. 6d.
- THE ELEMENTS OF GEOMETRICAL DRAWING:** an Elementary Text-book on Practical Plane Geometry, including an Introduction to Solid Geometry. By HENRY J. SPOONER, C.E., M.Inst.M.E. Crown 8vo., 3s. 6d.

MPHy
P

99429

Plimmer, Robert Henry Aders

The chemical changes and products resulting from
fermentations.

**University of Toronto
Library**

**DO NOT
REMOVE
THE
CARD
FROM
THIS
POCKET**

Acme Library Card Pocket
LOWE-MARTIN CO. LIMITED

